

# Analytic Properties of Shintani Zeta Functions

Frank Thorne

Stanford University and the University of South Carolina

January 19, 2010

# A famous problem

What is the distribution of field discriminants?

# A famous problem

What is the distribution of field discriminants?

## Theorem

*There are only finitely many fields  $K$  with discriminant  $|\Delta_K| < X$ .*

# A famous problem

What is the distribution of field discriminants?

## Theorem

*There are only finitely many fields  $K$  with discriminant  $|\Delta_K| < X$ .*

Some questions:

# A famous problem

What is the distribution of field discriminants?

## Theorem

*There are only finitely many fields  $K$  with discriminant  $|\Delta_K| < X$ .*

Some questions:

- ▶ How many  $K$  are there with fixed degree  $d$  and  $|\Delta_K| < X$ ?

# A famous problem

What is the distribution of field discriminants?

## Theorem

*There are only finitely many fields  $K$  with discriminant  $|\Delta_K| < X$ .*

Some questions:

- ▶ How many  $K$  are there with fixed degree  $d$  and  $|\Delta_K| < X$ ?
- ▶ How many with fixed degree and  $|\Delta_K| = X$ ?

# Asymptotics for cubic fields

**Folk Conjecture:** For each  $d$ ,

$$n_d(X) \sim C_d X,$$

for some constant  $C_d$ .

# Counting fields of low degree

Results:



# Counting fields of low degree

Results:

►  $d = 2$ :  $C_2 = 6/\pi^2$ .

# Counting fields of low degree

Results:

- ▶  $d = 2$ :  $C_2 = 6/\pi^2$ .
- ▶  $d = 3$ : (Davenport-Heilbronn)  $C_3 = \frac{1}{3\zeta(3)}$ .

# Counting fields of low degree

Results:

- ▶  $d = 2$ :  $C_2 = 6/\pi^2$ .
- ▶  $d = 3$ : (Davenport-Heilbronn)  $C_3 = \frac{1}{3\zeta(3)}$ .
- ▶  $d = 4$ : (Bhargava)  $C_4 = \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})$ .

# Counting fields of low degree

Results:

- ▶  $d = 2$ :  $C_2 = 6/\pi^2$ .
- ▶  $d = 3$ : (Davenport-Heilbronn)  $C_3 = \frac{1}{3\zeta(3)}$ .
- ▶  $d = 4$ : (Bhargava)  $C_4 = \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})$ .
- ▶  $d = 5$ : (Bhargava)  $C_5 = \frac{13}{120} \prod_p (1 + p^{-2} - p^{-4} - p^{-5})$ .

# Counting fields of low degree

Results:

- ▶  $d = 2$ :  $C_2 = 6/\pi^2$ .
- ▶  $d = 3$ : (Davenport-Heilbronn)  $C_3 = \frac{1}{3\zeta(3)}$ .
- ▶  $d = 4$ : (Bhargava)  $C_4 = \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})$ .
- ▶  $d = 5$ : (Bhargava)  $C_5 = \frac{13}{120} \prod_p (1 + p^{-2} - p^{-4} - p^{-5})$ .
- ▶  $d > 5$ : Open.  $C_d$  conjectured by Bhargava;  $n_d(X) \ll X^{n^\epsilon}$  due to Ellenberg-Venkatesh.

# From cubic fields to cubic forms

To count cubic fields:

# From cubic fields to cubic forms

To count cubic fields:

- ▶ First count cubic *rings* and then apply a sieve method;

# From cubic fields to cubic forms

To count cubic fields:

- ▶ First count cubic *rings* and then apply a sieve method;
- ▶ Understand cubic rings by means of binary cubic forms.



# Binary cubic forms

Let  $V$  be the space of binary cubic forms:

$$V := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 u v^2 + x_4 v^3; \ x_1, x_2, x_3, x_4 \in \mathbb{R}\}.$$

# Binary cubic forms

Let  $V$  be the space of binary cubic forms:

$$V := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 u v^2 + x_4 v^3; \ x_1, x_2, x_3, x_4 \in \mathbb{R}\}.$$

$\mathrm{GL}_2(\mathbb{R})$  acts on  $V$  by

$$(gx)(u, v) = x((u, v)g).$$

# Binary cubic forms

Let  $V$  be the space of binary cubic forms:

$$V := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 u v^2 + x_4 v^3; \ x_1, x_2, x_3, x_4 \in \mathbb{R}\}.$$

$\mathrm{GL}_2(\mathbb{R})$  acts on  $V$  by

$$(gx)(u, v) = x((u, v)g).$$

This makes  $V$  into a **prehomogeneous vector space**: There are  $\mathrm{GL}_2$ -orbits of the same dimension as  $V$ .

# The parameterization of cubic rings

## Definition

A **cubic ring** is a ring which is free of rank 3 as a  $\mathbb{Z}$ -module.

# The parameterization of cubic rings

## Definition

A **cubic ring** is a ring which is free of rank 3 as a  $\mathbb{Z}$ -module.

Nondegenerate cubic rings are: Orders in cubic fields;  $Q \oplus n\mathbb{Z}$  for  $Q$  a quadratic order,  $n_1\mathbb{Z} \oplus n_2\mathbb{Z} \oplus n_3\mathbb{Z}$ .

# The parameterization of cubic rings

## Definition

A **cubic ring** is a ring which is free of rank 3 as a  $\mathbb{Z}$ -module.

Nondegenerate cubic rings are: Orders in cubic fields;  $Q \oplus n\mathbb{Z}$  for  $Q$  a quadratic order,  $n_1\mathbb{Z} \oplus n_2\mathbb{Z} \oplus n_3\mathbb{Z}$ .

## Theorem (Delone-Faddeev, 1964)

*There is a canonical, explicit bijection between the set of cubic rings up to isomorphism and the set of  $\mathrm{GL}_2(\mathbb{Z})$ -equivalence classes of integral binary cubic forms.*

# The parameterization of cubic rings

## Definition

A **cubic ring** is a ring which is free of rank 3 as a  $\mathbb{Z}$ -module.

Nondegenerate cubic rings are: Orders in cubic fields;  $Q \oplus n\mathbb{Z}$  for  $Q$  a quadratic order,  $n_1\mathbb{Z} \oplus n_2\mathbb{Z} \oplus n_3\mathbb{Z}$ .

## Theorem (Delone-Faddeev, 1964)

*There is a canonical, explicit bijection between the set of cubic rings up to isomorphism and the set of  $\mathrm{GL}_2(\mathbb{Z})$ -equivalence classes of integral binary cubic forms.*

This bijection preserves the discriminant.

# Quartic and quintic rings and fields

Bhargava: Related but more sophisticated techniques apply.



# Dirichlet series for cubic rings

Shintani 1972: Associate a Dirichlet series.

# Dirichlet series for cubic rings

Shintani 1972: Associate a Dirichlet series.

Let  $a(n) :=$  number of cubic rings of discriminant  $n$ .

## Theorem (Shintani, 1972)

*The Dirichlet series  $\sum_{n \geq 1} a(n)n^{-s}$  and  $\sum_{n \geq 1} a(-n)n^{-s}$  have meromorphic continuation to all of  $\mathbb{C}$  and satisfy functional equations.*

# Binary cubic forms, cont.

$L$  is the lattice of **integral** binary cubic forms:

$$L := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 u v^2 + x_4 v^3; \ x_1, x_2, x_3, x_4 \in \mathbb{Z}\}.$$

# Binary cubic forms, cont.

$L$  is the lattice of **integral** binary cubic forms:

$$L := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 u v^2 + x_4 v^3; \ x_1, x_2, x_3, x_4 \in \mathbb{Z}\}.$$

$\widehat{L}$  is the **dual lattice**

$$L := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 u v^2 + x_4 v^3 \in L; \ x_2, x_3 \in 3\mathbb{Z}\}.$$

$\mathrm{SL}_2(\mathbb{Z})$  acts on  $L$  and  $\widehat{L}$  and preserves the discriminant.

# Binary cubic forms, cont.

$L$  is the lattice of **integral** binary cubic forms:

$$L := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 u v^2 + x_4 v^3; \ x_1, x_2, x_3, x_4 \in \mathbb{Z}\}.$$

$\widehat{L}$  is the **dual lattice**

$$L := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 u v^2 + x_4 v^3 \in L; \ x_2, x_3 \in 3\mathbb{Z}\}.$$

$\mathrm{SL}_2(\mathbb{Z})$  acts on  $L$  and  $\widehat{L}$  and preserves the discriminant.

## Definition

The class number  $h(n)$  (resp.  $\widehat{h}(n)$ ) is the number of  $\mathrm{SL}_2(\mathbb{Z})$ -orbits on  $L$  (resp.  $\widehat{L}$ ) of discriminant  $n$ .

# Binary cubic forms, cont.

$L$  is the lattice of **integral** binary cubic forms:

$$L := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 uv^2 + x_4 v^3; \ x_1, x_2, x_3, x_4 \in \mathbb{Z}\}.$$

$\widehat{L}$  is the **dual lattice**

$$L := \{x(u, v) = x_1 u^3 + x_2 u^2 v + x_3 uv^2 + x_4 v^3 \in L; \ x_2, x_3 \in 3\mathbb{Z}\}.$$

$\mathrm{SL}_2(\mathbb{Z})$  acts on  $L$  and  $\widehat{L}$  and preserves the discriminant.

## Definition

The class number  $h(n)$  (resp.  $\widehat{h}(n)$ ) is the number of  $\mathrm{SL}_2(\mathbb{Z})$ -orbits on  $L$  (resp.  $\widehat{L}$ ) of discriminant  $n$ .

Technical point: Actually, we need to adjust for orbits with nontrivial stabilizers.

# The Shintani zeta functions defined

Define

$$\xi_+(L, s) := \sum_{n=1}^{\infty} \frac{h(n)}{n^s}, \quad \xi_-(L, s) := \sum_{n=1}^{\infty} \frac{h(-n)}{n^s},$$

$$\xi_+(\widehat{L}, s) := \sum_{n=1}^{\infty} \frac{\widehat{h}(n)}{n^s}, \quad \xi_-(\widehat{L}, s) := \sum_{n=1}^{\infty} \frac{\widehat{h}(-n)}{n^s},$$

the **Shintani zeta functions**.

# Relation to cubic rings

We have the formula

$$\xi_{\pm}(s) = 2 \sum_{\mathcal{O}} \frac{1}{c(\mathcal{O})} |\text{disc } \mathcal{O}|^{-s},$$

where



# Relation to cubic rings

We have the formula

$$\xi_{\pm}(s) = 2 \sum_{\mathcal{O}} \frac{1}{c(\mathcal{O})} |\text{disc } \mathcal{O}|^{-s},$$

where

- ▶  $c(\mathcal{O}) = 6$  if  $\mathcal{O}$  is a cubic order,

# Relation to cubic rings

We have the formula

$$\xi_{\pm}(s) = 2 \sum_{\mathcal{O}} \frac{1}{c(\mathcal{O})} |\text{disc } \mathcal{O}|^{-s},$$

where

- ▶  $c(\mathcal{O}) = 6$  if  $\mathcal{O}$  is a cubic order,
- ▶  $c(\mathcal{O}) = 2$  if  $\mathcal{O}$  is a sum of a quadratic order and a multiple of  $\mathbb{Z}$ ,

# Relation to cubic rings

We have the formula

$$\xi_{\pm}(s) = 2 \sum_{\mathcal{O}} \frac{1}{c(\mathcal{O})} |\text{disc } \mathcal{O}|^{-s},$$

where

- ▶  $c(\mathcal{O}) = 6$  if  $\mathcal{O}$  is a cubic order,
- ▶  $c(\mathcal{O}) = 2$  if  $\mathcal{O}$  is a sum of a quadratic order and a multiple of  $\mathbb{Z}$ ,
- ▶  $c(\mathcal{O}) = 3$  if  $\mathcal{O}$  is three multiples of  $\mathbb{Z}$ .

# Relation to cubic rings

We have the formula

$$\xi_{\pm}(s) = 2 \sum_{\mathcal{O}} \frac{1}{c(\mathcal{O})} |\text{disc } \mathcal{O}|^{-s},$$

where

- ▶  $c(\mathcal{O}) = 6$  if  $\mathcal{O}$  is a cubic order,
- ▶  $c(\mathcal{O}) = 2$  if  $\mathcal{O}$  is a sum of a quadratic order and a multiple of  $\mathbb{Z}$ ,
- ▶  $c(\mathcal{O}) = 3$  if  $\mathcal{O}$  is three multiples of  $\mathbb{Z}$ .
- ▶ If we limit  $\mathcal{O} \otimes \mathbb{Q}$  to direct sums of cyclic Galois extensions, this can be understood by other means.

# Shintani's Main Theorem

## Theorem (Shintani, 1972)

*The above series converge absolutely for  $\Re(s) > 1$ , have meromorphic continuation to all of  $\mathbb{C}$  with poles at  $s = 1$  and  $s = 5/6$ , and satisfy the functional equation*

$$\begin{pmatrix} \xi_+(L, 1-s) \\ \xi_-(L, 1-s) \end{pmatrix} = \Gamma\left(s - \frac{1}{6}\right) \Gamma(s)^2 \Gamma\left(s + \frac{1}{6}\right) 2^{-1} 3^{6s-2} \pi^{-4s} \times \\ \begin{pmatrix} \sin 2\pi s & \sin \pi s \\ 3 \sin \pi s & \sin 2\pi s \end{pmatrix} \begin{pmatrix} \xi_+(\widehat{L}, s) \\ \xi_-(\widehat{L}, s) \end{pmatrix}. \quad (1)$$

# Diagonalization and Simplification

Datskovsky and Wright diagonalized this matrix.

# Diagonalization and Simplification

Datskovsky and Wright diagonalized this matrix.

Also, Ohno conjectured, and Nakagawa proved, the relations

$$\xi_+(\widehat{L}, s) = 3^{-3s} \xi_-(L, s),$$

$$\xi_-(\widehat{L}, s) = 3^{1-3s} \xi_+(L, s).$$

# Diagonalization and Simplification

Datskovsky and Wright diagonalized this matrix.

Also, Ohno conjectured, and Nakagawa proved, the relations

$$\xi_+(\widehat{L}, s) = 3^{-3s} \xi_-(L, s),$$

$$\xi_-(\widehat{L}, s) = 3^{1-3s} \xi_+(L, s).$$

Putting all this together...



# The “Nice” Shintani zeta function

Write

$$\xi_{\text{add}}(s) := 3^{1/2} \xi_+(L, s) + \xi_-(L, s),$$

$$\xi_{\text{sub}}(s) := 3^{1/2} \xi_+(L, s) - \xi_-(L, s),$$

# The “Nice” Shintani zeta function

Write

$$\xi_{\text{add}}(s) := 3^{1/2} \xi_+(L, s) + \xi_-(L, s),$$

$$\xi_{\text{sub}}(s) := 3^{1/2} \xi_+(L, s) - \xi_-(L, s),$$

$$\Lambda_{\text{add}}(s) := \left( \frac{432}{\pi^4} \right)^{s/2} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s}{2} + \frac{1}{2}\right) \Gamma\left(\frac{s}{2} + \frac{1}{12}\right) \Gamma\left(\frac{s}{2} - \frac{1}{12}\right) \xi_{\text{add}}(s),$$

$$\Lambda_{\text{sub}}(s) := \left( \frac{432}{\pi^4} \right)^{s/2} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s}{2} + \frac{1}{2}\right) \Gamma\left(\frac{s}{2} + \frac{5}{12}\right) \Gamma\left(\frac{s}{2} + \frac{7}{12}\right) \xi_{\text{sub}}(s).$$

# The “Nice” Shintani zeta function

Write

$$\xi_{\text{add}}(s) := 3^{1/2} \xi_+(L, s) + \xi_-(L, s),$$

$$\xi_{\text{sub}}(s) := 3^{1/2} \xi_+(L, s) - \xi_-(L, s),$$

$$\Lambda_{\text{add}}(s) := \left( \frac{432}{\pi^4} \right)^{s/2} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s}{2} + \frac{1}{2}\right) \Gamma\left(\frac{s}{2} + \frac{1}{12}\right) \Gamma\left(\frac{s}{2} - \frac{1}{12}\right) \xi_{\text{add}}(s),$$

$$\Lambda_{\text{sub}}(s) := \left( \frac{432}{\pi^4} \right)^{s/2} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s}{2} + \frac{1}{2}\right) \Gamma\left(\frac{s}{2} + \frac{5}{12}\right) \Gamma\left(\frac{s}{2} + \frac{7}{12}\right) \xi_{\text{sub}}(s).$$

Then

$$\Lambda_{\text{add}}(s) = \Lambda_{\text{add}}(1 - s),$$

$$\Lambda_{\text{sub}}(s) = \Lambda_{\text{sub}}(1 - s).$$

# The “Nice” Shintani zeta function

Write

$$\xi_{\text{add}}(s) := 3^{1/2} \xi_+(L, s) + \xi_-(L, s),$$

$$\xi_{\text{sub}}(s) := 3^{1/2} \xi_+(L, s) - \xi_-(L, s),$$

$$\Lambda_{\text{add}}(s) := \left( \frac{432}{\pi^4} \right)^{s/2} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s}{2} + \frac{1}{2}\right) \Gamma\left(\frac{s}{2} + \frac{1}{12}\right) \Gamma\left(\frac{s}{2} - \frac{1}{12}\right) \xi_{\text{add}}(s),$$

$$\Lambda_{\text{sub}}(s) := \left( \frac{432}{\pi^4} \right)^{s/2} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s}{2} + \frac{1}{2}\right) \Gamma\left(\frac{s}{2} + \frac{5}{12}\right) \Gamma\left(\frac{s}{2} + \frac{7}{12}\right) \xi_{\text{sub}}(s).$$

Then

$$\Lambda_{\text{add}}(s) = \Lambda_{\text{add}}(1 - s),$$

$$\Lambda_{\text{sub}}(s) = \Lambda_{\text{sub}}(1 - s).$$

Both have poles at  $s = 1$ , and  $\Lambda_{\text{add}}$  also has a pole at  $s = 5/6$ .

# Overview of today's talk

These are nice zeta functions, according to the usual axioms.

# Overview of today's talk

These are nice zeta functions, according to the usual axioms.  
But they **don't** have Euler products.

# Overview of today's talk

These are nice zeta functions, according to the usual axioms.

But they **don't** have Euler products.

Today's questions:

# Overview of today's talk

These are nice zeta functions, according to the usual axioms.

But they **don't** have Euler products.

Today's questions:

- ▶ Where do they fit into the general theory of zeta functions?



# Overview of today's talk

These are nice zeta functions, according to the usual axioms.

But they **don't** have Euler products.

Today's questions:

- ▶ Where do they fit into the general theory of zeta functions?
- ▶ Where are the zeroes – and does this tell us anything?

# Overview of today's talk

These are nice zeta functions, according to the usual axioms.

But they **don't** have Euler products.

Today's questions:

- ▶ Where do they fit into the general theory of zeta functions?
- ▶ Where are the zeroes – and does this tell us anything?
- ▶ Have we described all of the Shintani zeta functions?

# Overview of today's talk

These are nice zeta functions, according to the usual axioms.

But they **don't** have Euler products.

Today's questions:

- ▶ Where do they fit into the general theory of zeta functions?
- ▶ Where are the zeroes – and does this tell us anything?
- ▶ Have we described all of the Shintani zeta functions?

Today: *begin* to answer these questions.

# The location of the zeros

**Notation:**  $\xi(s) = \sum_{n \geq 1} a(n)n^{-s}$  is  $\xi_{\text{add}}(s)$  or  $\xi_{\text{sub}}(s)$ .

# The location of the zeros

**Notation:**  $\xi(s) = \sum_{n \geq 1} a(n)n^{-s}$  is  $\xi_{\text{add}}(s)$  or  $\xi_{\text{sub}}(s)$ .

Let  $N(T)$  count the nontrivial zeroes  $s = \sigma + it$  with  $|t| < T$ .  
Then  $\sigma$  must be close to the critical strip. Also,

# The location of the zeros

**Notation:**  $\xi(s) = \sum_{n \geq 1} a(n)n^{-s}$  is  $\xi_{\text{add}}(s)$  or  $\xi_{\text{sub}}(s)$ .

Let  $N(T)$  count the nontrivial zeroes  $s = \sigma + it$  with  $|t| < T$ .  
Then  $\sigma$  must be close to the critical strip. Also,

$$N(T) = \frac{T}{\pi} \log \left( \frac{432 T^4}{(2\pi e)^4} \right) + O(\log T).$$

# The location of the zeros

**Notation:**  $\xi(s) = \sum_{n \geq 1} a(n)n^{-s}$  is  $\xi_{\text{add}}(s)$  or  $\xi_{\text{sub}}(s)$ .

Let  $N(T)$  count the nontrivial zeroes  $s = \sigma + it$  with  $|t| < T$ .  
Then  $\sigma$  must be close to the critical strip. Also,

$$N(T) = \frac{T}{\pi} \log \left( \frac{432 T^4}{(2\pi e)^4} \right) + O(\log T).$$

Folllows by Stirling's formula and contour integration.

# The Riemann Hypothesis

**Question:** Does  $\xi(s)$  satisfy the Riemann hypothesis?



# The Riemann Hypothesis

**Question:** Does  $\xi(s)$  satisfy the Riemann hypothesis?

Using Rubinstein's "L" and Dokchitser's "ComputeL" software,  $\xi(s)$  has zeros

$$0.5 + 4.745125599327 \dots i$$

$$0.5 + 6.962286575567 \dots i$$

$$0.5 + 8.4742944491274 \dots i$$

$$0.5 + 10.152261066735 \dots i$$

$\dots$

# The Riemann Hypothesis

**Question:** Does  $\xi(s)$  satisfy the Riemann hypothesis?

Using Rubinstein's "L" and Dokchitser's "ComputeL" software,  $\xi(s)$  has zeros

$$0.5 + 4.745125599327 \dots i$$

$$0.5 + 6.962286575567 \dots i$$

$$0.5 + 8.4742944491274 \dots i$$

$$0.5 + 10.152261066735 \dots i$$

...

$$0.81420 \dots + 7.05984 \dots i.$$

**General heuristic:**  $L$ -functions with Euler products should satisfy the Riemann hypothesis.

# Euler products and non-Euler products

**General heuristic:**  $L$ -functions with Euler products should satisfy the Riemann hypothesis. Zeta functions without products shouldn't.

# Euler products and non-Euler products

**General heuristic:**  $L$ -functions with Euler products should satisfy the Riemann hypothesis. Zeta functions without products shouldn't.

Other examples of zeta functions without Euler products:

# Euler products and non-Euler products

**General heuristic:**  $L$ -functions with Euler products should satisfy the Riemann hypothesis. Zeta functions without products shouldn't.

Other examples of zeta functions without Euler products:

- ▶ Sums of  $L$ -functions with the same FE,

# Euler products and non-Euler products

**General heuristic:**  $L$ -functions with Euler products should satisfy the Riemann hypothesis. Zeta functions without products shouldn't.

Other examples of zeta functions without Euler products:

- ▶ Sums of  $L$ -functions with the same FE,
- ▶ Epstein zeta functions

$$\sum_{(u,v) \neq (0,0)} (au^2 + buv + cv^2)^{-s}.$$

# Euler products and non-Euler products

**General heuristic:**  $L$ -functions with Euler products should satisfy the Riemann hypothesis. Zeta functions without products shouldn't.

Other examples of zeta functions without Euler products:

- ▶ Sums of  $L$ -functions with the same FE,
- ▶ Epstein zeta functions

$$\sum_{(u,v) \neq (0,0)} (au^2 + buv + cv^2)^{-s}.$$

For  $a, b, c \in \mathbb{Z}$  and  $b^2 - 4ac < 0$ , special case of above.



# Zeros of Epstein zeta functions

**Working hypothesis:** The zeros of Shintani zeta functions behave like the zeros of Epstein zeta functions.

# Zeros of Epstein zeta functions

**Working hypothesis:** The zeros of Shintani zeta functions behave like the zeros of Epstein zeta functions.

Epstein zeta functions satisfy:

# Zeros of Epstein zeta functions

**Working hypothesis:** The zeros of Shintani zeta functions behave like the zeros of Epstein zeta functions.

Epstein zeta functions satisfy:

- ▶ At least  $\gg T$  zeros are on the critical line.

# Zeros of Epstein zeta functions

**Working hypothesis:** The zeros of Shintani zeta functions behave like the zeros of Epstein zeta functions.

Epstein zeta functions satisfy:

- ▶ At least  $\gg T$  zeros are on the critical line.
- ▶ At most  $O(T)$  zeros are to the right of any line  $\operatorname{Re}(s) > \sigma > 1/2$ .

# Zeros of Epstein zeta functions

**Working hypothesis:** The zeros of Shintani zeta functions behave like the zeros of Epstein zeta functions.

Epstein zeta functions satisfy:

- ▶ At least  $\gg T$  zeros are on the critical line.
- ▶ At most  $O(T)$  zeros are to the right of any line  $\operatorname{Re}(s) > \sigma > 1/2$ .
- ▶ At least  $\gg T$  nontrivial zeros are outside the critical strip.

# Zeros of Epstein zeta functions

**Working hypothesis:** The zeros of Shintani zeta functions behave like the zeros of Epstein zeta functions.

Epstein zeta functions satisfy:

- ▶ At least  $\gg T$  zeros are on the critical line.
- ▶ At most  $O(T)$  zeros are to the right of any line  $\operatorname{Re}(s) > \sigma > 1/2$ .
- ▶ At least  $\gg T$  nontrivial zeros are outside the critical strip.

Open question: Are almost all the zeros on  $\operatorname{Re}(s) = 1/2$ ?

# Zeros of Epstein zeta functions

**Working hypothesis:** The zeros of Shintani zeta functions behave like the zeros of Epstein zeta functions.

Epstein zeta functions satisfy:

- ▶ At least  $\gg T$  zeros are on the critical line.
- ▶ At most  $O(T)$  zeros are to the right of any line  $\operatorname{Re}(s) > \sigma > 1/2$ .
- ▶ At least  $\gg T$  nontrivial zeros are outside the critical strip.

Open question: Are almost all the zeros on  $\operatorname{Re}(s) = 1/2$ ?

See Hejhal, *Epstein zeta functions and supercomputers*.

# Zeros of Shintani zeta functions

The bad news:



# Zeros of Shintani zeta functions

The bad news:

- ▶ The proofs of these statements don't generalize well.

# Zeros of Shintani zeta functions

The bad news:

- ▶ The proofs of these statements don't generalize well.
- ▶ We can't seem to use much particular information about the Shintani zeta function.

# Zeros of Shintani zeta functions

The bad news:

- ▶ The proofs of these statements don't generalize well.
- ▶ We can't seem to use much particular information about the Shintani zeta function.

The good news:

# Zeros of Shintani zeta functions

The bad news:

- ▶ The proofs of these statements don't generalize well.
- ▶ We can't seem to use much particular information about the Shintani zeta function.

The good news:

- ▶ We can try to work in general.

# Zeros outside the critical strip

## Theorem (Soundararajan-T.)

*The Shintani zeta functions  $\xi_{\text{add}}$ ,  $\xi_{\text{sub}}$ , and  $\xi_{-}$  have infinitely many zeros to the right of  $\Re(s) = 1$ .*

# Zeros outside the critical strip

## Theorem (Soundararajan-T.)

*The Shintani zeta functions  $\xi_{\text{add}}$ ,  $\xi_{\text{sub}}$ , and  $\xi_{-}$  have infinitely many zeros to the right of  $\Re(s) = 1$ .*

$\xi_{+}$  should too, and we have “numerical evidence”.

# Zeros outside the critical strip

## Theorem (Soundararajan-T.)

*The Shintani zeta functions  $\xi_{\text{add}}$ ,  $\xi_{\text{sub}}$ , and  $\xi_{-}$  have infinitely many zeros to the right of  $\Re(s) = 1$ .*

$\xi_{+}$  should too, and we have “numerical evidence”.

Our results follow from a more general method.

If a zeta function has zeros in  $\Re s > 1$ , then this *proves* the zeta function doesn't have an Euler product.



If a zeta function has zeros in  $\Re s > 1$ , then this *proves* the zeta function doesn't have an Euler product.

Stark: Solve the class number 1 problem this way.

# Stark's challenge

## Theorem (Folk Theorem)

Write

$$\zeta(s, Q) = \sum_{(u,v) \neq (0,0)} (u^2 + uv + cv^2)^{-s}.$$

*If  $c > 41$ ,  $\zeta(s, Q)$  has a zero  $s$  with  $\sigma > 1$ .*

# Stark's challenge

## Theorem (Folk Theorem)

Write

$$\zeta(s, Q) = \sum_{(u,v) \neq (0,0)} (u^2 + uv + cv^2)^{-s}.$$

*If  $c > 41$ ,  $\zeta(s, Q)$  has a zero  $s$  with  $\sigma > 1$ .*

Problems:

# Stark's challenge

## Theorem (Folk Theorem)

Write

$$\zeta(s, Q) = \sum_{(u,v) \neq (0,0)} (u^2 + uv + cv^2)^{-s}.$$

If  $c > 41$ ,  $\zeta(s, Q)$  has a zero  $s$  with  $\sigma > 1$ .

Problems:

- ▶ The only “proof” uses the class number one determination.

# Stark's challenge

## Theorem (Folk Theorem)

Write

$$\zeta(s, Q) = \sum_{(u,v) \neq (0,0)} (u^2 + uv + cv^2)^{-s}.$$

*If  $c > 41$ ,  $\zeta(s, Q)$  has a zero  $s$  with  $\sigma > 1$ .*

Problems:

- ▶ The only “proof” uses the class number one determination.
- ▶ It hasn't been proved at all for algebraic numbers other than integers.

# Stark's challenge (cont.)

“So the challenge is clear: prove the folk theorem, but better still,  
FIND A PURELY ANALYTIC PROOF OF THE FOLK  
THEOREM.”

## Stark's challenge (cont.)

“So the challenge is clear: prove the folk theorem, but better still,  
FIND A PURELY ANALYTIC PROOF OF THE FOLK  
THEOREM.”

Stark's all caps, not mine.

# Sketch proof of theorem

It suffices to find *one* zero outside the critical strip.



# Sketch proof of theorem

It suffices to find *one* zero outside the critical strip.

Proof:

# Sketch proof of theorem

It suffices to find *one* zero outside the critical strip.

Proof:

- ▶ Uniformly in  $\Re(s) \geq 1 + \delta$  for any  $\delta > 0$ ,  $\xi(s)$  is “almost periodic” in  $t$ .

# Sketch proof of theorem

It suffices to find *one* zero outside the critical strip.

Proof:

- ▶ Uniformly in  $\Re(s) \geq 1 + \delta$  for any  $\delta > 0$ ,  $\xi(s)$  is “almost periodic” in  $t$ .
- ▶ By Rouché’s theorem, this zero repeats in every almost period.

# Sketch proof of theorem

It suffices to find *one* zero outside the critical strip.

Proof:

- ▶ Uniformly in  $\Re(s) \geq 1 + \delta$  for any  $\delta > 0$ ,  $\xi(s)$  is “almost periodic” in  $t$ .
- ▶ By Rouché’s theorem, this zero repeats in every almost period.
- ▶ For  $\xi_{\text{add}}$  and  $\xi_{\text{sub}}$ , we easily found a zero numerically.

# Sketch proof of theorem

It suffices to find *one* zero outside the critical strip.

Proof:

- ▶ Uniformly in  $\Re(s) \geq 1 + \delta$  for any  $\delta > 0$ ,  $\xi(s)$  is “almost periodic” in  $t$ .
- ▶ By Rouché’s theorem, this zero repeats in every almost period.
- ▶ For  $\xi_{\text{add}}$  and  $\xi_{\text{sub}}$ , we easily found a zero numerically.
- ▶ For  $\xi_-$ , we have to work harder.

# A zero of $\xi_-(s)$ in $\Re s > 1$

To “find a zero”:

# A zero of $\xi_-(s)$ in $\Re s > 1$

To “find a zero”:

- ▶ If  $\chi(n)$  is completely multiplicative, taking values in  $\pm 1$ , define the twisted zeta function

$$\xi(s, \chi) := \sum_{n>0} a(n) \chi(n) n^{-s}.$$

# A zero of $\xi_-(s)$ in $\Re s > 1$

To “find a zero”:

- ▶ If  $\chi(n)$  is completely multiplicative, taking values in  $\pm 1$ , define the twisted zeta function

$$\xi(s, \chi) := \sum_{n>0} a(n)\chi(n)n^{-s}.$$

- ▶ Twisting by such  $\chi$  is “like” shifting up or down the imaginary axis.



# A zero of $\xi_-(s)$ in $\Re s > 1$

To “find a zero”:

- ▶ If  $\chi(n)$  is completely multiplicative, taking values in  $\pm 1$ , define the twisted zeta function

$$\xi(s, \chi) := \sum_{n>0} a(n)\chi(n)n^{-s}.$$

- ▶ Twisting by such  $\chi$  is “like” shifting up or down the imaginary axis.
- ▶ Find a  $\chi$  and  $\sigma > 1$  with  $\xi(\sigma, \chi) = 0$ .

# A zero of $\xi_-(s)$ in $\Re s > 1$

To “find a zero”:

- ▶ If  $\chi(n)$  is completely multiplicative, taking values in  $\pm 1$ , define the twisted zeta function

$$\xi(s, \chi) := \sum_{n>0} a(n)\chi(n)n^{-s}.$$

- ▶ Twisting by such  $\chi$  is “like” shifting up or down the imaginary axis.
- ▶ Find a  $\chi$  and  $\sigma > 1$  with  $\xi(\sigma, \chi) = 0$ .
- ▶ If we can, then we have zeros outside the critical strip.

# Finding $\chi$ and $\sigma$

$$\xi(s) = \xi_-(s) = \frac{1}{3^s} + \frac{1}{4^s} + \frac{1}{7^s} + \frac{1}{8^s} + \cdots$$

To find  $\chi$  and  $\sigma$  with  $\xi(\sigma, \chi) = 0$ :

# Finding $\chi$ and $\sigma$

$$\xi(s) = \xi_-(s) = \frac{1}{3^s} + \frac{1}{4^s} + \frac{1}{7^s} + \frac{1}{8^s} + \cdots$$

To find  $\chi$  and  $\sigma$  with  $\xi(\sigma, \chi) = 0$ :

- ▶ Assuming  $\chi(3) = 1$ , enough if  $\xi(\sigma, \chi) < 0$ .

# Finding $\chi$ and $\sigma$

$$\xi(s) = \xi_-(s) = \frac{1}{3^s} + \frac{1}{4^s} + \frac{1}{7^s} + \frac{1}{8^s} + \cdots$$

To find  $\chi$  and  $\sigma$  with  $\xi(\sigma, \chi) = 0$ :

- ▶ Assuming  $\chi(3) = 1$ , enough if  $\xi(\sigma, \chi) < 0$ .
- ▶ Compute the first million or so  $a(n)$ ; determine a useful  $\chi$  by inspection.

# Finding $\chi$ and $\sigma$

$$\xi(s) = \xi_-(s) = \frac{1}{3^s} + \frac{1}{4^s} + \frac{1}{7^s} + \frac{1}{8^s} + \cdots$$

To find  $\chi$  and  $\sigma$  with  $\xi(\sigma, \chi) = 0$ :

- ▶ Assuming  $\chi(3) = 1$ , enough if  $\xi(\sigma, \chi) < 0$ .
- ▶ Compute the first million or so  $a(n)$ ; determine a useful  $\chi$  by inspection.

For appropriate  $\chi$ ,

$$\sum_{n \leq 10^6} a(n)\chi(n)n^{-1.3} = -.162 \cdots, \quad \sum_{n > 10^6} |a(n)|n^{-1.3} < 0.1.$$

# The method for $\xi_+(s)$

$$\xi_+(s) = 1/3 + \frac{1}{4^s} + \frac{1}{5^s} + \frac{1}{8^s} + \cdots$$

# The method for $\xi_+(s)$

$$\xi_+(s) = 1/3 + \frac{1}{4^s} + \frac{1}{5^s} + \frac{1}{8^s} + \cdots$$

Bad luck: The first two coefficients are squares.



# The method for $\xi_+(s)$

$$\xi_+(s) = 1/3 + \frac{1}{4^s} + \frac{1}{5^s} + \frac{1}{8^s} + \dots$$

Bad luck: The first two coefficients are squares.

Setting  $\chi(p) = -1$  for  $p \neq 7, 11, 19, 23$ ,

$$\sum_{n \leq 10^6} a(n)\chi(n)n^{-1.1} = -.103\dots$$

# The method for $\xi_+(s)$

$$\xi_+(s) = 1/3 + \frac{1}{4^s} + \frac{1}{5^s} + \frac{1}{8^s} + \dots$$

Bad luck: The first two coefficients are squares.

Setting  $\chi(p) = -1$  for  $p \neq 7, 11, 19, 23$ ,

$$\sum_{n \leq 10^6} a(n) \chi(n) n^{-1.1} = -.103 \dots$$

$$\sum_{n > 10^6} |a(n)| n^{-1.1} = 7.217 \dots$$

# The method for $\xi_+(s)$

$$\xi_+(s) = 1/3 + \frac{1}{4^s} + \frac{1}{5^s} + \frac{1}{8^s} + \dots$$

Bad luck: The first two coefficients are squares.

Setting  $\chi(p) = -1$  for  $p \neq 7, 11, 19, 23$ ,

$$\sum_{n \leq 10^6} a(n) \chi(n) n^{-1.1} = -.103 \dots$$

$$\sum_{n > 10^6} |a(n)| n^{-1.1} = 7.217 \dots$$

Conclusion: Compute 100,000,000,000,000 coefficients and try again.

# The inverse question

## Conjecture

Zeta functions with no zeros in the region of absolute convergence must “more or less” be Euler products.

# The inverse question

## Conjecture

Zeta functions with no zeros in the region of absolute convergence must “more or less” be Euler products.

Caution: We haven’t defined “more or less”.

# The inverse question

## Conjecture

Zeta functions with no zeros in the region of absolute convergence must “more or less” be Euler products.

Caution: We haven’t defined “more or less”.

We have technical partial results in this direction.

# All the Shintani zeta functions

Have we defined all the “Shintani zeta functions”?

# All the Shintani zeta functions

Have we defined all the “Shintani zeta functions”?

**No**, there are also:



# All the Shintani zeta functions

Have we defined all the “Shintani zeta functions”?

**No**, there are also:

- ▶ **Adelic** Shintani zeta functions associated to any global field (Datskovsky-Wright),

# All the Shintani zeta functions

Have we defined all the “Shintani zeta functions”?

**No**, there are also:

- ▶ **Adelic** Shintani zeta functions associated to any global field (Datskovsky-Wright),
- ▶ Shintani zeta functions associated to **other invariant lattices** (Ohno, Taniguchi, Wakatsuki),

# All the Shintani zeta functions

Have we defined all the “Shintani zeta functions”?

**No**, there are also:

- ▶ **Adelic** Shintani zeta functions associated to any global field (Datskovsky-Wright),
- ▶ Shintani zeta functions associated to **other invariant lattices** (Ohno, Taniguchi, Wakatsuki),
- ▶ Shintani zeta functions associated to **quartic** rings (Yukie),

# All the Shintani zeta functions

Have we defined all the “Shintani zeta functions”?

**No**, there are also:

- ▶ **Adelic** Shintani zeta functions associated to any global field (Datskovsky-Wright),
- ▶ Shintani zeta functions associated to **other invariant lattices** (Ohno, Taniguchi, Wakatsuki),
- ▶ Shintani zeta functions associated to **quartic** rings (Yukie),
- ▶ A theory for **general** prehomogeneous vector spaces (Sato-Shintani),

# All the Shintani zeta functions

Have we defined all the “Shintani zeta functions”?

**No**, there are also:

- ▶ **Adelic** Shintani zeta functions associated to any global field (Datskovsky-Wright),
- ▶ Shintani zeta functions associated to **other invariant lattices** (Ohno, Taniguchi, Wakatsuki),
- ▶ Shintani zeta functions associated to **quartic** rings (Yukie),
- ▶ A theory for **general** prehomogeneous vector spaces (Sato-Shintani),
- ▶ ...?

# Cubic rings and sieve methods

Preliminary results; to be carefully checked.

# Cubic rings and sieve methods

Preliminary results; to be carefully checked.

**Goal:** Apply classical sieve methods to cubic field discriminants.

# Cubic rings and sieve methods

Preliminary results; to be carefully checked.

**Goal:** Apply classical sieve methods to cubic field discriminants.

Typical sieve results:



# Cubic rings and sieve methods

Preliminary results; to be carefully checked.

**Goal:** Apply classical sieve methods to cubic field discriminants.

Typical sieve results:

- ▶ There are  $\ll X / \log X$  cubic fields of prime discriminant  $< X$ .

# Cubic rings and sieve methods

Preliminary results; to be carefully checked.

**Goal:** Apply classical sieve methods to cubic field discriminants.

Typical sieve results:

- ▶ There are  $\ll X/\log X$  cubic fields of prime discriminant  $< X$ .
- ▶ There are  $\gg X/\log X$  cubic fields of discriminant  $< X$  with at most  $r$  prime factors.

# Discriminants in progressions

Let  $b(n) :=$  number of cubic rings of discriminant  $n > 0$  or  $-n < 0$ .

# Discriminants in progressions

Let  $b(n) :=$  number of cubic rings of discriminant  $n > 0$  or  $-n < 0$ .

We can easily determine  $b(n)$  from the Shintani coefficients  $a(n)$ .

# Discriminants in progressions

Let  $b(n) :=$  number of cubic rings of discriminant  $n > 0$  or  $-n < 0$ .

We can easily determine  $b(n)$  from the Shintani coefficients  $a(n)$ .

The central proposition:

# Discriminants in progressions

Let  $b(n) :=$  number of cubic rings of discriminant  $n > 0$  or  $-n < 0$ .

We can easily determine  $b(n)$  from the Shintani coefficients  $a(n)$ .

The central proposition:

$$\sum_{n \leq X; d|n} b(n) = C\omega(d)X + O(X^\alpha),$$

for a **multiplicative**  $\omega(d)$  and any  $\alpha < 1$ .

# Discriminants in progressions

Let  $b(n) :=$  number of cubic rings of discriminant  $n > 0$  or  $-n < 0$ .

We can easily determine  $b(n)$  from the Shintani coefficients  $a(n)$ .

The central proposition:

$$\sum_{n \leq X; d|n} b(n) = C\omega(d)X + O(X^\alpha),$$

for a **multiplicative**  $\omega(d)$  and any  $\alpha < 1$ .

Or even better:

$$\sum_{n \leq X; d|n} b(n) = C\omega(d)X + C'\nu(d)X^{5/6} + O(X^\alpha),$$

where  $\nu(d)$  is also multiplicative and  $\alpha < 5/6$ .

# Discriminants in progressions (II)

Strategies of proof:



# Discriminants in progressions (II)

Strategies of proof:

- ▶ Use the Delone-Faddeev parameterization, as in works of Bhargava.

# Discriminants in progressions (II)

Strategies of proof:

- ▶ Use the Delone-Faddeev parameterization, as in works of Bhargava.
- ▶ Our proposal: Give an analytic proof, using Shintani zeta functions.

# Discriminants in progressions (II)

Strategies of proof:

- ▶ Use the Delone-Faddeev parameterization, as in works of Bhargava.
- ▶ Our proposal: Give an analytic proof, using Shintani zeta functions.
- ▶ An analytic proof should detect the secondary term.

# Discriminants in progressions (III)

## Proposition (preliminary)

We have

$$\sum_{n \leq X; d|n} b(n) = C\omega(d)X + C'\nu(d)X^{5/6} + O(d^3 X^\alpha)$$

as before, with  $\alpha = 3/5 + \epsilon$ .

# Discriminants in progressions (III)

## Proposition (preliminary)

We have

$$\sum_{n \leq X; d|n} b(n) = C\omega(d)X + C'\nu(d)X^{5/6} + O(d^3 X^\alpha)$$

as before, with  $\alpha = 3/5 + \epsilon$ .

**Ongoing work:** Reduce or eliminate the dependence on  $d$ .

# The $d$ -divisible Shintani zeta function

Start with Datskovsky-Wright's **adelic** Shintani zeta function.

# The $d$ -divisible Shintani zeta function

Start with Datskovsky-Wright's **adelic** Shintani zeta function.

This allows us to introduce  $p$ -adic conditions.

# The $d$ -divisible Shintani zeta function

Start with Datskovsky-Wright's **adelic** Shintani zeta function.

This allows us to introduce  $p$ -adic conditions.

Define the  $d$ -divisible Shintani zeta function

$$\xi_{d,\pm} := \sum_{d|n} a(n) n^{-s}.$$



# The $d$ -divisible Shintani zeta function

Start with Datskovsky-Wright's **adelic** Shintani zeta function.

This allows us to introduce  $p$ -adic conditions.

Define the  $d$ -divisible Shintani zeta function

$$\xi_{d,\pm} := \sum_{d|n} a(n) n^{-s}.$$

By Datskovsky-Wright, this has analytic continuation and a functional equation.

# The $d$ -divisible Shintani zeta function

Start with Datskovsky-Wright's **adelic** Shintani zeta function.

This allows us to introduce  $p$ -adic conditions.

Define the  $d$ -divisible Shintani zeta function

$$\xi_{d,\pm} := \sum_{d|n} a(n) n^{-s}.$$

By Datskovsky-Wright, this has analytic continuation and a functional equation.

**Note:** Taniguchi has also considered this.

# The $d$ -divisible Shintani zeta function (II)

The good news:

# The $d$ -divisible Shintani zeta function (II)

The good news:

- ▶ By Sato-Shintani's Tauberian theorem, we get our proposition.

# The $d$ -divisible Shintani zeta function (II)

The good news:

- ▶ By Sato-Shintani's Tauberian theorem, we get our proposition.
- ▶ This implies the sieve results stated earlier.

# The $d$ -divisible Shintani zeta function (II)

The good news:

- ▶ By Sato-Shintani's Tauberian theorem, we get our proposition.
- ▶ This implies the sieve results stated earlier.
- ▶ Possible generalization to quartic and quintic fields.

# The Fourier transform

We must analyze the Fourier transform of a certain  $p$ -adic test function.

# The Fourier transform

We must analyze the Fourier transform of a certain  $p$ -adic test function.

Previous proposition: Trivially estimate this Fourier transform.



# The Fourier transform

We must analyze the Fourier transform of a certain  $p$ -adic test function.

Previous proposition: Trivially estimate this Fourier transform.

Our goal: Understand this Fourier transform better.

# The Fourier transform

We must analyze the Fourier transform of a certain  $p$ -adic test function.

Previous proposition: Trivially estimate this Fourier transform.

Our goal: Understand this Fourier transform better.

Question: Is this Fourier transform something “nice”?

# Bounds for cubic fields

Question: How many cubic fields of a single discriminant  $n$  are there?

# Bounds for cubic fields

Question: How many cubic fields of a single discriminant  $n$  are there? Almost equivalently, what bounds do we have for  $a(n)$ ?

# Bounds for cubic fields

Question: How many cubic fields of a single discriminant  $n$  are there? Almost equivalently, what bounds do we have for  $a(n)$ ?

Ellenberg and Venkatesh:

$$a(n) \ll n^{1/3+\epsilon}.$$

Improving this would allow all sorts of analytic techniques to work.

# Improving Ellenberg-Venkatesh

# Improving Ellenberg-Venkatesh

One idea to improve this (perhaps conditionally):

One idea to improve this (perhaps conditionally):

- ▶ For any  $d$  with  $a(d) \asymp d^{1/3}$ , consider the  $d$ -divisible Shintani zeta function.



One idea to improve this (perhaps conditionally):

- ▶ For any  $d$  with  $a(d) \asymp d^{1/3}$ , consider the  $d$ -divisible Shintani zeta function.
- ▶ Its first Fourier coefficient is much larger than we were expecting.

One idea to improve this (perhaps conditionally):

- ▶ For any  $d$  with  $a(d) \asymp d^{1/3}$ , consider the  $d$ -divisible Shintani zeta function.
- ▶ Its first Fourier coefficient is much larger than we were expecting.
- ▶ This will greatly affect the analytic behavior.

One idea to improve this (perhaps conditionally):

- ▶ For any  $d$  with  $a(d) \asymp d^{1/3}$ , consider the  $d$ -divisible Shintani zeta function.
- ▶ Its first Fourier coefficient is much larger than we were expecting.
- ▶ This will greatly affect the analytic behavior.
- ▶ The hope: use this to prove something.