

Quantifier Examples

January 19, 2024

Abraham Lincoln

“You can fool some of the people all of the time, and all of the people some of the time, but you can not fool all of the people all of the time.”

Abraham Lincoln

“You can fool some of the people all of the time, and all of the people some of the time, but you can not fool all of the people all of the time.”

Formalize and negate each part.

Extra Credit Question From Hell



Weierstrass's theorem

Theorem (Weierstrass)

There exists a real-valued function which is continuous everywhere and differentiable nowhere.

The setup

Given:

The setup

Given:

- ▶ We write $\lim_{x \rightarrow a} f(x) = L$ if for every $\epsilon > 0$ there exists $\delta > 0$ such that whenever $|x - a| < \delta$ we have $|f(x) - L| < \epsilon$.

The setup

Given:

- ▶ We write $\lim_{x \rightarrow a} f(x) = L$ if for every $\epsilon > 0$ there exists $\delta > 0$ such that whenever $|x - a| < \delta$ we have $|f(x) - L| < \epsilon$.
- ▶ We say that $f(x)$ is continuous at $x = a$ if $\lim_{x \rightarrow a} f(x)$ exists and equals $f(a)$.

The setup

Given:

- ▶ We write $\lim_{x \rightarrow a} f(x) = L$ if for every $\epsilon > 0$ there exists $\delta > 0$ such that whenever $|x - a| < \delta$ we have $|f(x) - L| < \epsilon$.
- ▶ We say that $f(x)$ is continuous at $x = a$ if $\lim_{x \rightarrow a} f(x)$ exists and equals $f(a)$.
- ▶ We say that $f(x)$ is differentiable at $x = a$ if the limit

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

exists.

The setup

Given:

- ▶ We write $\lim_{x \rightarrow a} f(x) = L$ if for every $\epsilon > 0$ there exists $\delta > 0$ such that whenever $|x - a| < \delta$ we have $|f(x) - L| < \epsilon$.
- ▶ We say that $f(x)$ is continuous at $x = a$ if $\lim_{x \rightarrow a} f(x)$ exists and equals $f(a)$.
- ▶ We say that $f(x)$ is differentiable at $x = a$ if the limit

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

exists.

- ▶ We say that $f(x)$ is continuous everywhere if, for all $a \in \mathbb{R}$, $f(x)$ is continuous at $x = a$. Analogously for 'differentiable everywhere'.

The extra credit problem

The extra credit problem

- Formalize, and then negate Weierstrass's theorem in predicate logic. Your predicates should only involve inequalities and should not use any calculus terminology.