

# An Overview of Number Field Counting

Frank Thorne

University of South Carolina

December 11, 2024

## Definition

For any integer  $d \geq 1$ , write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\mathrm{Disc}(K)| < X\}.$$

## Definition

For any integer  $d \geq 1$ , write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X\}.$$

and for each transitive subgroup  $G \subseteq S_d$ ,

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X, \text{Gal}(K^c/\mathbb{Q}) = G\},$$

## Definition

For any integer  $d \geq 1$ , write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X\}.$$

and for each transitive subgroup  $G \subseteq S_d$ ,

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X, \text{Gal}(K^c/\mathbb{Q}) = G\},$$

so that

$$N_d(X) = \sum_{\substack{G \subseteq S_d \\ \text{transitive}}} N_d(X, G).$$

## Definition

For any integer  $d \geq 1$ , write

$$N_d(X) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X\}.$$

and for each transitive subgroup  $G \subseteq S_d$ ,

$$N_d(X, G) := \#\{K : [K : \mathbb{Q}] = d, |\text{Disc}(K)| < X, \text{Gal}(K^c/\mathbb{Q}) = G\},$$

so that

$$N_d(X) = \sum_{\substack{G \subseteq S_d \\ \text{transitive}}} N_d(X, G).$$

## Theorem (Finiteness – Hermite)

*For each  $d$  and  $X$ ,  $N_d(X)$  is finite.*

## Theorem (Finiteness – Hermite)

*For each  $d$  and  $X$ ,  $N_d(X)$  is finite.*

## Theorem (Minkowski's Lower Bound)

*If  $[K : \mathbb{Q}] = d$ , then*

$$|\mathrm{Disc}(K)| \geq \left(\frac{d^d}{d!}\right)^2 \left(\frac{\pi}{4}\right)^d.$$

## Theorem (Finiteness – Hermite)

*For each  $d$  and  $X$ ,  $N_d(X)$  is finite.*

## Theorem (Minkowski's Lower Bound)

*If  $[K : \mathbb{Q}] = d$ , then*

$$|\mathrm{Disc}(K)| \geq \left(\frac{d^d}{d!}\right)^2 \left(\frac{\pi}{4}\right)^d.$$

*In other words,*

$$N_d(X) = 0 \quad \text{for } X < (5.803 \cdots + o(1))^d.$$



# The Inverse Galois Problem

## Conjecture

For every  $d$  and transitive subgroup  $G \subseteq S_d$ ,

$$X \text{ big enough} \implies N_d(X, G) \neq 0.$$

# The Inverse Galois Problem

## Conjecture

For every  $d$  and transitive subgroup  $G \subseteq S_d$ ,

$$X \text{ big enough} \implies N_d(X, G) \neq 0.$$

Proof.

(Your Name Here)



# Malle's Conjecture

## Conjecture

In fact we have

$$N_d(X, G) \sim c(G)X^{1/a(G)}(\log X)^{b(G)},$$

where  $a(G) \geq 1$  and  $b(G) \geq 0$  are **explicitly described** integers.

# Malle's Conjecture

## Conjecture

In fact we have

$$N_d(X, G) \sim c(G) X^{1/a(G)} (\log X)^{b(G)},$$

where  $a(G) \geq 1$  and  $b(G) \geq 0$  are **explicitly described** integers.

Indeed,

$$a(G) := \min_{1 \neq \sigma \in G} (d - \#\text{cycles of } \sigma).$$

# Malle's Conjecture

## Conjecture

In fact we have

$$N_d(X, G) \sim c(G) X^{1/a(G)} (\log X)^{b(G)},$$

where  $a(G) \geq 1$  and  $b(G) \geq 0$  are **explicitly described** integers.

Indeed,

$$a(G) := \min_{1 \neq \sigma \in G} (d - \#\text{cycles of } \sigma).$$

So,

$$a(S_d) = 1, \quad a(A_d) = 2, \quad a(D_4) = 1, \quad a(C_p) = p - 1, \dots$$

# Malle's Conjecture – Very Rough Explanation

$$a(G) := \min_{1 \neq \sigma \in G} (d - \#\text{cycles of } \sigma).$$

# Malle's Conjecture – Very Rough Explanation

$$a(G) := \min_{1 \neq \sigma \in G} (d - \#\text{cycles of } \sigma).$$

If  $p \mid \text{Disc}(K)$ , then  $p^{a(G)} \mid \text{Disc}(K)$ .

# Malle's Conjecture – Very Rough Explanation

$$a(G) := \min_{1 \neq \sigma \in G} (d - \#\text{cycles of } \sigma).$$

If  $p \mid \text{Disc}(K)$ , then  $p^{a(G)} \mid \text{Disc}(K)$ .

Very **very** **very** roughly,

$$\sum_{\text{Gal}(\widehat{K}/\mathbb{Q}) \simeq G} |\text{Disc}(K)|^{-s} \sim \prod_p \left( 1 + \frac{b(G)}{p^{a(G)}} + \frac{??}{p^{a(G)+1}} + \dots \right)$$



# Four Methods to Count Number Fields

- ▶ Generator methods (Schmidt, Ellenberg-Venkatesh, ...)

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)  
Use **class field theory** and/or **Kummer theory**.

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)  
Use **class field theory** and/or **Kummer theory**.  
Get good results when  $G$  is abelian.

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)  
Use **class field theory** and/or **Kummer theory**.  
Get good results when  $G$  is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)  
Use **class field theory** and/or **Kummer theory**.  
Get good results when  $G$  is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)  
Count **lattice points** in **prehomogeneous representations**.



# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)  
Use **class field theory** and/or **Kummer theory**.  
Get good results when  $G$  is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)  
Count **lattice points** in **prehomogeneous representations**.  
Brilliant when it works.

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)  
Use **class field theory** and/or **Kummer theory**.  
Get good results when  $G$  is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)  
Count **lattice points** in **prehomogeneous representations**.  
Brilliant when it works.
- ▶ **Inductive methods** (Klüners, Cohen-Diaz-Olivier, ...)

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)  
Use **class field theory** and/or **Kummer theory**.  
Get good results when  $G$  is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)  
Count **lattice points** in **prehomogeneous representations**.  
Brilliant when it works.
- ▶ **Inductive methods** (Klüners, Cohen-Diaz-Olivier, ...)  
Obtain **old results from new**.

# Four Methods to Count Number Fields

- ▶ **Generator methods** (Schmidt, Ellenberg-Venkatesh, ...)  
Count in terms of appropriate **algebraic integers**.  
Usually seeking upper bounds.
- ▶ **Abelian methods** (Cohn, Wright, Mäki, ...)  
Use **class field theory** and/or **Kummer theory**.  
Get good results when  $G$  is abelian.
- ▶ **Parametrization methods** (Davenport-Heilbronn, Bhargava, ...)  
Count **lattice points** in **prehomogeneous representations**.  
Brilliant when it works.
- ▶ **Inductive methods** (Klüners, Cohen-Diaz-Olivier, ...)  
Obtain **old results from new**.  
Expand the scope of existing methods.

If  $\alpha \in \mathcal{O}_K$  is a generator of  $K/\mathbb{Q}$ , then  $\mathbb{Z}[\alpha] \subseteq \mathcal{O}_K$  and

$$\begin{aligned} |\mathrm{Disc}(\mathcal{O}_K)| &= \mathrm{Disc}(\mathbb{Z}[\alpha]) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2} \\ &= \mathrm{Disc}(\mathrm{minpoly}_\alpha) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2}. \end{aligned}$$

If  $\alpha \in \mathcal{O}_K$  is a generator of  $K/\mathbb{Q}$ , then  $\mathbb{Z}[\alpha] \subseteq \mathcal{O}_K$  and

$$\begin{aligned} |\mathrm{Disc}(\mathcal{O}_K)| &= \mathrm{Disc}(\mathbb{Z}[\alpha]) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2} \\ &= \mathrm{Disc}(\mathrm{minpoly}_\alpha) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^{-2}. \end{aligned}$$

## Theorem (Schmidt)

For each  $d$  we have

$$N_d(X) \ll X^{\frac{d+2}{4}}.$$

# Schmidt's proof

- By [Minkowski's theory](#), there exists  $\alpha \in \mathcal{O}_K$  with [trace 0](#) and  $||\alpha||_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$  for all embeddings  $\sigma : K \mapsto \mathbb{C}$ .



# Schmidt's proof

- ▶ By Minkowski's theory, there exists  $\alpha \in \mathcal{O}_K$  with trace 0 and  $||\alpha||_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$  for all embeddings  $\sigma : K \hookrightarrow \mathbb{C}$ .
- ▶ Assume that  $\mathbb{Q}(\alpha) = K$ . (If not, induct.)

# Schmidt's proof

- ▶ By Minkowski's theory, there exists  $\alpha \in \mathcal{O}_K$  with trace 0 and  $||\alpha||_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$  for all embeddings  $\sigma : K \mapsto \mathbb{C}$ .
- ▶ Assume that  $\mathbb{Q}(\alpha) = K$ . (If not, induct.)
- ▶ The minimal polynomial of  $\alpha$  is

$$\text{minpoly}_\alpha(x) = \prod_{\sigma} (x - \sigma(\alpha)) = x^n + a_2(\alpha)x^{n-2} + \cdots + a_n(\alpha),$$

with  $a_i(\alpha) \in \mathbb{Z}, \quad |a_i(\alpha)| \ll |\text{Disc}(K)|^{\frac{i}{2n-2}}.$

# Schmidt's proof

- ▶ By Minkowski's theory, there exists  $\alpha \in \mathcal{O}_K$  with trace 0 and  $||\alpha||_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$  for all embeddings  $\sigma : K \hookrightarrow \mathbb{C}$ .
- ▶ Assume that  $\mathbb{Q}(\alpha) = K$ . (If not, induct.)
- ▶ The minimal polynomial of  $\alpha$  is

$$\text{minpoly}_\alpha(x) = \prod_{\sigma} (x - \sigma(\alpha)) = x^n + a_2(\alpha)x^{n-2} + \cdots + a_n(\alpha),$$

with  $a_i(\alpha) \in \mathbb{Z}$ ,  $|a_i(\alpha)| \ll |\text{Disc}(K)|^{\frac{i}{2n-2}}$ .

- ▶ There are at most  $O(X^{\frac{d+2}{4}})$  possibilities.

# Schmidt's proof

- ▶ By Minkowski's theory, there exists  $\alpha \in \mathcal{O}_K$  with trace 0 and  $||\alpha||_\sigma \ll |\text{Disc}(K)|^{\frac{1}{2n-2}}$  for all embeddings  $\sigma : K \hookrightarrow \mathbb{C}$ .
- ▶ Assume that  $\mathbb{Q}(\alpha) = K$ . (If not, induct.)
- ▶ The minimal polynomial of  $\alpha$  is

$$\text{minpoly}_\alpha(x) = \prod_{\sigma} (x - \sigma(\alpha)) = x^n + a_2(\alpha)x^{n-2} + \cdots + a_n(\alpha),$$

with  $a_i(\alpha) \in \mathbb{Z}$ ,  $|a_i(\alpha)| \ll |\text{Disc}(K)|^{\frac{i}{2n-2}}$ .

- ▶ There are at most  $O(X^{\frac{d+2}{4}})$  possibilities.

Given a base number field  $K$ .

Given a base number field  $K$ .

## Theorem (Class Field Theory)

*Abelian extensions  $L/K$  of degree  $d$  and conductor dividing  $\mathfrak{m}$  are in bijection with index  $d$  quotients of  $\text{Cl}_{\mathfrak{m}}(L)$ .*

Given a base number field  $K$ .

## Theorem (Class Field Theory)

*Abelian extensions  $L/K$  of degree  $d$  and conductor dividing  $\mathfrak{m}$  are in bijection with index  $d$  quotients of  $\text{Cl}_{\mathfrak{m}}(L)$ .*

## Theorem (Kummer Theory)

*If in addition  $\mu_d \subseteq K$ , then abelian extensions  $L/K$  of exponent  $d$  are in bijection with subgroups of  $K^{\times}/(K^{\times})^d$ .*

# Cyclic cubic fields

## Theorem (Cohn, 1954)

*We have*

$$\sum_{K \text{ cyclic cubic}} \frac{1}{\text{Disc}(K)^s} = -\frac{1}{2} + \frac{1}{2} \left( 1 + \frac{1}{3^{4s}} \right) \prod_{p \equiv 1 \pmod{6}} \left( 1 + \frac{2}{p^{2s}} \right).$$

## Corollary

*We have*

$$N_3(X, C_3) \sim \frac{11\sqrt{3}}{36\pi} \prod_{p \equiv 1 \pmod{6}} \frac{(p+2)(p-1)}{p(p+1)}.$$



# General abelian number fields

Theorem (Wright, Mäki, but read Wood's treatment)

Let  $G$  be any abelian group of order  $n$ . Then we have

$$\sum_{\text{Gal}(K/\mathbb{Q}) \simeq G} \frac{1}{\text{Disc}(K)^s} = \text{finite sum of Euler products} .$$

Corollary

We have

$$N_{|G|}(X, G) \sim c(G) X^{1/a(G)} (\log X)^{b(G)},$$

where  $a(G)$  and  $b(G)$  are explicit and  $c(G)$  is 'explicit'.

# Prime degree (Cohen, Diaz y Diaz, Olivier 2002)

*"It is claimed that this constant can be explicitly computed as a finite product of local adelic integrals, but in practice this has not been done, even for the simplest Abelian groups  $G$ , except for  $G = C_2$ ..."*

# Prime degree (Cohen, Diaz y Diaz, Olivier 2002)

*"It is claimed that this constant can be explicitly computed as a finite product of local adelic integrals, but in practice this has not been done, even for the simplest Abelian groups  $G$ , except for  $G = C_2 \dots$ "*

Cohen, Diaz y Diaz and Olivier, Cyclic extensions 171

**Theorem 1.1.** *Let  $K$  be a number field of signature  $(r_1, r_2)$ . Let  $\mathcal{R}$  (resp.  $\mathcal{D}$ ) be the set of prime ideals of  $K$  which are ramified (resp. totally split) in  $K_z/K$ . Then*


$$N_{K, \ell}(C_\ell, X^{\ell-1}) \sim c_1 c_2 c_3 c_4 X \log^{q_z-1} X$$

with

$$c_1 = \frac{\left( \prod_{d|d_z} \zeta_{K_z[d]}(d)^{\mu(d)} \right)^{q_z}}{d_z^{\ell r_2 + r_z} q_z!},$$
$$c_2 = \prod_{\mathfrak{p} \in \mathcal{D}} \left( \left( 1 + \frac{\ell-1}{\mathcal{N}\mathfrak{p}} \right) \prod_{d|d_z} \left( 1 - \frac{1}{\mathcal{N}\mathfrak{p}^d} \right)^{(\ell-1)\mu(d)/d} \right),$$
$$c_3 = \left( \prod_{\mathfrak{p} \in \mathcal{R}} \prod_{d|d_z} \left( 1 - \frac{1}{\mathcal{N}\mathfrak{p}^{df}(\mathfrak{p}_d/\mathfrak{p})} \right)^{g(\mathfrak{p}_d/\mathfrak{p})\mu(d)} \right)^{q_z},$$
$$c_4 = \prod_{\mathfrak{p} \nmid \ell, \mathfrak{p} \notin \mathcal{D}} \left( 1 + \frac{\ell-1}{\mathcal{N}\mathfrak{p}} - \frac{\ell-1-r(e(\mathfrak{p}))(1-1/\mathcal{N}\mathfrak{p})}{\mathcal{N}\mathfrak{p}^{\lceil e(\mathfrak{p})/(\ell-1) \rceil}} \right),$$

where  $r_z = 0$  if  $\zeta_\ell \in K$ , while  $r_z = r_1 - 1$  otherwise, and by abuse of notation, for any number field  $L$  we write  $\zeta_L(1)$  for the residue of the Dedekind zeta function  $\zeta_L(s)$  at  $s = 1$ .

# The parametrization method



# Intersections of conics

Example. Solve  $x^4 - x^3 + 3x^2 - 5x + 1 = 0$ .

# Intersections of conics


Example. Solve  $x^4 - x^3 + 3x^2 - 5x + 1 = 0$ .

$$y = x^2, \quad y^2 - yx + 3y - 5x + 1 = 0.$$

# Intersections of conics

Example. Solve  $x^4 - x^3 + 3x^2 - 5x + 1 = 0$ .

$$y = x^2, \quad y^2 - yx + 3y - 5x + 1 = 0.$$



# A sample theorem

Theorem (Bhargava, *Annals*, 2004)

*There exists an explicit, discriminant preserving bijection between the following two sets:*



## Theorem (Bhargava, *Annals*, 2004)

*There exists an explicit, discriminant preserving bijection between the following two sets:*

- ▶  $\mathrm{GL}_3(\mathbb{Z}) \times \mathrm{GL}_2(\mathbb{Z})$ -orbits on the lattice  $(\mathrm{Sym}^2 \mathbb{Z}^3 \otimes \mathbb{Z}^2)$  of pairs of integral ternary quartic forms.

## Theorem (Bhargava, *Annals*, 2004)

*There exists an explicit, discriminant preserving bijection between the following two sets:*

- ▶  $\mathrm{GL}_3(\mathbb{Z}) \times \mathrm{GL}_2(\mathbb{Z})$ -orbits on the lattice  $(\mathrm{Sym}^2 \mathbb{Z}^3 \otimes \mathbb{Z}^2)$  of pairs of integral ternary quartic forms.
- ▶ Pairs  $(Q, R)$ , where  $Q$  is a quartic ring and  $R$  is a cubic resolvent of  $Q$ .

# A Bhargava-style metatheorem

## Theorem

*There exists an explicit, discriminant preserving bijection between the following two sets:*

# A Bhargava-style metatheorem

## Theorem

*There exists an explicit, discriminant preserving bijection between the following two sets:*

- ▶  *$G(\mathbb{Z})$ -orbits on a lattice  $V(\mathbb{Z})$ ; where  $G$  is an **algebraic group** acting (often **prehomogeneously**) on a vector space  $V$ ;*

# A Bhargava-style metatheorem

## Theorem

*There exists an explicit, discriminant preserving bijection between the following two sets:*

- ▶  $G(\mathbb{Z})$ -orbits on a lattice  $V(\mathbb{Z})$ ; where  $G$  is an **algebraic group** acting (often **prehomogeneously**) on a vector space  $V$ ;
- ▶ Some nice class of arithmetic objects we want to count.

# How to prove Bhargava-style theorems

# How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration.

# How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration. (Or Omar Khayyam!)



# How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration. (Or Omar Khayyam!)
- ▶ Pick your favorite complex representation  $(G, V)$  (which should be defined over  $\mathbb{Z}$ , and for which the invariant theory should be nice).

# How to prove Bhargava-style theorems

- ▶ Read old papers in representation theory, invariant theory, and commutative algebra for inspiration. (Or Omar Khayyam!)
- ▶ Pick your favorite complex representation  $(G, V)$  (which should be defined over  $\mathbb{Z}$ , and for which the invariant theory should be nice).
- ▶ Try to prove that the  $G(\mathbb{Z})$ -orbits on  $V(\mathbb{Z})$  parametrize something. Hope to get lucky.

# Counting Low Degree Fields

Theorem (Davenport-Heilbronn, Bhargava, et al.)

*We have*

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3}(\log X)^{2.09}),$$

# Counting Low Degree Fields

Theorem (Davenport-Heilbronn, Bhargava, et al.)

*We have*

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3}(\log X)^{2.09}),$$

$$N_4(X, S_4) \sim \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})X,$$

# Counting Low Degree Fields

Theorem (Davenport-Heilbronn, Bhargava, et al.)

*We have*

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3}(\log X)^{2.09}),$$

$$N_4(X, S_4) \sim \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})X,$$

$$N_5(X) \sim \frac{13}{130} \prod_p (1 + p^{-2} - p^{-4} - p^{-5})X.$$

# Counting Low Degree Fields

Theorem (Davenport-Heilbronn, Bhargava, et al.)

*We have*

$$N_3(X) = \frac{1}{3\zeta(3)}X + \frac{4(1 + \sqrt{3})\zeta(1/3)}{5\Gamma(2/3)^3\zeta(5/3)}X^{5/6} + O(X^{2/3}(\log X)^{2.09}),$$

$$N_4(X, S_4) \sim \frac{5}{24} \prod_p (1 + p^{-2} - p^{-3} - p^{-4})X,$$

$$N_5(X) \sim \frac{13}{130} \prod_p (1 + p^{-2} - p^{-4} - p^{-5})X.$$

These are now **lattice point** counting problems.

# Inductive Methods (New Results From Old)

## Theorem (Cohen, Diaz y Diaz, Olivier)

*We have*

$$N_4(X, D_4) \sim X \cdot \frac{3}{\pi^2} \sum_D \frac{2^{-r_2(D)}}{D^2} \frac{L(1, D)}{L(2, D)},$$

*where the sum ranges over all fundamental discriminants  $\neq 1$ .*

# Inductive Methods (New Results From Old)

## Theorem (Cohen, Diaz y Diaz, Olivier)

*We have*

$$N_4(X, D_4) \sim X \cdot \frac{3}{\pi^2} \sum_D \frac{2^{-r_2(D)}}{D^2} \frac{L(1, D)}{L(2, D)},$$

*where the sum ranges over all fundamental discriminants  $\neq 1$ .*



# Counting by other invariants

## Theorem (Belabas-Fouvry, Bhargava-Wood)

*We have*

$$N_6(X, S_3) \sim \frac{2}{9} \left( \frac{4}{3} + \frac{1}{3^{5/3}} + \frac{2}{3^{7/3}} \right) \prod_{p \neq 3} (1 + p^{-1} + p^{-4/3}) \cdot X^{1/3}.$$

# Counting by other invariants

## Theorem (Belabas-Fouvry, Bhargava-Wood)

*We have*

$$N_6(X, S_3) \sim \frac{2}{9} \left( \frac{4}{3} + \frac{1}{3^{5/3}} + \frac{2}{3^{7/3}} \right) \prod_{p \neq 3} (1 + p^{-1} + p^{-4/3}) \cdot X^{1/3}.$$

**Idea:** If  $K$  is an  $S_3$ -cubic with  $\text{Disc}(K) = Dn^2$ , then  $\text{Disc}(\tilde{K}) = D^3 n^4$  apart from the 2- and 3-adic factors.

## Theorem (Wang, Masri-T.-Tsai-Wang)

*Let  $d \in \{3, 4, 5\}$  and let  $A$  be any abelian group. Then*

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A)X^{1/|A|}.$$

## Theorem (Wang, Masri-T.-Tsai-Wang)

Let  $d \in \{3, 4, 5\}$  and let  $A$  be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A)X^{1/|A|}.$$

**Idea:** If  $K$  is an  $S_d$ -field and  $L$  is an  $A$ -field, then **usually**  $K$  and  $L$  are linearly disjoint with

## Theorem (Wang, Masri-T.-Tsai-Wang)

Let  $d \in \{3, 4, 5\}$  and let  $A$  be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A) X^{1/|A|}.$$

**Idea:** If  $K$  is an  $S_d$ -field and  $L$  is an  $A$ -field, then **usually**  $K$  and  $L$  are linearly disjoint with

$$\text{Disc}(KL) = * \cdot \text{Disc}(K)^{|A|} \text{Disc}(L)^d,$$

## Theorem (Wang, Masri-T.-Tsai-Wang)

Let  $d \in \{3, 4, 5\}$  and let  $A$  be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A) X^{1/|A|}.$$

**Idea:** If  $K$  is an  $S_d$ -field and  $L$  is an  $A$ -field, then **usually**  $K$  and  $L$  are linearly disjoint with

$$\text{Disc}(KL) = * \cdot \text{Disc}(K)^{|A|} \text{Disc}(L)^d,$$

where  $*$  is divisible only primes **ramified in both  $K$  and  $L$** .

## Theorem (Wang, Masri-T.-Tsai-Wang)

Let  $d \in \{3, 4, 5\}$  and let  $A$  be any abelian group. Then

$$N_{d|A|}(X, S_d \times A) \sim c(S_d \times A) X^{1/|A|}.$$

**Idea:** If  $K$  is an  $S_d$ -field and  $L$  is an  $A$ -field, then **usually**  $K$  and  $L$  are linearly disjoint with

$$\text{Disc}(KL) = * \cdot \text{Disc}(K)^{|A|} \text{Disc}(L)^d,$$

where  $*$  is divisible only primes **ramified in both  $K$  and  $L$** .

(This doesn't happen too often.)

Theorem (Klüners + $\epsilon$ )

*Assume* a 'weak Malle conjecture' of the form

$$N_d(X, G) \ll X^{3/2}.$$



Theorem (Klüners + $\epsilon$ )

*Assume a 'weak Malle conjecture' of the form*

$$N_d(X, G) \ll X^{3/2}.$$

*Then we have*

$$N_{2d}(X, C_2 \wr G) \sim c(C_2 \wr G)X.$$

Theorem (Klüners + $\epsilon$ )

*Assume a 'weak Malle conjecture' of the form*

$$N_d(X, G) \ll X^{3/2}.$$

*Then we have*

$$N_{2d}(X, C_2 \wr G) \sim c(C_2 \wr G)X.$$

**Idea.** A quadratic extension of a  $G$ -extension usually has Galois group  $C_2 \wr G$ .

# Wreath products

Theorem (Klüners + $\epsilon$ )

*Assume a 'weak Malle conjecture' of the form*

$$N_d(X, G) \ll X^{3/2}.$$

*Then we have*

$$N_{2d}(X, C_2 \wr G) \sim c(C_2 \wr G)X.$$

**Idea.** A quadratic extension of a  $G$ -extension usually has Galois group  $C_2 \wr G$ .

**Note.**  $D_4 \simeq C_2 \wr C_2$ ; subsumes Cohen-Diaz-Olivier as a special case.

## Theorem (Alberts, 2018)

Assume that *“the  $m$ -torsion in class groups is small on average”*.  
Then, for every solvable transitive subgroup  $G \subseteq S_d$  we have

$$N_d(X, G) \ll X^{1/a(G)+\epsilon}.$$

## Theorem (Alberts, 2018)

Assume that *“the  $m$ -torsion in class groups is small on average”*.  
Then, for every solvable transitive subgroup  $G \subseteq S_d$  we have

$$N_d(X, G) \ll X^{1/a(G)+\epsilon}.$$

Further development of the same family of ideas.

## Theorem (Alberts, 2018)

Assume that *“the  $m$ -torsion in class groups is small on average”*.  
Then, for every solvable transitive subgroup  $G \subseteq S_d$  we have

$$N_d(X, G) \ll X^{1/a(G)+\epsilon}.$$

Further development of the same family of ideas.

**Also:** See further related works by [Altuğ](#), [Lemke Oliver](#), [Mehta](#), [Shankar](#), [Taniguchi](#), [Varma](#), [Wilson](#), and previously named authors (in various permutations).

Happy Holidays!