

Curried Functional Spaces


Roman Maksimovich

Hong Kong University of Science and Technology

May 10, 2026

Premise


Multivariable functions



Multivariate calculus

Curried functions

$$\mathbb{R} \rightarrow (\mathbb{R} \rightarrow (\dots (\mathbb{R} \rightarrow \mathbb{R})))$$




λ -calculus, functional programming

Premise

Curried functions

$$\mathbb{R} \rightarrow (\mathbb{R} \rightarrow (\dots (\mathbb{R} \rightarrow \mathbb{R})))$$



λ -calculus,
functional programming

Multivariate calculus

Continuous Functions

$C_n(\mathbb{R})$ — the space of continuous n -ary curried functions:

1. $C_0(\mathbb{R}) := \mathbb{R}$


2. $C_{n+1}(\mathbb{R}) := \{f: \mathbb{R} \rightarrow C_n(\mathbb{R}) \mid f \text{ is continuous}\}$

Continuous Functions

$C_n(\mathbb{R})$ — the space of continuous n -ary curried functions:

1. $C_0(\mathbb{R}) := \mathbb{R}$

2. $C_{n+1}(\mathbb{R}) := \{f: \mathbb{R} \rightarrow C_n(\mathbb{R}) \mid f \text{ is } \underline{\text{continuous}}\}$



⇒ what is the topology on $C_n(\mathbb{R})$?


Continuous Functions

$C_n(\mathbb{R})$ — the space of continuous n -ary curried functions:

1. $C_0(\mathbb{R}) := \mathbb{R}$

2. $C_{n+1}(\mathbb{R}) := \{f: \mathbb{R} \rightarrow \underline{C_n(\mathbb{R})} \mid f \text{ is continuous}\}$

⇒ compact-open topology: $C_n(\mathbb{R}) = \mathcal{C}(\mathbb{R}, C_{n-1}(\mathbb{R}))$



Topological Vector Spaces


Remark: If $f_1, f_2 \in C_n(\mathbb{R})$, then $\alpha f_1 + \beta f_2 \in C_n(\mathbb{R})$.

Topological Vector Spaces

Remark: If $f_1, f_2 \in C_n(\mathbb{R})$, then $\alpha f_1 + \beta f_2 \in C_n(\mathbb{R})$.


Definition: X is a **TVS** $\Leftrightarrow X$ has **topological** and **vector** structure, and they are *compatible*:

1. $(+): X \times X \rightarrow X$ is continuous;
2. $(\cdot): \mathbb{R} \times X \rightarrow X$ is continuous.




Properties of Topological Vector Spaces


Metrizability



Local convexity




Completeness




Fréchet space

Properties of Topological Vector Spaces


Metrizability



Local convexity



Completeness



Fréchet space



Curried function spaces $C_n(\mathbb{R})$

Continuous Functions

$C_n(\mathbb{R})$ — the space of continuous n -ary curried functions:

1. $C_0(\mathbb{R}) := \mathbb{R}$;
 2. $C_{n+1}(\mathbb{R}) := \mathcal{C}(\mathbb{R}, C_n(\mathbb{R}))$, with the compact-open topology.
-

Results:

1. $C_n(\mathbb{R})$ is isomorphic to $\mathcal{C}(\mathbb{R}^n, \mathbb{R})$.
 2. $C_n(\mathbb{R})$ is a topological vector space.
 3. $C_n(\mathbb{R})$ is a Fréchet space.
-

$\Rightarrow$ Theorems from functional analysis apply:

1. Uniform boundedness principle;
2. Open mapping theorem;
3. ...

Differentiable functions

$D_n^m(\mathbb{R})$ — the space of n -ary m times differentiable functions:

1. $D_0^m(\mathbb{R}) := \mathbb{R};$

2. $D_n^0(\mathbb{R}) := C_n(\mathbb{R});$

3. If $n, m > 0$, then $D_n^m(\mathbb{R})$ consists of all such $f \in \mathcal{C}(\mathbb{R}, D_{n-1}^m(\mathbb{R}))$, that there is $f' \in D_n^{m-1}(\mathbb{R})$ with

$$\frac{f(x+h) - f(x)}{h} \xrightarrow{h \rightarrow 0} f'(x).$$

Differentiable functions


$D_n^m(\mathbb{R})$ — the space of n -ary m times differentiable functions:

1. $D_0^m(\mathbb{R}) := \mathbb{R};$

2. $D_n^0(\mathbb{R}) := C_n(\mathbb{R});$

3. If $n, m > 0$, then $D_n^m(\mathbb{R})$ consists of all such $f \in \mathcal{C}(\mathbb{R}, D_{n-1}^m(\mathbb{R}))$, that there is $f' \in D_n^{m-1}(\mathbb{R})$ with

$$\frac{f(x+h) - f(x)}{h} \xrightarrow{h \rightarrow 0} f'(x).$$



pointwise convergence!

$$f_k \xrightarrow{k \rightarrow \infty} f \Leftrightarrow f_k(x_1) \dots (x_n) \xrightarrow{k \rightarrow \infty} f(x_1) \dots (x_n)$$

Differentiable functions

$D_n^m(\mathbb{R})$ — the space of n -ary m times differentiable functions:

1. $D_0^m(\mathbb{R}) := \mathbb{R}$;
2. $D_n^0(\mathbb{R}) := C_n(\mathbb{R})$;
3. If $n, m > 0$, then $D_n^m(\mathbb{R})$ consists of all such $f \in \mathcal{C}(\mathbb{R}, D_{n-1}^m(\mathbb{R}))$, that there is $f' \in D_n^{m-1}(\mathbb{R})$ with

$$\frac{f(x+h) - f(x)}{h} \xrightarrow{h \rightarrow 0} f'(x).$$

The topology on D_n^m is *coarsest* such that the maps

$$\begin{array}{ccc} i: D_n^m(\mathbb{R}) \rightarrow \mathcal{C}(\mathbb{R}, D_{n-1}^m(\mathbb{R})), & d: D_n^m(\mathbb{R}) \rightarrow D_n^{m-1}(\mathbb{R}) \\ f \mapsto f & f \mapsto f' \end{array}$$