



# Master Thesis

Thorvald Demuth Jørgensen

## Provability Logic

An Investigation of the Relationship Between Modal Logic and  
Arithmetics

Date: December 2, 2022

Advisor: Asger Dag Törnquist & Rasmus Kræmmer Rendsvig



## Abstract

# Contents

|  |           |
|--|-----------|
| <b>0. Introduction</b>   | <b>1</b>  |
| 0.1. Notation . . . . .  | 2         |
| 0.2. Historical Introduction . . . . .                                 | 3         |
| 0.2.1. Modal Logic . . . . .   | 3         |
| 0.2.2. Arithmetics and Recursion Theory . . . . .                      | 4         |
| 0.2.3. Provability Logic . . . . .                                     | 5         |
| <b>1. Preliminary</b>  | <b>7</b>  |
| 1.1. Modal Logic . . . . .   | 7         |
| 1.1.1. The language of modal logic . . . . .                           | 7         |
| 1.1.2. Semantics . . . . .   | 8         |
| 1.1.3. The modal logic GL . . . . .                                    | 10        |
| 1.2. Gödel's Incompleteness Theorems . . . . .                         | 12        |
| 1.2.1. Primitive Recursive Functions . . . . .                         | 12        |
| 1.2.2. <b>PRA</b> . . . . .  | 14        |
| 1.2.3. Arithmetization of the syntax . . . . .                         | 17        |
| 1.2.4. The proof predicate and the theorems . . . . .                  | 18        |
| <b>2. Recursion Theory</b>   | <b>21</b> |
| 2.1. Partial Recursive Functions and Recursive Functions . . . . .     | 21        |
| 2.2. Turing Computable Functions . . . . .                             | 22        |
| 2.3. The $s$ - $m$ - $n$ -Theorem . . . . .                            | 24        |
| 2.4. Recursively Enumerable sets and the Graph of a function . . . . . | 27        |
| 2.5. The Recursion Theorem . . . . .                                   | 30        |
| 2.5.1. Application of the Recursion Theorem . . . . .                  | 31        |
| <b>3. The Arithmetical Hierarchy and Fragments of Arithmetics</b>      | <b>33</b> |
| 3.1. The arithmetical hierarchy . . . . .                              | 33        |
| 3.2. Fragments of arithmetic . . . . .                                 | 36        |
| 3.3. <b>PRA</b> and the other fragments of Arithmetics . . . . .       | 38        |
| 3.3.1. Induction In <b>PRA</b> . . . . .                               | 39        |
| 3.4. Further extensions of the arithmetical hierarchy . . . . .        | 40        |
| 3.5. Sigma Completeness . . . . .                                      | 41        |

|  |           |
|--|-----------|
| <b>4. General Results on GL</b>                              | <b>43</b> |
| 4.1. Tress and <b>GL</b>                                     | 43        |
| 4.1.1. Trees   | 43        |
| 4.1.2. The Finite Tree Theorem                               | 46        |
| 4.2. The Continuity Theorem                                  | 48        |
| 4.3. The Modal Logic <b>GLS</b>                              | 49        |
| <b>5. Fixed Point Theorem</b>                                | <b>55</b> |
| 5.1. Prerequisite Results and Definitions                    | 55        |
| 5.2. Proof of the Fixed Point Theorem                        | 58        |
| 5.3. Calculation of Fixed Points                             | 62        |
| <b>6. Solovays Completeness Theorems</b>                     | <b>67</b> |
| 6.1. Soundness   | 67        |
| 6.2. The First Theorem                                       | 68        |
| 6.3. The Second Theorem                                      | 77        |
| 6.3.1. Using the Second Completeness Theorem                 | 80        |
| 6.4. Generalisations of the Completeness Theorems            | 81        |
| 6.5. Solovays Completeness Theorems and Fixed Points         | 81        |
| 6.6. Concluding remarks: Implications of Solovay's Theorems  | 82        |
| <b>7. Further Results in Provability Logic</b>               | <b>85</b> |
| 7.1. Multi-modal provability logic                           | 85        |
| 7.1.1. The system <b>GLB</b>                                 | 85        |
| 7.1.2. The system <b>IDzh</b> and completeness of <b>GLB</b> | 88        |
| 7.1.3. The system <b>GLP</b>                                 | 90        |
| 7.2. Quantified provability logic                            | 91        |
| <b>8. Bibliography</b>                                       | <b>93</b> |
| <b>A. Necessitation free definition of GL and GLS</b>        | <b>97</b> |

# 0. Introduction

If modern modal logic was  
conceived in sin, then it has been  
redeemed through Gödliness

---

George. S Boolos 1979 page 1

{chap:intro}

This project is about the subject provability logic from mathematical logic, and the main result will be Solovay's completeness theorems. These theorems states that the  $\Box$ -operator in modal logic **GL** axiomatizes the proof predicate  $\text{Pr}(\cdot)$  from a wide range of fragments of arithmetics, and that the modal logic **GLS** axiomatizes the proof predicate of true arithmetics. For proving these theorems we will need some results from recursion theory and general knowledge about fragments of arithmetics and the modal logics **GL** and **GLS**. We will further prove the fixed point theorem for **GL**. The project has the following structure:

- Chapter 1:** This chapter will list some prerequisite results and definitions about the modal logic **GL** and the arithmetical system primitive recursive arithmetic.
- Chapter 2:** In this chapter there will be a short introduction to recursion theory and the main goal is to show the recursion theorem.
- Chapter 3:** Here we will introduce the arithmetical hierarchy and fragments of arithmetics. We will further comment on the amount of induction we have available in primitive recursive arithmetics
- Chapter 4:** We will start of by strengthening the completeness theorem for **GL**, with the finite tree completeness theorem. Further we will look at an expansions of **GL** known as **GLS**, which we will use later on for Solovay's second completeness theorem.
- Chapter 5:** The main goal of this chapter is to state and prove the fixed point theorem. We will also use it to calculate a few fixed points.
- Chapter 6:** In this chapter we will prove both of Solovay's completeness theorems. After the proofs of these we will look at applications of them.

**Chapter 7:** Here we will round up the project with some perspective to further results in provability logic.

In the rest of this introduction we will introduce a bit of notational conventions and give a short historical overview of the different logical fields, that will play a role in the project.

The bibliography of this project consist of a wide range of logical, mathematical and philosophical texts that in one way or another has a connection to of is referenced in the project. The main sources for this project is (Smorynski 1985), (George. S Boolos 1993) and (Soare 1987).

### 0.1. Notation

We denote the non-negative integers with  $\omega$ ; i.e  $\omega = \{0, 1, 2, 3, \dots\}$ , and  $\omega \times \omega = \omega^2$  and so on. Subsets of  $\omega$  will be denoted by  $A, B, C \dots$  and arbitrary sets will be denoted by  $\Gamma, \Phi, \Theta \dots$ . Lower case Latin letter  $a, b, c$  and  $x, y, z$  will denote integers, and  $\vec{x}$  will be shorthand for a finite sequences, e.g.  $x_1, \dots, x_n$ . Further recursive functions will be denoted by  $\varphi, \psi, \vartheta, \dots$ , the graphs of recursive functions will be denoted by  $\tau, \xi$  and  $\zeta$ . Functions that are primitive recursive will be denoted by  $f, g, h, \dots$ .

We will further introduce a system of arithmetic called *primitive recursive arithmetic* (**PRA**, for short) and the formulas of this system will be denoted with  $F, G$  and  $G$ .

For a partial function  $\varphi(x) \downarrow$  will denote that  $\varphi(x)$  is defined and  $\varphi(x) \downarrow = y$  denotes that  $\varphi(x)$  is defined and has value  $y$ .  $\varphi(x) \uparrow$  denotes that  $\varphi(x)$  is undefined.  $\text{dom}(\varphi)$  and  $\text{im}(\varphi)$  denotes the domain and image of  $\varphi(x)$ . A few special functions have their own symbols:  $S$  for the successor function,  $Z$  for the zero function and  $P$  for the projection. The formulae of the language **PRA** will be denoted by  $F, G$  and  $H$ .

Relations will be denoted by  $R$  and  $<, >, \leq, \geq$  will be used in the usual sense on integers.

We will use the standard symbols of propositional logic  $\wedge, \vee, \rightarrow, \neg$ . Further we will sometimes use  $\exists$  and  $\forall$  in the metalanguage of modal logic, and use them in the language of first-order arithmetics. The differences of these two uses should be clear from the context. We will argument the propositional logic with the modal unary connective  $\Box$  and its dual  $\Diamond := \neg\Box\neg$ . We will denote modal formulas by the greek letters  $\alpha, \beta, \gamma, \sigma$ . The symbol  $\mathcal{H}$  will denote a Hintikka frame which is a tuple  $\langle W, R \rangle$  where  $W$  is a set of "nodes" and  $R$  is a relation on  $W$  and  $\mathcal{K}$  will denote a Kripke model, which is a Hintikka frame with a valuation  $\phi$  on.

## 0.2. Historical Introduction

Humans have been interested in numbers and calculation with these since before the beginning of history; i.e arithmetics.

### 0.2.1. Modal Logic

The Greek philosopher Aristotle did try to develop some kind of modal logic about necessary and possibility in both his logical work *De Interpretatione* and in his *Metaphysic*.

With our modern eyes his thoughts about necessitation and possibility seems a bit confusing. But he gets some implications about these notions right. A good overview of this part of his logical work can be found in (Lemmon 1977) and in (Łukasiewicz 1957). In the late antiquity and the middle ages there was done some work building on Aristotle earlier work.

The next big step for modal logic came in the enlightenment when the rationalist philosopher Gottfried W. Leibniz came up with the idea of possible worlds. Leibniz thought that we live in the best possible world, since it is this world God has chosen to create. With this idea of a metaphysical possible worlds, it was possible to make a clear definition of when a proposition is necessary true; and a proposition is necessary true, if it is true in all possible worlds. But Leibniz did not any further works on modal logic as a whole.

Modern modal logic is said to be started by Clarence Irving Lewis. He set out to develop a formal system without the paradoxes of material implication. This led to his development of a wide range of different modal logical system in the first part of the 20th century. His method was syntactical, since there were yet to be develop a "smart" semantic for modal logic. But in the start of the 20th century, C. I. Lewis had started the modern modal logic, and a lot of other philosophers and logicians began to have an interest in the subject.

In the 1950s there had been a lot of research into the syntax of modal logic. But there was still not a clear definition of the semantics of modal logic, even though philosophers and logicians still had the intuitive definition of necessary true from Leibniz. The answer came in the late 1950s, when the philosophers and logicians Jaako Hintikka and Saul Kripke developed the concept of a Kripke model.<sup>1</sup> Hintikka came up with the idea of a frame; i.e a pair  $\langle W, R \rangle$  where  $W$  is a set of worlds and  $R$  is an arbitrary relation on these. With this construction it was possible to explain for a given world, what other worlds was accessible to it; i.e possible worlds for it. Kripke came up with the same idea, but added a valuation function to the frame, which to each formula of the language ascribed a

---

<sup>1</sup>They were not alone in this development. But for simplicity sake **ja**

## 0. Introduction

---

set of worlds in which it was true. This construction made it possible for Kripke to prove some completeness results about modal logics.

When the concept of the Kripke model became wider known, it started a *Golden* age of research in modal logic. A lot of completeness results about different modal logic was proven in the 1960s and early 1970s; including the weak completeness theorem for the logic that would later be known as provability logic; this was done by Krister Segerberg without him having knowledge of what the interpretation of this logic could be.

A deeper look into the development of modal logic can be found in (Goldblatt 2003).

### 0.2.2. Arithmetics and Recursion Theory

Even though humans have been using arithmetics for millennia, it had not been axiomatized.

It was first in the second half of the 1800s there was a development in the axiomatization of arithmetics. Richard Dedekind and Giuseppe Peano developed axioms system for arithmetics, these was later evolved into what today is known as the Peano axioms for arithmetics.

Another important achievement at the turn of the century, was David Hilbert's proof that the consistency of Euclidean geometry could be proven by proving that arithmetics was consistent. **Hans geometri værk** This was one of the first steps in what later would be known as the Hilbert program. The Hilbert program was not really a program until the end of 1920, where it became clear that its goal was to prove that the *ideal* transfinite mathematics was consistent, and that the proof of this should be conducted in a *real* finite mathematics system.

An example of such a finite system could be the *Primitive recursive arithmetics* that was developed by Toralf Skolem; the definition of this system will be given in the next chapter. This system can be seen as a (induction-wise) weaker version of Peano arithmetics. Tait argues that such a system fulfills the finitist conditions of the Hilbert program. This system will be the main system of this project and the definition of it will come in the next chapter.

Another (mostly historically) important mathematical system of arithmetics was the one developed by Bertrand Russell and Alfred North Whitehead in their work: *Principia Mathematica*. The goal of this work was somewhat different than the one of the Hilbert program. Russell had a thesis that shortly (and not the most precis) says that all of mathematics could be reduce to logic. In Kantian terms he thought that mathematics was analytical *a priori*.

In 1931 Kurt Gödel's paper: *Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme I* it was shown that the system of Russell and Whitehead was not complete, and under the assumption that the system



was consistent that it could not prove its own consistency. This result can be generalized to systems strong enough to do simple arithmetic. The original proof was build on the system of Russel and Whitehead and had a assumption of  $\omega$ -consistency. This was a blow to the Hilbert program, since the result showed that the goal of it could not be fulfilled. Later on John Barkley Rosser in his article (Rosser 1936) improved the result so that it consistency was enough.

Later on in the 1930s Gödel extended the primitive recursive function to the recursive functions. A few different definitions of intuitively computable functions was set forward by Kleene, Turing and Church and it was shown that all these different notions of intuitively computable functions gave rise to the same class of functions.

This lead Church to state the so called *Church-Turing Thesis*, that states that this class of functions are the computable functions.

### 0.2.3. Provability Logic

The first time that someone made a connection between modal logic and the proof predicate of some form of arithmetic, was in 1933 when Kurt Gödel published his very short paper: *Eine Interpretation des intuitionistischen Aussagenkalküls*. Here Gödel stated a hypothesis that it was possible to axiomatize the what the intuitionist meant by a proof in a version of modal logic.

After Löb's derivability conditions was found and after the development of the semantics of modal in the 1960s, it became clear that there was a connection between the logic that Segerberg had proven to be weakly complete and the proof predicate from arithmetics.

This lead to research done in different places around the from, from The Netherlands, Italy and the United States. It was clear early on that if  $\alpha$  was a theorem of provability logic, then for all interpretations of  $\alpha$  in a fragment of arithmetics was also a theorem of this fragment; but the other way was harder to show. In the same time there was proven a number of important theorems about provability; one of these, called the Fixed Point Theorem, will be proven in chapter 5. A short overview of this early development can be found in (G. Boolos and Sambin 1991).

It was proven in 1976 by Solovay that the modal logic now known as provability logic, is the logic of the provability predicate in Peano Arithmetics; i.e if for all interpretations of a formula  $\alpha$ , if this interpretation was a theorem of Peano arithmetics, then the formula  $\alpha$  was a theorem of provability logic. This also means that provability logic can be seen as an axiomatization of the proof predicate. Solovays result will be the main result in this project; it will be stated precisely and proven in Chapter 6.

After Solovay's article was published, the research focus of provability logic was to generalize Solovay's result to other fragments of arithmetics and if it did also

## 0. Introduction

---

hold for multi-modal provability logic and quantified provability logic. The results of these results will be commented on in the last part of chapter 6 and in chapter 7

The most of the main results about provability logic was proven in the late 70s, 80s and early 90s. This project will not go beyond these early results.

# 1. Preliminary

In this section I will state a few results and definitions from my two project I have written; one about modal logic and one about Gödel's incompleteness theorems. These results will be given without proof. The proofs can be found in two projects I have earlier written (Jørgensen 2021) and (Jørgensen 2022).

The main points of this section will be the definition of the predicate  $\text{Pr}(\cdot)$  which is constructed in the proof of Gödel's Incompleteness Theorems, and the definition of the modal logic **GL** (The name comes from Gödel and Löb) and a few of the properties of this modal logic. The main result of this project will be Solovay's arithmetic completeness theorem, that simply put states that these two predicates (from different mathematical systems) "behaves" in the same way.

## 1.1. Modal Logic

Modal logic will play an important rule in this project. The main point here, is that provability can be seen as a modality. This will be swon later on in the project.

### 1.1.1. The language of modal logic

In this subsection we will define our language  $\mathcal{L}_\Box$  of modal logic. We will have start of with a set called  $\Phi$ , which consists of propositional letter  $p, q, \dots$ . We can further define the primitive symbols of our modal language  $\mathcal{L}_\Box$ :

**Definition 1.1** The following symbols are the primitive in our language  $\mathcal{L}_\Box$

1. Every letter from the set  $\Phi$ .
2. The logical constants  $\perp$  (zero-ary) and  $\rightarrow$  (binary)
3. The modal operator  $\Box$  (unary).

—

We can now define the well formed formulas of  $\mathcal{L}_\Box$ , or as we will call them for the: *modal formulas*.

## 1. Preliminary

---

**Definition 1.2** We define a *modal formula* recursively in the following way:

- i Each  $p \in \Phi$  is a modal formula
- ii  $\perp$  is a modal formula
- iii If  $\alpha$  and  $\beta$  are modal formulas then so is  $\alpha \rightarrow \beta$ .
- iv If  $\alpha$  is a modal formula then so is  $\Box\alpha$
- v Nothing is a modal formula except as prescribed by (i)-(iv)

—

Further we will define the connectives  $\wedge, \neg, \vee, \leftrightarrow, \Diamond, \Box^n, \Diamond^n$  and  $\top$  in the usual way. We will read  $\Box\alpha$  as " $\alpha$  is provable. Normally in the alethic modal logic the  $\Box$  reads " $\alpha$  is necessarily true. Further the symbol  $\perp$  stands for falsehood. We will read  $\Diamond\alpha$  as " $\alpha$ " is consistent.

### 1.1.2. Semantics

In this section we will define the notation of truth for our modal logic. We will begin by defining a Hintikka frame and a Kripke model

**Definition 1.3 (Hintikka frame)** A *Hintikka frame* is a tuple  $\mathcal{H} = \langle W, R \rangle$  where  $W$  is non-empty set (we will call its members for nodes<sup>1</sup>) and where  $R \subset W \times W$  is a relation that we will call the accessibility relation. We will use the notation  $wRv$  to denote that the note  $w$  sees the note  $v$ ; i.e  $(w, v) \in R$ . —

We do not at the given time give the relation  $R$  any properties. It could be reflexive, transitive or anti-symmetric.

**Definition 1.4 (Kripke model)** A *Kripke model* is a tuple  $\mathcal{K} = \langle \mathcal{H}, \phi \rangle$  where  $\mathcal{H}$  is a Hintikka frame and  $\phi$  is a valuation function that to each propositional letter  $p \in \Phi$  assigns a subset  $\phi(p)$  of  $W$ . Formally:

$$\phi : \Phi \rightarrow \mathcal{P}(W)$$

—

---

<sup>1</sup>Normally the members of this set will be called worlds, but this interpretation do not make senses in provability logic

We can now define the notion of truth in a given Kripke model  $\mathcal{K} = \langle \mathcal{H}, \phi \rangle$ . If a modal formula  $\alpha$  is true at a node  $w$  in a Kripke model  $\mathcal{K} = \langle \mathcal{H}, \phi \rangle$  we will write:

$$\models_w^{\mathcal{K}} \alpha$$

This notation is taken from (Lemmon 1977). Another notation instead of  $\models_w^{\mathcal{K}} \alpha$  is  $\mathcal{K}, w \models \alpha$ .

**Definition 1.5 (Truth definition)** We define the notion of *truth* in a Kripke modal  $\mathcal{K} = \langle \mathcal{H}, \phi \rangle$  in the following way:

1. If  $\alpha$  is propositional  $p$  then:

$$\models_w^{\mathcal{K}} \alpha \text{ iff } w \in \phi(p)$$

2. If  $\alpha$  is  $\perp$  then:

$$\models_w^{\mathcal{K}} \alpha \text{ iff never}$$

3. If  $\alpha$  is  $\beta \rightarrow \gamma$  then:

$$\models_w^{\mathcal{K}} \alpha \text{ iff if } \models_w^{\mathcal{K}} \beta \text{ then } \models_w^{\mathcal{K}} \gamma$$

4. if  $\alpha$  is  $\Box\beta$  then:

$$\models_w^{\mathcal{K}} \alpha \text{ iff for all } v \text{ such that } wRv \text{ we have } \models_v^{\mathcal{K}} \beta$$

→

We will end this section with definitions of a *valid* and *satisfiable* formula.

**Definition 1.6** For a given Hintikka frame  $\mathcal{H} = \langle W, R \rangle$  we say that  $\alpha$  is *valid* in  $\mathcal{H}$  and write  $\models^{\mathcal{H}} \alpha$  if and only if  $\models_w^{\mathcal{K}} \alpha$  for all Kripke models on our frame  $\mathcal{H}$  and all nodes  $w \in W$ . Further we say that  $\alpha$  is *satisfiable* in  $\mathcal{H}$  if and only if  $\models_w^{\mathcal{K}} \alpha$  for some Kripke model  $\mathcal{K}$  on the frame  $\mathcal{H}$  and some node  $w \in W$ .

$\alpha$  is called *valid* if and only if  $\alpha$  is valid in all frames  $\mathcal{H}$ , if this is the case we simply write  $\models \alpha$ .  $\alpha$  is called *satisfiable* if and only if  $\alpha$  is satisfiable on all frames  $\mathcal{H}$ .

A formula  $\alpha$  is called valid on a class of frames  $\mathcal{C}$  if for all  $\mathcal{H} \in \mathcal{C}$  we have that  $\models^{\mathcal{H}} \alpha$ . We write this as  $\vdash^{\mathcal{C}} \alpha$ .

A formula  $\alpha$  is *valid* in a Kripke model  $\mathcal{K} = \langle W, R, \phi \rangle$  if and only if  $\models_w^{\mathcal{K}} \alpha$  for all nodes  $w \in W$ . We write this as  $\models^{\mathcal{K}} \alpha$

If a Kripke model  $\mathcal{K}$  has a *minimal node*  $w_0$ , i.e a node such that for all  $v \in W$  we have  $w_0 R v$ , then a modal formula  $\alpha$  is called *true* in  $\mathcal{K}$  if we have that  $\models_{w_0}^{\mathcal{K}} \alpha$ . →

### 1.1.3. The modal logic GL

In this section we will define the notion of a modal logic and define the modal logic **GL**, which will be our base modal logic in this project.

**Definition 1.7** A *modal logic*  $\Lambda$  is a set of modal formulas that contains all propositional tautologies, and is closed under *modus ponens* (MP) and uniform substitution. If  $\alpha \in \Lambda$  we say that  $\alpha$  is a theorem of  $\Lambda$  and write  $\vdash_{\Lambda} \alpha$ , else we write  $\not\vdash_{\Lambda} \alpha$ .

We will also define the set  $\Lambda_S = \{\alpha \models^S \alpha, \text{ for all structures } S \in S\}$ . Where  $S$  is any class of frames.  $\dashv$

We will further define when a given modal formula is deducible in a modal logic:

**Definition 1.8** Let  $\Lambda$  be a modal logic, let  $\beta_1, \dots, \beta_n$  be modal formulas in  $\Lambda$  and let  $\alpha$  be a modal formula. We say that  $\alpha$  is *deducible* from  $\beta_1, \dots, \beta_n$  if  $(\beta_1 \wedge \dots \wedge \beta_n) \rightarrow \alpha$  is a tautology.

If  $\Gamma \cup \{\alpha\}$  is a set of modal formulas, then  $\alpha$  is *deducible* in  $\Lambda$  from  $\Gamma$  if  $\vdash_{\Lambda} \alpha$  or if there are formulas  $\beta_1, \dots, \beta_n \in \Gamma$  such that:

$$\vdash_{\Lambda} (\beta_1 \wedge \dots \wedge \beta_n) \rightarrow \alpha$$

In this case we write  $\Gamma \vdash_{\Lambda} \alpha$  else we write  $\Gamma \not\vdash_{\Lambda} \alpha$   $\dashv$

We will now define what a *normal modal logic* is. We will later look at a modal logic that is not normal. But the modal logic **GL** is a normal one.

**Definition 1.9** A modal logic  $\Lambda$  is called *normal* if it has the following axioms and deduction rules:

Tau All propositional tautologies.

$$\text{K } \Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q)$$

MP  $p, p \rightarrow q \vdash_{\Lambda} q$  (*modus ponens*)

Nec  $p \vdash_{\Lambda} \Box p$  (*necessitation*)

If  $\Gamma$  is a set of modal formulas we call the smallest normal logic containing  $\Gamma$  the normal modal logic axiomatized by  $\Gamma$ . The normal modal logic axiomatized by empty set is called **K** and this is the smallest normal modal logic.<sup>2</sup>  $\dashv$

A few standard results follows from these definitions:

---

<sup>2</sup>Here **K** stands for Kripke

**Proposition 1.1** If  $\Lambda$  is a normal modal logic then:

1. If  $\vdash_{\Lambda} \alpha \rightarrow \beta$  then  $\vdash_{\Lambda} \Box \alpha \rightarrow \Box \beta$
2.  $\vdash_{\Lambda} \Box(\alpha \wedge \beta) \leftrightarrow \Box \alpha \wedge \Box \beta$
3.  $\vdash_{\Lambda} \Box(\alpha_1 \wedge \cdots \wedge \alpha_n) \leftrightarrow \Box \alpha_1 \wedge \cdots \wedge \Box \alpha_n$  for  $n \geq 2$ .
4. If  $\vdash_{\Lambda} \beta_1 \wedge \cdots \wedge \beta_n \rightarrow \alpha$  then  $\vdash_{\Lambda} \Box \beta_1 \wedge \cdots \wedge \Box \beta_n \rightarrow \Box \alpha$ , for  $n \geq 0$ .

It should further be noted that the axiom **K** is equivalent to the following formula:

$$\Box \alpha \wedge \Box(\alpha \rightarrow \beta) \rightarrow \Box \beta$$

Smorynski uses this formula as an axiom instead of the axiom **K**. We can extend the logic **K** in the following way, to get the logic **GL** and the logic **4**.

**Definition 1.10** **K4** is the modal logic extending **K** by adding the following axiom:

$$4 \quad \Box \alpha \rightarrow \Box \Box \alpha$$

**GL** is the modal logic extending **K** by adding the two following axioms:

$$4 \quad \Box p \rightarrow \Box \Box p$$

$$L \quad \Box(\Box p \rightarrow p) \rightarrow \Box p$$

+

**Remark 1.1** Smorynski calls the modal logic **K4** for **BML** (Basic modal logic).

It can be shown that the axiom 4 is redundant. It is now possible to define the notions of soundness, strong completeness and weak completeness.

**Definition 1.11** Let  $S$  be a class of frames or models.

1. A normal modal logic  $\Lambda$  is sound with respect to  $S$  if  $\Lambda \subseteq \Lambda_S$ , i.e if  $\vdash_{\Lambda} \alpha$  implies  $\models^S \alpha$  for all  $S \in S$ .
2. A modal logic  $\Lambda$  is strongly complete with respect to  $S$  if for any set of formulas  $\Gamma \cup \{\alpha\}$ , if  $\Gamma \models^S \alpha$  then  $\Gamma \vdash_{\Lambda} \alpha$  for all  $S \in S$  and it is weakly complete with respect to  $S$  if for any formula  $\alpha$  if  $\models^S \alpha$  then  $\vdash_{\Lambda} \alpha$  for all  $S \in S$ .

+

## 1. Preliminary

---

To show completeness results you will create a *canonical* model and then show that the given modal logic is complete with respect to this model. It can be shown that **K** is sound and strongly complete with respect to the class of all frames. A lot of different completeness results can be found in (Lemmon 1977) and Blackburn 2002. Here we will just state a few of the results about **GL**.

**Theorem 1.1** **GL** is not sound and strongly complete with respect to any class of frames.

There is another result concerning the completeness and soundness of **GL**. To state this we first need some definitions concerning relations:

**Definition 1.12** A relation  $R$  on frame  $\mathcal{H} = \langle W, R \rangle$  is said to be *transitive* if for all  $w_1, w_2, w_3 \in W$ , whenever  $w_1 R w_2$  and  $w_2 R w_3$  then  $w_1 R w_3$ .

The relation  $R$  is said to be *well-founded* on  $\mathcal{H}$  if every non-empty subset  $V \subseteq W$  has a minimal element with respect to  $R$ . In other words  $R$  is well-founded if there is no infinite sequence  $\dots R w_2 R w_1 R w_0$

Further the relation  $R$  is said to be *conversely well-founded* on  $\mathcal{H}$  if the converse  $R^{-1}$  of  $R$  is well-founded; i.e if there is no infinite sequence such that  $w_0 R w_1 R w_2 R \dots$   $\neg$

We can now state the following theorem:

**Theorem 1.2** **GL** is sound and weakly complete with respect to the class of transitive and conversely well-founded frames.

In chapter 4 we will show some further results about **GL** and in chapter 5 we will show a fixed point theorem about **GL**. We will end this section with the following proposition that will be needed later:

**Proposition 1.2**  $\vdash_{\mathbf{GL}} \Box(\alpha \leftrightarrow \beta) \rightarrow (\Box\alpha \leftrightarrow \Box\beta)$

## 1.2. Gödels Incompleteness Theorems

In this section we will state and sketch the proof of Gödels incompleteness theorems. In this sketch we will define a predicate  $\text{Pr}(\cdot)$  that says a given formula has a proof in arithmetics.

### 1.2.1. Primitive Recursive Functions

We will start of by defining a class of function that will both play a crucial role in the proof of Gödels Incompleteness Theorems, and in the rest of this project.



**Definition 1.13** The class of primitive recursive functions is the smallest class closed under the following schemata:

- I.  $S(x) = x + 1$  is primitive recursive.
- II.  $Z(x) = 0$  is primitive recursive.
- III.  $P_i^n(x_1, \dots, x_n) = x_i$  is primitive recursive.
- IV. If  $g, h_1, \dots, h_m$  are primitive recursive then so is

$$f(x_1, \dots, x_n) = g(h_1(x_1, \dots, x_n), \dots, h_m(x_1, \dots, x_n))$$

- V. If  $g$  and  $h$  are primitive recursive and  $n \geq 1$  then  $f$  is also primitive recursive where:

$$\begin{aligned} f(0, x_1, \dots, x_n) &= g(x_1, \dots, x_n) \\ f(x_1 + 1, x_2, \dots, x_n) &= h(x_1, f(x_1, \dots, x_n), x_2, \dots, x_n) \end{aligned}$$

—

We can also define relations as being primitive recursive:

**Definition 1.14** A relation  $R \subseteq \omega^n$  is primitive recursive if its characteristic function :

$$\chi_R(\vec{x}) = \begin{cases} 0 & \text{if } R(\vec{x}) \\ 1 & \text{if } R(\vec{x}) \end{cases}$$

—

So with these definitions we can show that some well known functions are primitive recursive. The proofs of these can be rather tiresome and have been left out.

## 1. Preliminary

---

**Proposition 1.3** The following list of functions are all primitive recursive:

- |    |   |                     |
|----|---|---------------------|
| 1. | $K_k^n(x_1, \dots, x_n) = k$  | Constant            |
| 2. | $A(x, y) = x + y$   | Addition            |
| 3. | $M(x, y) = x \cdot \dots \cdot y$   | Multiplication      |
| 4. | $E(x, y) = x^y$   | Exponentiation      |
| 5. | $pd(x) = \begin{cases} x - 1, & x > 0 \\ 0, & x = 0 \end{cases}$          | Predecessor         |
| 6. | $x \dot{-} y = \begin{cases} x - y & x \geq y \\ 0, & x < y \end{cases}$  | Cut-off subtraction |
| 7. | $sg(x) = \begin{cases} 0, & x = 0 \\ 1, & x > 0 \end{cases}$              | Signum              |
| 8. | $\overline{sg}(x) = \begin{cases} 1, & x = 0 \\ 0, & x > 0 \end{cases}$   | Signum complement   |
| 9. | $ x - y  = \begin{cases} x - y, & x \geq y \\ y - x, & x < y \end{cases}$ | absolute value      |

### 1.2.2. PRA

In this section we will specify the rules and language of **PRA**. First of we will define the language  $\mathcal{L}_{\mathbf{PRA}}$  of **PRA**.

**Definition 1.15** The language  $\mathcal{L}_{\mathbf{PRA}}$  consists of the following symbols:

- Variables :  $v_0, v_1, \dots$
- Constant :  $\bar{0}$
- Function symbols :  $\bar{f}$  for each primitive recursive function  $f$
- Relation symbols :  $=$
- propositional connectives :  $\neg, \wedge, \vee, \rightarrow$
- Quantifiers :  $\forall, \exists$

⊢

We will now define the notions of terms and formulae of **PRA**. This part is mostly done to settle notations.

**Definition 1.16** 1. The set of *terms* of the language of **PRA** is defined inductively by:

- a)  $\bar{0}$  is a term and each  $v_i$  is a term.
- b) If  $f$  is an  $n$ -ary function symbol and  $t_1, \dots, t_n$  are terms then  $\bar{f}t_1 \dots t_n$  is a term.
- 2. The set of *formulae* of the language of **PRA** is defined inductively by:
  - a) If  $t_1$  and  $t_2$  are terms then  $= t_1 t_2$  is a formula.
  - b) If  $F$  and  $G$  are formulae, so are  $\neg F$ ,  $\wedge FG$ ,  $\vee FG$  and  $\rightarrow FG$ .
  - c) If  $F$  is a formula and  $v$  is a variable, then  $\exists v F$  and  $\forall v F$  are also formulae.

→

We use Polish notation so we do not have parentheses in our language. In practice we will use parentheses and infix notation. We will also need the definition of a *sentence*.

**Definition 1.17** A formula  $F$  of **PRA** is called a sentence if it has no free variables; i.e if all variables  $v$  that occurs in  $F$  are bound. →

Having defined the language of **PRA** we can now state the different axioms of **PRA**:

**Definition 1.18** The axioms **PRA** are the following:

- 1. Propositional axioms
  - a)  $F \rightarrow (G \rightarrow F)$
  - b)  $(F \rightarrow (G \rightarrow H)) \rightarrow ((F \rightarrow G) \rightarrow (F \rightarrow H))$
  - c)  $F \wedge G \rightarrow F$
  - d)  $F \wedge G \rightarrow G$
  - e)  $F \rightarrow (G \rightarrow F \wedge G)$
  - f)  $F \rightarrow F \vee G$
  - g)  $G \rightarrow F \vee G$
  - h)  $(F \rightarrow H) \rightarrow ((G \rightarrow H) \rightarrow (F \vee G \rightarrow H))$
  - i)  $(F \rightarrow G) \rightarrow ((F \rightarrow \neg G) \rightarrow \neg F)$
  - j)  $\neg \neg F \rightarrow F$
- 2. Quantifier axioms
  - a)  $\forall v Fv \rightarrow Ft$
  - b)  $Ft \rightarrow \exists v Fv$

## 1. Preliminary

---

Where  $t$  is substitutable for  $v$  in  $Fv$  in both cases.

### 3. Equality axioms

- a)  $v_0 = v_0$
- b)  $v_0 = v_1 \rightarrow v_1 = v_0$
- c)  $v_0 = v_1 \wedge v_1 = v_2 \rightarrow v_0 = v_2$
- d)  $v_i = w \rightarrow \bar{f}(v_1, \dots, v_i, \dots, v_n) = \bar{f}(v_1, \dots, w, \dots, v_n)$

Where  $1 \leq i \leq n$  and  $\bar{f}$  is an  $n$ -ary function symbol.

### 4. Non-logical axioms

#### a) Initial functions

- i.  $\bar{Z}(v_0) = \bar{0}$
- ii.  $\neg(\bar{0} = \bar{S}(v_0))$
- iii.  $\bar{S}(v_0) = \bar{S}(v_1) \rightarrow v_0 = v_1$
- iv.  $\bar{P}_i^n(v_1, \dots, v_n) = v_1$  for  $1 \leq i \leq n$

#### b) Derived functions

- i.  $\bar{f}(v_1, \dots, v_n) = \bar{g}(\bar{h}_1(v_1, \dots, v_n), \dots, \bar{h}_m(v_1, \dots, v_n))$ . Here  $f$  is defined by composition of  $g, h_1, \dots, h_m$
- ii. Let  $f$  be defined by primitive recursion from the primitive recursive function  $g$  and  $h$ , then the following two things hold:  
 $\bar{f}(\bar{0}, v_1, \dots, v_n) = \bar{g}(v_1, \dots, v_n)$  and  
 $\bar{f}(\bar{S}v_0, v_1, \dots, v_n) = \bar{h}(\bar{f}(v_0, v_1, \dots, v_n), v_0, v_1, \dots, v_n)$

#### c) Induction

$F(\bar{0}) \wedge \forall v(\delta v \rightarrow F(\bar{S}v)) \rightarrow \forall v F(v)$  where  $F(v)$  is

$$\exists v_n(\bar{f}(v, v_0, v_1, \dots, v_n) = \bar{0})$$

⊢

The induction axiom seems a bit strange at first. If we had allowed full induction; i.e induction on every formula, the system we would have defined would be known as **PA**; Peano Arithmetics. Having written down the axioms we will move on to the inference rules of **PRA**.

**Definition 1.19** The inference rules of **PRA** are the following:

1. From  $F, F \rightarrow G$ , derive  $G$ . (modus ponens)

2. From  $Fv \rightarrow G$  derive  $\exists F \rightarrow G$ , under the assumption that no  $v$  occurs free in  $G$ .
3. From  $G \rightarrow Fv$  derive  $G \rightarrow \forall v Fv$ , under the assumption that no  $v$  occurs free in  $G$ .

A formal derivation in **PRA** is a sequence of formulas of **PRA**  $F_0, F_1, \dots, F_k$  such that each  $F_i$  is either an axiom of **PRA** or follows from two other formulas  $F_j, F_l$  where  $j, l < i$  by one of the three inferences rules.  $\dashv$

We will end this section with a theorem that states that **PRA** can compute the primitive recursive functions.

**Theorem 1.3** Let  $f$  be an  $n$ -ary primitive recursive function and let  $\bar{f}$  be the function symbol representing it in **PRA**. Then for all  $n_1, \dots, n_m, n \in \omega$  we have:

$$f(n_1, \dots, n_m) = n \Rightarrow \mathbf{PRA} \vdash \bar{f}(\bar{n}_1, \dots, \bar{n}_m) = \bar{n}$$

That **PRA** can compute the primitive recursive functions makes it possible to encode the syntax of **PRA**. This will be explained in the next section.

### 1.2.3. Arithmetication of the syntax

In this section we will discuss how each expression in the language  $F$  of **PRA** can be given a unique code-number  $\ulcorner F \urcorner$  called the Gödel number. This makes it possible for **PRA** to express things about it self. For this encoding to work we will need the fundamental theorem of arithmetic that states that every natural number  $a \geq 2$  has a unique representation:

$$a = p_{i_0}^{n_0} \cdots p_{i_k}^{n_k}$$

Where each  $p$  is a distinct prime and all the  $n_i$  are positive. If given a sequence  $(j_0, \dots, j_t)$  we can code it with a unique code in the following way:

$$c = 2^{j_0+1} 3^{j_1+1} \cdots p^{j_t+1}$$

So by this it is possible to give each formula of **PRA** a unique code. We will start of by listing the codes for some of our symbols of **PRA**:

1.  $\ulcorner \bar{0} \urcorner$  is (0).
2.  $\ulcorner = \urcorner$  is (1).
3.  $\ulcorner \neg \urcorner$  is (2);  $\ulcorner \wedge \urcorner$  is (3);  $\ulcorner \vee \urcorner$  is (4) and  $\ulcorner \rightarrow \urcorner$  is (5).

## 1. Preliminary

---

4.  $\ulcorner \forall \urcorner$  is (6) and  $\ulcorner \exists \urcorner$  is (7).

5.  $\ulcorner v_i \urcorner$  is  $(8, i)$ .

These are the easy one to gives. The hard ones are the codes for the functions symbols. We will define the codes for these inductively as seen in the following table:

| Function  | Index                                      |
|---|--|
| $Z(x) = 0$  | $(9, 1, 1)$                                |
| $S(x) = x + 1$  | $(9, 2, 1)$                                |
| $P_i^n(x_1, \dots, x_n) = x_i$  | $(9, 3, n, i)$                             |
| $f(\vec{x}) = g(h_1(\vec{x}), \dots, h_m(\vec{x}))$                                     | $(9, 4, n, m, (g^*, h_1^*, \dots, h_m^*))$ |
| $f(0, x_1, \vec{x}) = g(\vec{x})$<br>$f(x + 1, \vec{x}) = h(f(x, \vec{x}), x, \vec{x})$ | $(9, 5, n + 1, g^*, h^*)$                  |

Table 1.1.: The Codes for the Function Symbols

This encoding is primitive recursive so **PRA**, can do this encoding. But the proof of that is very tedious, so it is left out. A lot of different relations and functions can be shown to be primitive recursive, one of the important ones is the function  $subst(x; y, z)$  which substitutes  $z$  for  $y$  in the formula  $x$ . With this it is possible to show primitive recursively define our axioms of **PRA**. This means that **PRA** is capable of identifying its own axioms. Further it can also identify that if a given sequence of formulas is a proof of another given formula; i.e the relation that says "y codes a derivation of the formula with code x". We shorten this as  $Prov(x, y)$  and we further define the non primitive recursive relation "x codes a provable formula" as:  $Pr(x) \leftrightarrow \exists y(Prov(y, x))$

### 1.2.4. The proof predicate and the theorems

The relation  $Pr(x)$  has the following properties called Löbs derivability conditions:

$$D1 \text{ PRA} \vdash F \Rightarrow \vdash Pr(\ulcorner F \urcorner)$$

$$D2 \text{ PRA} \vdash Pr(\ulcorner F \rightarrow G \urcorner) \rightarrow (Pr(\ulcorner F \urcorner) \rightarrow Pr(\ulcorner G \urcorner))$$

$$D3 \text{ PRA} \vdash Pr(\ulcorner F \urcorner) \rightarrow Pr(\ulcorner Pr(\ulcorner F \urcorner) \urcorner)$$

it is also clear that  $\neg Pr(\perp)$ , where  $\perp$  is some false statement expresses consistence. We will denote this by  $Con$ . We can further with the use of the  $subst$  function prove the following important lemma:

**Lemma 1.1** (The fixed point lemma) Given any formula  $G$  where the only free variable is  $v$ , we can find a sentence  $F$  such that:

$$\mathbf{PRA} \vdash F \leftrightarrow G(\ulcorner F \urcorner)$$

This lemma makes it possible for us to show the first incompleteness theorem:

**Theorem 1.4** (The First Incompleteness Theorem) The predicate  $\text{Pr}(\cdot)$  have the following properties for all formulas  $G$ :

1.  $\mathbf{PRA} \vdash G \Rightarrow \vdash \text{Pr}(\ulcorner G \urcorner)$
2.  $\mathbf{PRA} \vdash \text{Pr}(\ulcorner G \urcorner) \Rightarrow \vdash G$

If we let  $\mathbf{PRA} \vdash F \leftrightarrow \neg \text{Pr}(\ulcorner F \urcorner)$ , we then have:

1.  $\mathbf{PRA} \not\vdash F$
2.  $\mathbf{PRA} \not\vdash \neg F$

I.e we have a sentence where neither it or its negation can be proven. So our system is incomplete.

Since  $\text{Pr}(\cdot)$  fullfills the derivability conditions the second theorem can also be proven.

**Theorem 1.5** (The Second Incompleteness Theorem) Under the assumption that  $\mathbf{PRA}$  is consistent we have that  $\mathbf{PRA} \not\vdash \text{Con}$

The main goal for the rest of this project is now to prove that the  $\Box$  in **GL** "behaves" in the same way as the  $\text{Pr}(\cdot)$  predicate from arithmetics. How to should be understood will be explained latter on. To prove this theorem we will first need a main result from Recursion Theory called the recursion theorem.

We will need one more result about the proof predicate in the project. It is called the formalized Löb's theorem, and this theorem together with the derivability conditions tells the full story about the predicate  $\text{Pr}(\cdot)$

**Theorem 1.6** (Formalized Löb's Theorem) Let  $F$  be any sentence of  $\mathbf{PRA}$ . Then:

$$\mathbf{PRA} \vdash \text{Pr}(\ulcorner \text{Pr}(\ulcorner F \urcorner) \rightarrow F \urcorner) \rightarrow \text{Pr}(\ulcorner F \urcorner)$$

This theorem was an original an answer to a question asked by Leon Henkin: "If  $\mathbf{PRA}$  proves that  $G \leftrightarrow \text{Pr}(\ulcorner G \urcorner)$  what does this say about  $G$ ? Löb's answer is then that if  $\mathbf{PRA}$  can only that  $\text{Pr}(\ulcorner G \urcorner \rightarrow G$  in the case where  $\mathbf{PRA}$  proves  $G$  itself.

We will in a later chapter need the following lemma:

## 1. Preliminary

**Lemma 1.2** Let  $\bar{f}$  be an  $n$ -ary primitive recursive function symbol. There is a function  $g$  depending on  $f$ , such that:

$$\mathbf{PRA} \vdash \bar{f}v_0 \dots v_{n-1} = v \rightarrow \text{Prov}(\bar{g}v_0 \dots v_{n-1}, \ulcorner f\dot{v}_0 \dots \dot{v}_{n-1} = \dot{v}_n \urcorner)$$

Where  $\ulcorner f\dot{v}_0 \dots \dot{v}_{n-1} = \dot{v}_n \urcorner$  denotes the code of the formula where we have substituted the variables with the numerals:  $\bar{v}_0, \dots, \bar{v}_{n-1}$ .

The Formalized Löb's Theorem and the derivability conditions can be shown to hold for a wide range of arithmetical systems. A overview over this subject will be given in chapter 3. If we write  $\vdash F$  instead of  $\mathbf{PRA} \vdash F$  it means we have not specified the arithmetical system we are looking at.

If we compare the properties of  $\text{Pr}(\cdot)$  with the properties of the  $\Box$ -operator of  $\mathbf{GL}$ , we get the following table:

|  |   |
|--|---|
| $\vdash_{\mathbf{GL}} \alpha \Rightarrow \vdash_{\mathbf{GL}} \Box \alpha$                             | $\vdash F \Rightarrow \vdash \text{Pr}(\ulcorner F \urcorner)$  |
| $\vdash_{\mathbf{GL}} \Box(\alpha \rightarrow \beta) \rightarrow (\Box \alpha \rightarrow \Box \beta)$ | $\vdash \text{Pr}(\ulcorner F \rightarrow G \urcorner) \rightarrow (\text{Pr}(\ulcorner F \urcorner) \rightarrow \text{Pr}(\ulcorner G \urcorner))$ |
| $\vdash_{\mathbf{GL}} \Box \alpha \rightarrow \Box \Box \alpha$  | $\vdash \text{Pr}(\ulcorner F \urcorner) \rightarrow \text{Pr}(\ulcorner \text{Pr}(\ulcorner F \urcorner) \urcorner)$                               |
| $\vdash_{\mathbf{GL}} \Box(\Box \alpha \rightarrow \alpha) \rightarrow \Box \alpha$                    | $\vdash \text{Pr}(\ulcorner \text{Pr}(\ulcorner F \urcorner) \rightarrow F \urcorner) \rightarrow \text{Pr}(\ulcorner F \urcorner)$                 |

Table 1.2.: The properties of  $\Box$  and the predicate  $\text{Pr}(\cdot)$

It is clear from this table that the predicate  $\text{Pr}(\cdot)$  and the  $\Box$ -operator of  $\mathbf{GL}$  have a lot of the same properties in common. Solovay's completeness theorems shows that the modal logic  $\mathbf{GL}$  in some way axiomatize the proof predicate of a wide range of different arithmetical systems. A somewhat fun fact is, that the first time the principle  $\Box(\Box \alpha \rightarrow \alpha) \rightarrow \Box \alpha$  was used in modal logic, was in 1963 in a paper by Timothy John Smiley about the logic basis of ethic (Smiley 1963). Further the modal logic now known as  $\mathbf{GL}$  was considered before the arithmetical interpretations of it was conjectured. Segerberg actually proved that it was completeness with respect to the conversely well-founded frames before he knew about its importance in arithmetic.



## 2. Recursion Theory

{chap:Recur}

**Note:** grafer kan ses som formler fra PRA, så bare brug F, G og H for disse We will begin this section with define two classes of functions; the *Partial Recursive Functions* and the *Turing Computable functions*. These two classes of functions gives rise to the same class of functions The idea behind these two classes of functions is to define what we intuitively mean with a computable function. The proofs in this section will be rather informal. The goal of this section will be to state and prove Kleene's recursion theorem, which will play a crucial role in our proof of Solovay's two completeness theore

The topic I have chosen to call *recursion theory* is today often known as *computability theory*. Robert Soares have stated some arguments for why this name is better suited for the discipline than the name recursion theory in his book (Soare 2016).

This sections will follow (Soare 1987), unless stated otherwise.

### 2.1. Partial Recursive Functions and Recursive Functions

The class of partial recursive functions is an enlargement of the primitive recursive functions. The class of primitive recursive functions capture a lot of the function we intuitively sees as computable. It does not contain any function that we would say was incomputable.

But the primitive recursive functions do not include all intuitively computable functions. For example the following function, called the Ackermann function, that is clearly computable is not in the class of primitive recursive functions:

$$\begin{aligned} \text{Ack}(0, n) &= n + 1 \\ \text{Ack}(m + 1, 0) &= \text{Ack}(m, 1) \\ \text{Ack}(m + 1, n + 1) &= \text{Ack}(m, \text{Ack}(m + 1, n)) \end{aligned}$$

This function grows *to fast* to be a primitive recursive function, i.e we have for all primitive recursive functions  $f(x)$  that for some  $y$  that  $\text{Ack}(x, y) > f(x)$  for all  $x \in \omega$ . proof of this fact can be found in **Find ref**.

## 2. Recursion Theory

---

So we will need to expand our class of functions, if we want to *capture* all intuitively computable functions. We will expand them in the following way:

**Definition 2.1** The class of *partial recursive* (from now on some times called (p.r) functions is the least class closed under schemata I through V from the definition of primitive recursive functions and the following VI schema .

VI. If  $\theta(x_1, \dots, x_n, y)$  is a partial recursive function of  $n + 1$  variables and

$$\psi(x_1, \dots, x_n) = \mu y [\theta(x_1, \dots, x_n, y) \downarrow = 0 \\ \wedge \forall z \leq y [\theta(x_1, \dots, x_n, z) \downarrow]]$$

Then  $\psi$  is a partial recursive function of  $n$  variables

A partial recursive function that is total is called a total recursive function; abbreviated to recursive function.  $\dashv$

This is one way in which one can define what the computable functions are. In the next section we will come with another definition of a type of functions, and it will be shown that these leads to the same class of functions.

We will end this section with the following definition:

**Definition 2.2** A relation  $R \subset \omega^n$  where  $n \geq 1$ , is recursive (primitive recursive) if its characteristic function  $\chi_R$  is recursive (primitive recursive). The case where  $n = 1$  is the case where  $R$  is a set  $A \subset \omega$  so we also have the definition of a set being recursive.  $\dashv$

## 2.2. Turing Computable Functions

Another way to describe the intuitively computable functions is via a Turing machine.

**Definition 2.3** A *Turing machine*  $M$  consists of a two-way infinite tape that is divided into different cells and a finite set of internal states  $Q = \{q_0, \dots, q_n\}$ ,  $n \geq 1$ . Each cell is either blank: B or has value 1. The following three things can happen in a single step:

1. Change form one state to another.
2. Change the scanned symbol  $s$  to another symbol  $s' \in S = \{1, B\}$
3. Move the reading head one cell to the right R or the left L.

The operation of  $M$  is controlled by a partial map:

$$\delta : Q \times S \rightarrow Q \times S \times \{R, L\}$$

Which may not be defined for all arguments. ⊥

The situation can be seen in the following figure:




Figure 2.1.: A Turing Machine

The way to understand this definition is the following: if  $(q, s, q', s', X) \in \delta$  it means that the machine  $M$  is in stage  $q$  where it scans symbol  $s$  then changes to state  $q'$  and replaces  $s$  by  $s'$ . Lastly it moves to the right if  $X = R$  and the left if  $X = L$ . The map  $\delta$  is called a Turing program if it can be views as a finite set of quintuples. If the input integer is  $x$  then it will be represented by a string of  $x + 1$  consecutive 1's, where all other cells are blank.

Further the machine  $M$  start in the state  $q_1$  scanning the left-most cell that contains a 1. The machine stops if it reaches the halting state  $q_0$ , and it will then output the number  $y$  which is the total number of 1's on the tape in the this state. If  $M$  with input  $x$  halts and outputs  $y$  we say that  $M$  computes the partial function  $\psi(x) = y$

The conditi [Indfør noget mere tekst her]on a is determined by:

1. The current state  $q_i$
2. The symbol  $s_0$  that is being scanned.
3. The symbols on the tape to the right of  $s_0$  up to the last 1. Denote this sequence by  $s_1, s_2, \dots s_n$

## 2. Recursion Theory

---

4. The symbols on the tape to the left of  $s_0$  up to the first 1. Denote this sequence by  $s_{-1}, s_{-2}, \dots s_{-m}$

This is called the configuration of the machine and we can write it as follows:

$$s_{-m} \cdots s_{-1} q_0 s_0 s_1 \cdots c_n$$

**Definition 2.4** A Turing computation according to the Turing program  $P$  with input  $x$  is a sequence of configurations  $c_0, c_1, \dots, c_n$  such that  $c_0$  represent the machine in the halting state  $q_0$ , and the transition  $c_i \rightarrow c_{i+1}$  for all  $i < n$  is giving by the Turing program  $P$ .  $\dashv$

**Proposition 2.1** Each Turing program  $P_e$  can be assigned a Gödel number  $e$ .

*Proof.* We will use the fact that each  $x \in \omega$  has a unique prime decomposition:

$$x = p_0^{x_0} \cdots p_n^{x_n} \cdots$$

We can assign a number to each quintuple  $(q_{i,j}, q_k, s_l, r_m)$  in a Turing program  $P$  in the following way[Omformuler denne sætning]:

$$p_0^{1+i} p_1^{1+j} p_2^{1+k} p_3^{1+l} p_4^{1+m}$$

Where we have that  $r_0 = R$  and  $r_1 = L$ . Since the prime decomposition is unique, each different state of the program has a unique code. Each Turing is a sequence of different states and we can thus for an arbitrary Turing program  $P_e$  we let  $e_0, \dots e_n$  denote the Gödel number of each different state and set:

$$e = p_0^{e_0} \cdots p_n^{e_n}$$

Thus each Turing program  $P_e$  has a unique Gödel number  $e$ .  $\dashv$

Since each Turing program has a unique code we can list them and be able to find any program  $P_e$  by its code  $e$ . This gives the following definition:

**Definition 2.5** The  $P_e$  be the Turing program with Gödel number  $e$  in the list and let  $a_e^{(n)}$  be the function of  $n$  variables computed by  $P_e$ . Further let  $a_e$  abbreviate  $a_e^{(1)}$ .  $\dashv$

### 2.3. The $s$ - $m$ - $n$ -Theorem

It can be proven that the two classes of functions; partial recursive and Turing computable functions gives rise to the same class of partial functions. This can

be seen as evidence for *Church's Thesis* which states that this class of functions coincide with the function that we see as intuitively computable. In the rest of this project we will assume that the Church's Thesis is true.

We will begin by proving the padding lemma, which states that each partial function  $\varphi_x$  has an infinite amount of indices.

**Lemma 2.1 (The Padding Lemma)** Each partial recursive function  $\varphi_i$  has  $\aleph_0$  indices, and for each  $x$  we can *effectively* find an infinite set  $A_x$  of indices for the same partial function.

*Proof.* For any program  $P_x$  that have internal states:  $\{q_0, \dots, q_n\}$  we can add extra instructions  $q_{n+1}B\ R, q_{n+2}B\ R, \dots$  such that we get a new program for the same computation.  $\dashv$

The following theorem will show that each Turing computable function is in fact partial recursive. The converse also holds and the proof of this fact can be found in (Kleene 1952)

**Theorem 2.1 (The Normal Form Theorem)** There exist a predicate  $T(e, x, y)$  and a function  $U(y)$  that are primitive recursive such that:

{thm:Normalform}

$$\varphi_e(x) = U(\mu y T(e, x, y))$$

*Proof.* We showing that the predicate  $T(e, x, y)$  exists and is primitive recursive. This predicate informally states that  $y$  is the code of Turing program  $P_e$  with input  $x$ . For each possible configuration  $c$ , we can assign a code:

$$\#(c) = 2^{1+i} 3^{1+\#(s_0)} 5^r 7^l$$

Where  $\#(s) = 0$  if  $s = B$  and is equal to 1 otherwise,  $r = \prod_{j \geq 1} p_j^{\#(s_j)}$  and  $l = \prod_{j \leq -1} p_j^{\#(s_j)}$ . We can now define the code of a Turing computation  $c_0, c_1, \dots, c_n$  according to  $P_e$  to be:

$$y = 2^e \prod_{i \leq n} p_{i+1}^{\#(s_i)}$$

We can now define  $T(e, x, y)$  to be [Læs i Kleene] Having defined the predicate  $T$  we can check if it holds. By proposition 2.1 we can "recover" the program  $P_e$  from  $e$ . Then we can recover the computation  $c_0, c_1, \dots, c_n$  from  $y$  if  $y$  codes such a thing. We can now check if  $c_0, c_1, \dots, c_n$  is a computation according to  $P_e$  with  $x$  as the input in the first configuration  $c_0$ . If this is true, then  $U(y)$  just outputs the number of 1's in the final configuration  $c_n$ . [Læs mere i Klenne]  $\dashv$

This theorem also gives us that each partial recursive function can be created by two primitive recursive functions, with a single application of the  $\mu$ -operator.

## 2. Recursion Theory

{thm:Emu}

**Theorem 2.2 (Enumeration Theorem)** There is a partial recursive function of 2 variables  $\varphi_z^{(2)}(e, x)$  such that  $\varphi_z^{(2)}(e, x) = \varphi_e(x)$ .

*Proof.* By Theorem 2.1 we will define  $\varphi_z^{(2)}(e, x) = U(\mu y T(e, x, y)) = \varphi_e(x)$   $\dashv$

We will need the following notation in the next proof:

**Definition 2.6** Set  $\langle x, y \rangle$  to be the image of  $(x, y)$  under the injective recursive pairing function:

$$\frac{1}{2}(x^2 + 2xy + y^2 + 3x + y)$$

This function is from  $\omega \times \omega$  onto  $\omega$ .  $\dashv$

**Theorem 2.3 (s-m-n theorem)** For every  $m, n \geq 1$  there exists an injective recursive function  $s_n^m$  of  $m + 1$  variable such that for all  $x, y_1, \dots, y_m$

$$\varphi_{s_n^m(x, y_1, \dots, y_m)}^{(n)} = \lambda z_1, \dots, z_n [\varphi_x^{(m+n)}(y_1, \dots, y_m, z_1, \dots, z_n)]$$

*Proof.* I will follow Soare and only proof the case where  $m = n = 1$ . I.e the case where we have to proof:

$$a_{s_1^1(x, y)}(z) = [a_x^2(y, z)]$$

Let  $x$  and  $y$  på given. Then  $s_1^1(x, y)$  can be described as follows:

1. Let  $P_x$  the Turing program with code  $x$ .
2. Change  $P_x$  into another Turing program  $P_{x'}$  such that:  $P_{x'}$  writes  $y + 1$  "1" left of the input, such that there is a B between these 1 and the other input. Further i places the head to the left of the new input and proceeds to run  $P_x$ .
3. outputs  $x'$

it is clear that  $P_{x'}$  on input  $z$  compute the same as  $p_e$  would on input  $(x, y)$ ; i.e  $\varphi_{x'} = \varphi_x^{(2)}(y, z)$ . Further we have that  $x' = s_1^1(x, y)$  By Church's Thesis the function  $s = s_1^1$  is recursive, since it can be computed effectively. If it is not injective it can be replaced by a injective recursive function  $s'$  such that  $\varphi_{s(x, y)} = \varphi_{s'(x, y)}$  by using the padding lemma and by defining  $s'(x, y)$  in increasing order of  $\langle x, y \rangle$ .  $\dashv$

A full proof of this statement can be found in Kleene. The  $s - m - n$  theorem plays a crucial role in both the proof and the use of the recursion theorem. **Skriv noget om intuition.** Therefore we will use the  $s - m - n$  theorem to prove the following proposition:

**Proposition 2.2** There is a recursive function  $g$  of two variables such that for all  $x, y$ :

$$\varphi_{g(x,y)} = \varphi_x \varphi_y$$

*Proof.* By Church's Thesis it is clear that  $\eta = \varphi_x \varphi_y$  is a partial recursive function since it is the product of two partial recursive functions.

To end the proof we must show that we can find Gödel number for  $\eta$  in an uniform effective way from  $x$  and  $y$  as  $x$  and  $y$  vary. We will start of by defining:

$$\theta(x, y, z) = \varphi_x(\varphi_y(z)) = \varphi_{x_1}(x, \varphi_{x_1}(y, z))$$

Where  $\varphi_{x_1}$  is the function  $\varphi_z^{(2)}$  from Theorem 2.2. By the church thesis this function is partial recursive and has an index  $e$ . So by applying the  $s - m - n$  Theorem we get:

$$\varphi_x \varphi_y = \lambda z [\varphi_e(x, y, z)] = \varphi_{s_1^2(e, x, y)}$$

And thus  $s_1^2(e, x, y)$  is our  $g$  (i.e  $\lambda xy[s_1^2(e, x, y)]$ ). ⊢

## 2.4. Recursively Enumerable sets and the Graph of a function

In this section we will introduce the two concepts *recursive enumerable* sets and the *graph* of a function. We will further show that there is connection between these two concepts.

**Definition 2.7** A set  $A$  is *recursively enumerable* (r.e.) if  $A$  is the domain of some primitive recursive function. Further we define the following two sets:

1. We let the  $e$ th r.e set be denoted by:

$$E_e = \text{dom}(\varphi_e) = \{x : f(x) \downarrow\} = \{x : \exists y T(e, x, y)\}$$

- 2.

$$E_{e,s} = \text{dom}(\varphi_{e,s})$$

⊢

A set can be recursively enumerable without being recursive, which the following two propositions shows:

**Proposition 2.3** Let  $K = \{x : \varphi_x(x) \text{ converges}\} = \{x : x \in E_x\}$ , then  $K$  is r.e.

## 2. Recursion Theory

---

*Proof.* We have that  $K$  is the domain of the following primitive recursive function:

$$\psi(x) = \begin{cases} x & \text{if } \varphi_x(x) \text{ converges} \\ \text{undefined} & \text{otherwise.} \end{cases}$$

This function is primitive recursive by Church Thesis, since  $\psi(x)$  can be computed by program  $P_x$  on input  $x$ , which outputs  $x$  only if  $\varphi_x(x)$  converges.  $\dashv$

**Proposition 2.4**  $K$  is not recursive

*Proof.* If  $K$  were recursive we would that it had a recursive characteristic function  $K_\chi$ , and thus the following would be recursive:

$$f(x) = \begin{cases} \varphi_x(x) + 1 & \text{if } x \in K \\ 0 & \text{if } x \notin K \end{cases}$$

But we this  $f$  can not be recursive since for all  $x$  we have  $f \neq \varphi_x$ .  $\dashv$

**Forklar kort meningen med dette.**

Next goal is to show that the definition of a r.e sets is equivalent to the definition that there is an algorithm that enumerates the members of it. We will further also show that the definition of a r.e set is equivalent to the set being  $\Sigma_1$ , which is a concepts that will play a bigger role later on in this project.

**Definition 2.8** A set  $A$  is the *projection* of some relation  $R \subseteq \omega \times \omega$  if  $A = \{x : \exists y : R(x, y)\}$ . We further say that a set  $A$  is in  $\Sigma_1$  form, if  $A$  is the projection of some recursive relation  $R \subseteq \omega \times \omega$ .  $\dashv$

We can now show the following theorem:

**Theorem 2.4** A set  $A$  is r.e iff  $A$  is  $\Sigma_1$ .

*Proof.*  $(\Rightarrow)$  Since  $A$  is r.e we have that  $A = W_e = \text{dom} f_e$  for some  $e$ . This means that:

$$x \in W_e \Leftrightarrow \exists s(x \in W_{e,s}) \Leftrightarrow \exists s(T(e, x, s))$$

Since the relation  $T$  is primitive recursive and we have that the set  $A$  is the projection a recursive relation.

$(\Leftarrow)$  Let  $A = \{x : \exists y(R(x, y))\}$  where  $R$  is recursive. We then have that  $A = \text{dom} f$  where  $f(x) = \mu y(R(x, y))$  and thus  $A$  is r.e  $\dashv$

**Skriv noget metatekst**



1:RecSigma}

**Theorem 2.5** If there is a recursive relation  $R \subseteq \omega^{n+1}$  and if we have the following set:

$$A = \{x \mid \exists y_1 \dots \exists y_n R(x, y_1, \dots, y_n)\}$$

Then the set  $A$  is  $\Sigma_1$

*Proof.* We will start of by defining the relation  $S \subseteq \omega^2$  as follows:

$$S(x, z) \Leftrightarrow R(x, (z)_1, \dots, (z)_n)$$

Where we have the following prime decomposition of  $z$ :

$$z = p_1^{(z)_1} \dots p_k^{(z)_k}$$

Then the following equivalences holds:

$$\begin{aligned} \exists z S(x, z) &\Leftrightarrow \exists z R(x, (z)_1, \dots, (z)_n) \\ &\Leftrightarrow \exists y_1 \dots \exists y_n R(x, y_1, \dots, y_n) \end{aligned}$$

And thus the set  $A$  is clearly  $\Sigma_1$ . ⊢

From this theorem we can easily get the following corollary:

**Corollary 2.1** The projection of an r.e relation is r.e

The next definition will also play a role in our proof of Solovay's completeness theorems.

**Definition 2.9** The graph of a (partial) function  $\varphi$  is the relation:

$$(x, y) \in \text{graph}\varphi \Leftrightarrow f(x) = y$$

⊢

The relation of a graph can also be seen as a function in the language of **PRA**; so we will often denote the graph of a function  $\varphi(x) = y$  with the following notation:  $Fxy$ , and say that  $(x, y) \in \text{graph}(\varphi)$  if and only if  $Fxy$  is true.

**Theorem 2.6** If  $R \subseteq \omega^2$  is an r.e relation, then there is a p.r function sel called the selector function for  $R$  such that:

$$\text{sel}(x) \text{ is defined} \Leftrightarrow \exists y (R(x, y))$$

and if this is the case we have that  $(x, f(x)) \in R$

## 2. Recursion Theory

---

*Proof.* Since  $R$  is r.e it is  $\Sigma_1$ . This means that there is a recursive relation  $S$  such that  $R(x, y)$  holds iff  $\exists z(S(x, y, z))$ . Thus we can define the following primitive recursive function:

$$g(x) = \mu u(S(x, (u)_1, (u)_2))$$

And now we put  $f(x) = (g(x))_1$  ←

It will be the following theorem we will use in our proof later on.

**Theorem 2.7** A partial function  $\varphi$  is partial recursive iff its graph is recursive enumerable.

*Proof.*  $(\Rightarrow)$  The graph of  $\varphi_e$  is r.e by theorem 2.5 and the definition of a graph.

$(\Leftarrow)$  Since the graph of  $\varphi$  is assumed to be r.e we can conclude that  $\varphi$  is its own primitive recursive selector function. This is that  $R = \text{graph}\varphi$  can only have  $\varphi$  as its selector function. [Whyyy?] ←

## 2.5. The Recursion Theorem

In this section we will state and prove the recursion theorem. It will be crucial in the next [chapter?], since we will need it to define a function.

**Theorem 2.8 (The Recursion Theorem)** For every recursive function  $f$  there exists a fixed point  $n$  such that  $\varphi_n = \varphi_{f(x)}$

*Proof.* We will start of be defining the following *diagonal* function  $d(u)$  as:

$$\{\text{eq:du}\} \quad \varphi_{d(u)}(z) = \begin{cases} \varphi_{\varphi_u(u)}(z) & \text{if } \varphi_u(u) \text{ converges} \\ \text{undefined} & \text{else} \end{cases} \quad (2.1)$$

By the  $s-m-n$  theorem we have that the function  $d$  is injective and total. Further it is clearly seen that  $d$  is independent of  $f$ .

Given an arbitrary  $f$  we will choose an index  $v$  such that:

$$\{\text{eq:fd}\} \quad \varphi_v = f \circ d \quad (2.2)$$

Now set  $n = d(v)$ . We will show that this is a fixed point for the function  $f$ . Since  $f$  is total we also have that  $f \circ d$  is total. This means that  $\varphi_v(v)$  converges and that  $\varphi_{d(v)} = \varphi_{\varphi_v(v)}$ . Thus we have:

$$\varphi_n = \varphi_{d(v)} = \varphi_{\varphi_v(v)} = \varphi_{f \circ d(v)} = \varphi_{f(n)}$$

The second equality sign follows from 2.1 and the third follows from 2.2. ←

Following [Owens, find ref], the argument in the proof can be seen as a digitalization argument that fails. Commonly when we apply a digitalization argument, we have a class of sequences, with terms from an set  $A$ , that we arranges as the rows in a square matrix. We then have a map  $h : A \rightarrow A$  that induces a operation  $h^*$  on the set of sequences such that if  $\langle s(i), i \in I \rangle$  is a sequences in our matrix then

$$h^*(\langle s(i), i \in I \rangle) = \langle h(s(i)), i \in I \rangle$$

After having defined this map we will use it on the sequences that consists of the elements of the diagonal of the matrix and show that the resulting sequences is not one of the original sequences.

The digitalization argument "fails" in our case, since the sequences of the diagonal is already already one of the rows and thus the image  $h^*$  of this sequences will also be one of the rows; i.e the  $h$  has a fixed point.

The start of the proof can be seen as the following lemma:

**Lemma 2.2** There is a diaognal function  $d(u)$  such that:

$$\varphi_{d(u)}(z) = \begin{cases} \varphi_{\varphi_u(u)}(z) & \text{if } \varphi_u(u) \text{ converges} \\ \text{undefined} & \text{else} \end{cases} \quad (2.3)$$

Most of the times where one uses the Recursion Theorem, one actually uses this lemma to construct the given function.

### 2.5.1. Application of the Recursion Theorem

The recursion theorem is a "powerful" tool. It enables us to define a partial recursive function, which uses its own index as part of its definition. This The recursion theorem overrides this "self-reference" because we are using the  $s-m-n$  theorem to define a function  $f(x)$  and  $\varphi_{f(x)}(z) = (\dots, x \dots)$  and then taking a fixed point:  $\varphi_n = \varphi_{f(n)}$ . When we are making constructions like this the only thing we cannot do is use specific properties of the function  $\varphi_n$ . We will use the theorem in this way in our proof of Solovay's Completeness Theorems to define a function with help of the functions own Gödel number; and the recursion theorem makes this a viable tactic.

The following examples will show a few uses of this theorem.

**Example 2.1** We will show that there is a  $n$  such that:

$$W_n = \{n\}$$

We start of by using the  $s-m-n$  theorem to define  $W_{f(x)} = \{x\}$  then by the

## 2. Recursion Theory

---

recursion theorem we can choose  $n$  such that we have:

$$W_n = W_{f(n)} = \{n\}$$

—

The next example will be a application of the Recursion Theorem that is in the same vein as the one we will use when we have to prove Solovay's completeness theorems.

**Example 2.2** Let  $\psi : \omega^2 \rightarrow \omega$  and  $\theta : \omega^3 \rightarrow \omega$  be recursive functions and define the function  $\varphi : \omega^2 \rightarrow \omega$  by:

$$\begin{aligned}\varphi(0, y) &= \psi(y) \\ \varphi(x+1, y) &= \theta(\varphi(x, y), x, y)\end{aligned}$$

We will now show that  $\varphi$  is recursive by using the recursion theorem.

Let  $\phi_0 v_0 v_1$  be an arbitrary graph and let  $\phi_1 v_0 v_1$  be the graph of  $\psi$  and  $\chi v_0 v_1 v_2$  be the graph of  $\theta$ . We will now look at the following formula:

$$\Phi(\phi_0) : (v_0 = \bar{0} \wedge \phi_1 v_0 v_1) \vee (v_0 > \bar{0} \wedge \exists v(\phi_0(v_+ - 1, v) \wedge \chi v v_0 v_1))$$

We can see  $\ulcorner \Phi(\phi_0) \urcorner$  as a primitive recursive function of  $\ulcorner \phi_0 \urcorner$ . I.e we have:

$$\ulcorner \Phi(\phi_0) \urcorner = \eta(\ulcorner \phi_0 \urcorner)$$

We can now use the Recursion Theorem to chose a  $n$  such that we have:  $\varphi_{\eta(n)}^{(2)} = \varphi_n^{(2)}$  and for  $\varphi = \varphi_n^{(2)}$  we have:

$$\varphi(x, y) = z \leftrightarrow (x = 0 \wedge \psi(y) = z \vee (x > 0 \wedge \theta(\varphi(x-1, y), x, y) = z))$$

I.e  $\varphi$  does exactly what we want it to do. WE just need to define  $\varphi$  and this can be done by a  $\Sigma_1$  induction. —

This ends our discussion about recursion theory. In the next chapter we will generalize the notion of  $\Sigma_1$  sets and in chapter 6 we will use the recursion theorem to define a important function.

## 3. The Arithmetical Hierarchy and Fragments of Arithmetics

{chap:PRA}

In this section we will generalize the class of  $\sigma_1$  sets to the classes of  $\sigma_n$ ,  $\pi_n$  and  $\delta_n$  sets. We will use these sets in later chapters, since they make it possible for us to classify different arithmetical formulas. These definitions will make it possible for us to look at different fragments of arithmetic, since we can distinguish these fragments by which class of sets they allow induction on. We will start off by defining a "base" fragment called **EA** and then augment it with stronger induction axioms. This means that some of these fragments contains the other fragments. With this knowledge it will be possible for us to place **PRA** in this hierarchy.

We will further look at how much induction we have in **PRA**; i.e which class of formulae from the arithmetical hierarchy we can do induction on in **PRA**.

We will also introduce a two other ways to augment the different fragments without the adding more induction. These will play a role in the generalization of Solovay's completeness theorems to other fragments than **PRA**.

Lastly we will prove a generalization of Löb's third derivability condition for **PRA**.

This chapter should be seen as overview of the subject, and a lot of statements will be left unproven; since these will not play a major role in what is to come.

### 3.1. The arithmetical hierarchy

In this section we will introduce the so called arithmetical hierarchy. Parts of it has already be defined; i.e the sets  $\Sigma_1$ . In the arithmetical hierarchy, we can classify sets with respect to their quantifier complexity in their syntactical definition.

In the next section about the fragments of arithmetics, we will use the arithmetical hierarchy as a tool to determine the amount of induction we will have in a given fragment.

The goal of this section is to introduce the arithmetical hierarchy and prove some results about it. The most of these results might not be used going forward.

**Definition 3.1** We define the sets  $\Sigma_n$  and  $\Pi_n$  in the following way:

1. A set  $A$  is in  $\Sigma_0$  ( $\Pi_0$ ) if and only if  $A$  is recursive.

### 3. The Arithmetical Hierarchy and Fragments of Arithmetics

---

2. For  $n \geq 1$  the set  $A$  is in  $\Sigma_n$  if there is a recursive relation  $R(x, y_1, \dots, y_n)$  such that:

$$x \in A \text{ iff } \exists y_1 \forall y_2 \exists y_3 \cdots Q y_n R(x, y_1, \dots, y_n)$$

Here  $Q$  is  $\exists$  if  $n$  is odd and  $Q$  is  $\forall$  if  $n$  is even. We define  $A$  being in  $\Pi_n$  likewise.  $A$  is in  $\Pi_n$  if:

$$x \in A \text{ iff } \forall y_1 \exists y_2 \forall y_3 \cdots Q y_n R(x, y_1, \dots, y_n)$$

3.  $A$  is in  $\Delta_n$  if  $A \in \Sigma_n \cap \Pi_n$

We further say that a formula  $F$  is  $\Sigma_n$  ( $\Pi_n$ ) if it is  $\Sigma_n$  ( $\Pi_n$ ) as a relation of the variables that are free in it.  $\dashv$

In the rest of this project we will mostly look at formulas that are either  $\Sigma_n$  or  $\Pi_n$  and not sets, that have this property. Furthermore we have that  $\Sigma_0 = \Pi_0 = \Delta_0$ , and a formula  $F$  that is  $\Delta_0$  is also called a bounded formula.

We can show a few properties of these sets.

**Proposition 3.1** 1.  $A \in \Sigma_n \Leftrightarrow \bar{A} \in \Pi_n$

2.  $A \in \Sigma_n(\Pi_n) \Rightarrow (\forall m > n)(A \in \Sigma_m \cap \Pi_m)$
3.  $A, B \in \Sigma_n(\Pi_n) \Rightarrow A \cup B, A \cap B \in \Sigma_n(\Pi_n)$
4.  $(R \in \Sigma_n \wedge n > 0 \wedge A = \{x : \exists y R(x, y)\}) \Rightarrow A \in \Sigma_n$
5.  $(B \leq_m A \wedge A \in \Sigma_n) \Rightarrow B \in \Sigma_n$
6. If  $R \in \Sigma_n(\Pi_n)$  and  $A$  and  $B$  are defined by:

$$\langle x, y \rangle \in A \Leftrightarrow \forall z < y R(x, y, z)$$

and

$$\langle x, y \rangle \in B \Leftrightarrow \exists z < y R(x, y, z)$$

Then we have  $A, B \in \Sigma_n(\Pi_n)$

*Proof.* 1. If we have that:

$$A = \{x : \exists y_1 \forall y_2 \cdots R(x, y_1, \dots)\}$$

Then we have:

$$\bar{A} = \{x : \forall y_1 \exists y_2 \cdots \neg R(x, y_1, \dots)\}$$

Which is clearly  $\Pi_n$ .

2. If for example  $A = \{x : \exists y_1 \forall y_2 R(x, y_1, y_2)\}$ , then we can make the following reformulation of  $A$ :

$$A = \{x : \exists y_1 \forall y_2 \exists y_3 (R(x, y_1, y_2) \wedge y_3 = y_2)\}$$

This kind of reformulation can be done for any set in  $\Sigma_n$  ( $\Pi_n$ )

3. Let the following to sets be defined:

$$A = \{x : \exists y_1 \forall y_2 \cdots R(x, y_1, y_2, \dots)\}$$

$$B = \{x : \exists z_1 \forall z_2 \cdots S(x, z_1, z_2, \dots)\}$$

Then we have:

$$\begin{aligned} x \in A \cup B &\Leftrightarrow \exists y_1 \forall y_2 \cdots R(x, y_1, y_2, \dots) \vee \exists z_1 \forall y_2 \cdots S(x, z_1, z_2, \dots) \\ &\Leftrightarrow \exists y_1 \exists z_1 \forall y_2 \forall z_2 \cdots (R(x, y_1, y_2, \dots) \vee S(x, z_1, z_2, \dots)) \\ &\Leftrightarrow \exists u_1 \forall u_2 \cdots (R(x, (u_1)_0, (u_2)_0, \dots) \vee S(x, (u_1)_1, (u_2)_1, \dots)) \end{aligned}$$

Which is clearly  $\Sigma_n$ . The same argument can be made for  $A \cap B$  and for  $\Pi_n$  sets.

4. This follows by quantifier contraction, in the same way as (3)  
5. Let

$$A = \{x : \exists y_1 \forall y_2 \cdots R(x, y_1, y_2, \dots)\}$$

And let  $B \leq_m A$  via the function  $f$ . Then we have:

$$B = \{x : \exists y_1 \forall y_2 \cdots R(f(x), y_1, y_2, \dots)\}$$

6. We will prove this by induction on  $n$ .

**Base case:** Let  $n = 0$ . Then  $A$  and  $B$  are clearly recursive.

**Induction step:** Now assume that  $n > 0$  and suppose that  $R \in \Sigma_n$ . Our induction hypothesis says that (6) is true for all  $m < n$ . Then by (4) we have that  $B \in \Sigma_n$ . Further we have  $S \in \Pi_{n+1}$  such that the following holds:

$$\begin{aligned} \langle x, y \rangle \in A &\Leftrightarrow (\forall z < y) R(x, y, z) \\ &\Leftrightarrow (\forall z < y) \exists u S(x, y, z, u) \\ &\Leftrightarrow \exists \sigma (\forall z < y) S(x, y, z, \sigma(z)) \end{aligned}$$

We have that  $\sigma$  range is in  $\omega^{<\omega}$ . By the induction hypothesis we have that  $(\forall z < y) S \Pi_{n+1}$  but by the above deduction we must then have that  $A \in \Sigma_n$ .

## 3.2. Fragments of arithmetic

In this section we will use the previous section to look at different fragments of arithmetic. These can be seen as different sub-theories of *Peano Arithmetics*. These different fragments are important, since a natural question to ask is; "is which axioms of arithmetic do we need for proving Gödel's incompleteness theorems or Solovay's completeness theorems?" We will comment on the bit about Solovay's completeness theorem going forward.

We will split the different fragments up in two categories:<sup>1</sup>

**Strong fragments:** The fragments that contain some induction axiom.

**Weak fragments:** The fragments which do not contain any induction axioms.

We will start of by defining a very weak fragment called Robinson's theory. This fragments was first considered in [Ref]. We will denote this theory with  $\mathbf{Q}$ . This theory have the following axioms:

$$\begin{aligned} &\forall x(\neg Sx \neq 0) \\ &\forall x\forall y(Sx = Sy \rightarrow x = y) \\ &\forall x(x \neq 0 \rightarrow \exists y(Sy = x)) \\ &\forall x(x + 0 = x) \\ &\forall x\forall y(x + Sy = S(x + y)) \\ &\forall x(x \cdot 0 = 0) \\ &\forall(x \cdot Sy = x \cdot y + x) \end{aligned}$$

This theory dose not have the inequality symbol. We will extend  $\mathbf{Q}$  with the following axiom:

$$x \leq y \leftrightarrow \exists z(x + z = y)$$

This extension of  $\mathbf{Q}$  is denoted by  $\mathbf{Q}_{\leq}$ . Both  $\mathbf{Q}$  and  $\mathbf{Q}_{\leq}$  are very weak fragments since they do not contain any induction schemas.

**Definition 3.2** Giving a class  $\Gamma$  of either  $\Sigma_n$  or  $\Pi_n$  formulas we define  $\Gamma$ -*induction* ( $\Gamma$ -IND) to be the following schema:

$$\varphi(0) \wedge \forall v(\varphi(x) \rightarrow \varphi(Sx)) \rightarrow \forall v\varphi(v)$$

---

<sup>1</sup>These definitions are not canonocal. Buss splits them up in three: strong fragments, weak fragments and very weak fragments. But for the scope of this project, we will only need two categories.



for  $\varphi \in \Gamma$ . Further we define  $\Gamma$ -Least Number Principle ( $\Gamma$ -LNP) to be the following schema:

$$\exists x \varphi(x) \rightarrow \exists (\varphi(x) \wedge \neg \exists y (y < x \wedge \varphi(y)))$$

For  $\varphi \in \Gamma$ . Lastly we will define the replacement axioms for  $\Gamma$ , ( $\Gamma$ -REPL) as the following formulas:

$$(\forall x \leq t) \exists y \varphi(x, y) \rightarrow \exists z (\forall x \leq t) (\exists y \leq z) \varphi(x, y)$$

?

—

From the above axioms we can create a hierarchy of different strong fragments of arithmetics. We will define the following theories:

**Definition 3.3** We define the following strong fragments:

1. The theory **EA** is the theory axiomatized by the axioms of  $\mathbf{Q}_{\leq}$  and the induction over bounded formulas  $F(x, \vec{a})$ .
2. The theory  $I\Sigma_n$  [ $\Pi_n$ ] is the theory that is axiomatized by the axioms of  $\mathbf{Q}_{\leq}$  and the  $\Sigma_n$  [ $\Pi_n$ ]-IND axioms.
3. The theory  $I\Delta_0$  is  $\mathbf{Q}_{\leq}$  plus the  $\Delta_0$ -IND axioms.
4. The theory  $L\Sigma_n$  is defined by the theory  $I\Delta_0$  plus the  $\Sigma_n$ -LNP axioms
5. The theory  $B\Sigma_n$  is defined by the theory  $I\Delta_0$  plus the  $\Sigma_n$ -REPL axioms.
6. The theory  $I\Sigma_n^R$  is defined as the closure of  $\mathbf{Q}_{\leq}$  under the  $\Sigma_n$  induction rule:

$$\frac{F(0), \forall x (F(x) \rightarrow \varphi(x+1))}{\forall x F(x)}$$

7. The theory  $I\Sigma_n^-$  [ $\Pi_n^-$ ] is the theory axiomatized by the axioms of  $\mathbf{Q}_{\leq}$  and the schema of induction of  $\Sigma_n$  [ $\Pi_n$ ] functions  $f(x)$ , which only has  $x$  as a free variable. Lastly the theory Peano arithmetics is defined as the theory  $\mathbf{Q}$  plus induction for all first order formulas.

—

Going forward the most important of these theories is  $I\delta_0$ , since we will later augment and for this agumention it can be shown that Solovay's completeness theorems holds. It can be shown that in  $\mathbf{Q}_Q$  a lot of the basic facts about arithmetic can be shown, and these facts holds in all the exetions of  $\mathbf{Q}$ . These facts will not be shown here, but a list of them can be found in [Buss].

### 3.3. PRA and the other fragments of Arithmetics

Having introduced the arithmetical hierarchy in the previous section, it is not obvious where to place **PRA** in this hierarchy. We know that **PRA** has  $\Sigma_1$  induction by axiom 1.18.2.c, so it. But the following theorem can be shown.

**Proposition 3.2** **PRA** is the equivalent to the theory  $I\Sigma_1^R$

The proof of this is non-trivial and will go way beyond the scope of this project. The statement will not be used going forward, and its just included to give some intuition about the placement of **PRA** in the arithmetical hierarchy. The proof can be found in (S. N. Artemov and Beklemishev n.d.). This makes it possible to place **PRA** in the arithmetical hierarchy. We can make a visualization of how all the different fragments relate to each other:




Figure 3.1.: How the fragments relate to each other.

The figure also how the other fragments relate to each other, this will not be shown.

We will now turn to another version of the Selection Theorem.

{thm:sel}

**Theorem 3.1** Let  $\tau v_0 \dots v_{n-1} v_n$  be  $\Sigma_1$ . There is then a  $\Sigma_1$ -formula  $\text{Sel}(\tau)$  with exactly the same free variables such that:

1.  $\text{Sel}(\tau) v_0 \dots v_n \rightarrow \tau v_0 \dots v_n$

2.  $\text{Sel}(\tau)v_0 \dots v_{n-1}v_n \wedge \text{Sel}(\tau)v_0 \dots v_{n-1}v^* \rightarrow v_n = v^*$
3.  $\exists v_n \tau v_0 \dots v_{n-1}v_n \rightarrow \exists v_n \text{Sel}(\tau v_0 \dots v_{n-1}v_n$

Further this can be proven in **PRA**.

*Proof.* Since  $\tau$  is  $\Sigma_1$  it is recursively enumerable. So we have that:

$$\tau v_0 \dots v_n : \exists v (f v v_0 \dots v_n = \bar{0})$$

⊥

### 3.3.1. Induction In PRA

We know that we can do induction on  $\Sigma_1$  formulae in **PRA**. A natural question to ask is then what other kinds of formulae can we do induction on in **PRA**? The following prop will give an answer to this question:

**Proposition 3.3** Over **EA** the following schemata are equivalent:

1.  $\Sigma_n\text{-Ind}$
2.  $\Pi_n\text{-Ind}$
3.  $\Sigma_n\text{-LNP}$
4.  $\Pi_n\text{-LNP}$

It shall be noted that we will only use the case of  $n = 1$  going forward.

*Proof.* The implications (1)  $\Rightarrow$  (4) and (2)  $\Rightarrow$  (3) was proven in proof of theorem 3.1. The converse of these are done in the same way.

We will now show that (1)  $\Rightarrow$  (2). The (2)  $\Rightarrow$  (1) is similar. We will prove the case where  $n = 1$ , and the cases where  $n > 1$  is identical, *modulo* the closure of  $\Sigma_n$  under bounded quantification.

So assume  $\Sigma_1\text{-IND}$  and suppose for  $Fv \in \Sigma_1$  that the following instance of  $\Pi_1\text{-IND}$  fails:

$$(\neg F(\bar{0}) \wedge \forall v (\neg F(v) \rightarrow \neg F(Sv))) \rightarrow \forall v \neg F(v)$$

This means that we have  $\neg F(\bar{0})$ ,  $\forall v (\neg F(v) \rightarrow \neg F(Sv))$  and  $\exists v F(v)$ . We chose  $v_0$  such that  $F(v_0)$ . We will use  $\Sigma_1$  induction on the variable  $v$  in  $F(v_0 \dot{-} v)$  to prove that  $F(\bar{0})$  and get a contradiction.

**Base step:** It is clear that  $F(v_0 \dot{-} \bar{0})$ , since this is just  $F(v_0)$ .

**Induction step:** Assume that  $F(v_0 \dot{-} v)$  is true. We have that  $S(v_0 \dot{-} S(v)) = v_0 \dot{-} v$  unless we already have that  $v = v_0$ , in which case we have that  $v_0 \dot{-} S(v) =$

### 3. The Arithmetical Hierarchy and Fragments of Arithmetics

---

$v_0 - v$ . So we can conclude that  $\forall v F(v_0 \dot{-} v)$  and thus we have  $F(\bar{0})$  and get our contradiction.

The converse of i.e (2)  $\Rightarrow$  (1) is done in a similar way.  $\dashv$

This means that **PRA** can do induction on  $\Pi_1$  formulas, since it has induction for  $\Sigma_1$  induction. It also tells us that **PRA** has the least number principle for  $\Sigma_1$  and  $\Pi_1$  formulas. For **PRA** the following theorem can be proven, but the proof will not be given here. But we will use theorem.

**Theorem 3.2** **PRA**  $\vdash$   $\text{Bool}(\Sigma_1) - \text{Ind}$  where  $\text{Bool}(\Sigma_1)$  is the class of combinations of  $\Sigma_1$  formulae.

We will use these results in chapter 6, since we will need to do induction on a few complicated formulae inside of **PRA**.

### 3.4. Further extensions of the arithmetical hierarchy

If we have that  $I\Gamma$  is a fragment of Peano Arithmetics, then we can look at the fragment  $I\Gamma + \text{EXP}$  which proves induction over  $\Gamma$ -formulas and proves that for all  $x$  the its power  $2^x$  exists. We can also look at  $I\Gamma + \Omega_1$  which is a weaker theory than  $I\Gamma + \text{EXP}$ . Here  $\Omega_1$  is a axiom that asserts that for all  $x$  its power  $x^{\log(x)}$  exists. It is also clear that since **PRA**  $= I\Sigma_1^R$  that the two fragments  $I\sigma_1 + \text{EXO}$  and  $I\sigma_1 + \Omega_1$  are weaker than **PRA**. We will return to these two fragments later on in chapter 6.

It should here be stated that if look at theories  $T$  such that  $Q_{\leq} \subseteq T$ , then it will be possible to encode the syntax of that theory in a similar way to what we have done with **PRA**; i.e in a way such that the proof predicate of that theory  $\text{Pr}_T(x)$  is a  $\Sigma_1$  formula. We will call such a theory for a *RE-theory*. We can then derive all the incompleteness result that we have obtained for **PRA** in that theory. Sometimes a little *tweaking* of the encode should be made, but we can get the following result that will not be proven here:

**Theorem 3.3** Let  $T$  be a *RE-theory* such that **PRA**  $\subseteq T$ . Then the following holds:

1. For any sentences  $F$  and  $G$  we get Löbs derivability conditions
2. Gödel's incompleteness theorems for any sentence such that **PRA**  $\vdash F \leftrightarrow \neg \text{Pr}_T(\ulcorner F \urcorner)$
3. Löb's theorem for any sentence  $F$

This shows that the provability predicate of such fragments  $T$  behaves in the same way as the one from **PRA**.

## 3.5. Sigma Completeness

We can make a generalization of  $D3$ . We use this generalization in some of our proofs later on. We will first need the following lemma which will not be proven:

**Lemma 3.1** Let  $Fv_0 \dots v_{n-1}$  be a  $\Sigma_1$ . Then there is a recursively enumerable formula such that

$$\mathbf{PRA} \vdash Fv_0 \dots v_{n-1} \leftrightarrow \exists v(fvv_0 \dots v_{n-1} = \bar{0})$$

{thm:DemoSig}

**Theorem 3.4** (Demonstrable  $\Sigma_1$  completeness) Let  $Fv_0 \dots v_{n-1}$  be a  $\Sigma_1$  formula with free variables. Then:

$$\mathbf{PRA} \vdash Fv_0 \dots v_{n-1} \rightarrow \text{Pr}(\ulcorner f\dot{v}_0 \dots \dot{v}_{n-1} \urcorner)$$

This theorem states that if a  $\Sigma_1$  formula  $F$  is true, then it can be proven to be true in **PRA** i.e **PRA** is complete with respect to  $\Sigma_1$  formulae.

*Proof.* By [ja, fra hvad?] we have that there is a  $\Sigma_1$  formula  $\exists v(gvv_0 \dots v_{n-1} = \bar{0})$  such that we have:

$$\mathbf{PRA} \vdash fv_0 \dots v_{n-1} \leftrightarrow \exists v(gvv_0 \dots v_{n-1} = \bar{0}) \quad (3.1) \quad \{\text{eq:Com1}\}$$

and by  $D1$  we have:

$$\mathbf{PRA} \vdash \text{Pr}(\ulcorner fv_0 \dots v_{n-1} \leftrightarrow \exists v(gvv_0 \dots v_{n-1} = \bar{0}) \urcorner) \quad (3.2) \quad \{\text{eq:Com2}\}$$

We can now make the following deductions:

$$\begin{aligned} \mathbf{PRA} \vdash fv_0 \dots v_{n-1} &\rightarrow \exists v(gvv_0 \dots v_{n-1} = \bar{0}) && \text{By 3.1} \\ &\rightarrow \exists v \text{Pr}(\ulcorner hvv_0 \dots v_{n-1} = \bar{0} \urcorner) && 1.2 \\ &\rightarrow \text{Pr}(\ulcorner \exists v(hvv_0 \dots v_{n-1} = \bar{0}) \urcorner) && \text{By D1 and D2} \\ &\rightarrow \text{Pr}(\ulcorner fv_0 \dots v_{n-1} \urcorner) && \text{By 3.2} \end{aligned}$$

—

This theorem gives a difference between  $\Sigma_1$  and  $\Pi_1$  sentences in **PRA**. If  $F$  is  $\Pi_1$  we can have that both  $F + \text{Con}(\mathbf{PRA} + F)$  and  $F + \text{Con}(\mathbf{PRA} + \neg F)$  are consistent, but if  $F$  is  $\Sigma_1$  we get that  $F + \text{Con}(\mathbf{PRA} + \neg F)$  is inconsistent.

In the next two chapter we will turn to prove some results about the modal logic **GL**. The next results in chapter 4 will be important for the proof of Solovay's completeness theorems, while the results in chapter 5 will important for using Solovay's completeness theorems to gain results about formulae of arithmetic. This

### 3. The Arithmetical Hierarchy and Fragments of Arithmetics

---

means that the results from chapter 2 and 3 will first be used in chapter 6, where they will be crucial in the proof of Solovay's theorems, but they do not play any role for the "pure" modal logic side of things.

## 4. General Results on GL

In this section we will prove some different properties about the modal logic **GL**, that was introduced in chapter 1. These properties will be used in the following chapters. Some of these properties might hold in other modal logics, but we will look at **GL** as our base logic, instead of a more common modal logic like **K**. Further in the end we will introduce the system **GLS**.

This section will follow (Smorynski 1985). But there will be a some small differences. Smorynski calls the logic **GL** for **PRL** which stands for *Provability Logic*, and he denotes the logic we have called **GLS** by **PRL** <sup>$\omega$</sup> .<sup>1</sup>

We start of by proving the finite tree theorem for **GL**. This theorem makes it possible for us to assume that for a Kripke model  $\mathcal{K} = \langle W, R, \phi \rangle$  of **GL**, that the set  $W$  is finite. For proving this theorem we will first need the to prove this theorem we will first need some knowledge about trees and prove König's lemma.

After this we will prove the continuity theorem which we will use a lot in chapter 5, for the proof of the fixed point theorem.

We will end this chapter by defining the logic **GLS** and show some few results about this logic.

This section (and the next one) dose not use the preceding sections about recursive functions. We will only focus on modal logic. In chapter 6 we will combine all three parts; the recursion theory, the arithmetics and the modal logic.

Finiteness of  $W$  in such a model?

### 4.1. Tress and GL

It is known that **GL** is weakly complete with respect to conversely well-founded transitive frames. There is another result that is in the same vein; **GL** is complete with respect to finite tress. The goal of this section is to prove this result. We will first start of by defining the notion of a tree and a few results about these.

#### 4.1.1. Trees

In this subsection the notion of a special type of graphs called tress will be introduced. The goal of this section is to prove Königs lemma about trees. We will

---

<sup>1</sup>He also have a slightly different definition of a Kripke model.

## 4. General Results on GL

---

start of by defining the notion of a tree.

**Definition 4.1** A *tree* is a tuple  $\mathcal{T} = \langle W, R, w_0 \rangle$  where  $(W, R)$  where:

1.  $<$  is transitive and asymmetric.
2.  $w_0$  is the minimal element of  $<$ . I.e  $w_0 < w$  for all  $w \in W$ .
3. The set of predecessors of any element is finite and linearly ordered by  $<$ .

A tree is *infinite* if the set  $W$  is infinite. ←

We will state some immediate definitions, that should be known from the course *Diskret matematik*:

**Definition 4.2** The *nodes* of a tree  $\mathcal{T} = \langle W, R, w_0 \rangle$  are the elements of the set  $W$ .  
If  $w, v \in W$  are nodes and  $wRv$  then we say that they are joined by an *edge*.

A *walk* is an altering series of vertices and edges in which for each  $v$  we have that  $wRv$  for the  $w$  immediate before it. We call a walk for a *trail*, if all the edges in the walk are distinct. A *path* is trail in which all the nodes are different. ←

It is know from the course *Diskret matematik* that between any two vertices there are exactly one path in a tree. As the concept of a tree is assumed know, we will not look at any concrete examples, We will need some further definitions to state and prove Könings lemma:

**Definition 4.3** Let  $w \in W$ . The *degree* of  $w$  in  $\mathcal{T}$  is the number of edges incident to it. ←

**Definition 4.4** Let  $w$  be a node of a tree  $\mathcal{T}$ , we then define the map  $\pi : \mathcal{T} \setminus \{w_0\} \rightarrow \mathcal{T}$  in the following way:

$$\pi(w) = \text{The node adjacent to } w \text{ on the path to } w_0$$

We call  $\pi(w)$  the *parent node* of  $w$ .

The *child nodes* of  $w$  are the elements of the set  $\{v \in \mathcal{T} : \pi(v) = w\}$ , i.e the child nodes of  $w$  are all the nodes of  $\mathcal{T}$  to which  $w$  is the parent. If  $w$  has no child nodes it is called a *leaf node*. ←

It is clear that if  $w$  is a leaf node, then degree of  $w$  is 1.

**Definition 4.5** A subset  $\Gamma$  of a tree  $\mathcal{T}$  is called a *branch* if and only if all the following conditions holds:

1.  $w_0 \in \Gamma$



2. The parent node of each  $w \in \Gamma \setminus \{w_0\}$  is in  $\Gamma$
3. Each node in  $\Gamma$  is either a leaf node in  $\mathcal{T}$  or has exactly one child node in  $\Gamma$ .

A branch  $\Gamma$  is called *infinite* if and only if it has no leaf node at the end.  $\dashv$

**Definition 4.6** A *fork* of a  $\mathcal{T}$  is a node of  $\mathcal{T}$  which is the end point of two or more branches.  $\dashv$

**Definition 4.7** A tree  $\mathcal{T}$  is called a *finitely branching tree* if every node of  $\mathcal{T}$  has finitely many child nodes.  $\dashv$

The proof of Königs lemma will depend on the following axiom:

**Axiom 4.1** (Axiom of Dependent Choice) Let  $R$  be an relation on a set  $W$  and suppose that:

$$\forall w \in W \exists v \in W : wRv$$

Then there exists a sequence  $(w_n)_{n \in \omega} \in W$  such that

$$\forall n \in \omega : w_n R w_{n+1}$$

This axiom is a weak form of the axiom of choice.

In the proof of Köning's lemma we will look at the subset of all nodes with infinitely many descendants and the relation  $R$  where we have that for nodes  $w, v \in W$ :  $wRv$  if and only if  $v$  is a child of  $w$ . This relation and set fulfills the conditions of the axiom and we can thus create the sequence  $(w_n)_{n \in \omega}$ .

We can now state Köning's lemma:

**Lemma 4.1** (Köning's lemma) Let  $\mathcal{T}$  be a finitely branching tree. Then it is infinite if and only if has an infinite path.

{lem:kong}

*Proof.* If  $\mathcal{T}$  has an infinite path, then it is trivial true that  $\mathcal{T}$  is infinite.

For the other way, assume that  $\mathcal{T}$  is infinite. To show that  $\mathcal{T}$  has an infinite path is the same as showing that it has an infinite branch. We will show that there is a sequence of nodes  $w = (w_0, w_1, w_2, \dots)$  in  $\mathcal{T}$  such that the following holds:

1.  $w_0$  is the root node
2.  $w_{n+1}$  is a child node of  $t_n$
3. each  $w_n$  has infinitely many descendants (definer dette, selvom det giver god mening).

## 4. General Results on GL

---

Then  $w$  is a branch of infinite length, and thus  $\mathcal{T}$  has an infinite path.

We start with the root node  $w_0$ . It has a finite number of child nodes. Suppose for contradiction that each of these child nodes had a finite number of descendants, but thus would mean that  $w_0$  would have a finite number of descendants and thus that  $\mathcal{T}$  would be finite. Therefore  $w_0$  has a child node with infinitely many descendants; let  $w_1$  be one of those.

Now suppose that node  $w_k$  has infinitely many descendants. By the same argument as before we get that  $w_k$  has at least one child node with finitely many descendants. Let  $w_{k+1}$  be one of these. We have thus shown how the sequence  $w$  can be constructed, and the lemma follows by the axiom of dependent choice  $\dashv$

It is clear that finite trees are conversely well-founded frames. This fact is crucial in the proof of theorem in the next subsection, that can be seen as strengthening of the weak completeness theorem of chapter 1:

### 4.1.2. The Finite Tree Theorem

The following theorem is a strengthening of the weak completeness theorem for **GL**, since trees are a subset of the transitive conversely well-founded models and we by this theorem get that we can with out loss of generality assume that our model is finite. This last feature will be crucial in the proof of main theorems of the next chapter.

**Theorem 4.2 (The Finite Tree Theorem)** Let  $\alpha$  be a modal formula. Then the following are equivalent:

1.  $\vdash_{\mathbf{GL}} \alpha$
2.  $\alpha$  is true in all models on finite trees
3.  $\alpha$  is valid in all models on finite trees

With this theorem it is possible to only consider finite frames. This will make the proof of our main theorem of chapter 6 possible. We will call the class of finite trees for FT, and call a formula valid in this class of models for FT-valid.

*Proof.* The implications (1)  $\Rightarrow$  (2) and (1)  $\Rightarrow$  (3) follows by the completeness theorem. We also have that (2)  $\Leftrightarrow$  (3) is true [Kom med et kort argument]. Thus we will just have to show (2)  $\Rightarrow$  (1)

We will show this by contraposition, i.e  $\neg(1) \Rightarrow \neg(2)$ . Assume that  $\not\vdash_{\mathbf{GL}} \alpha$  and let  $\mathcal{K}$  be the following model  $\mathcal{K} = \langle W, R, w_0, \phi \rangle$  be a counter model i.e  $\not\models_{w_0}^{K'} \alpha$ . Let  $S$  be the set of subformulas of  $\alpha$  The goal is now to define a finite tree model:  $\mathcal{K}_T = \langle W_T, <_T, \phi_T \rangle$ . We will do this by letting  $W_T$  consists of finite  $R$ -increasing sequences form  $K$ .

**Stage 0:** Let the sequence  $w_0$  be a part of  $K_T$ .

**Stage  $n + 1$ :** For each sequence  $(w_0, \dots, w_n) \in W_T$  we will look at  $\Gamma = \{\Box\beta \in S(\alpha) : w_n \in \phi(\Box\beta)\}$ . If  $\Gamma = \emptyset$  then we do not extend the sequence  $(w_0, \dots, w_n)$ . Otherwise we will for each  $\Box\beta \in \Gamma$  choose a node  $v \in W$  such that  $w_n R v$  and such that we have:

$$v \in \phi(\Box\beta), v \notin \phi(\beta)$$

We can do this because of axiom 3. We will then add the sequence  $(w_0, \dots, w_n, v)$  to the set  $W_T$

The model  $\mathcal{K}_T$  then further consists of  $<_T$  is the strict ordering by extension of finite sequences.  $\phi_T$  is the defined in the following way:

$$(w_0, \dots, w_n) \in \phi_T(p) \Leftrightarrow w_n \in \phi(p)$$

This leads way to the following notation that we will use in the rest of the proof:

$$\models_{(w_0, \dots, w_n)}^{\mathcal{K}_T} p \Leftrightarrow \models_{w_n}^{\mathcal{K}} p$$

We will now prove two claims, by which the theorem will follow

**Claim 1:**  $\mathcal{T} = (W_T, <_T, w_0)$  is a finite tree with origin  $(w_0)$ . Finiteness follows from lemma 4.1, since the  $\mathcal{T}$  is finitely branching since the branches correlated with the elements of  $S$ . Further there is no infinite paths in  $\mathcal{T}$ , since when we go from  $(w_0, \dots, w_n)$  to  $(w_0, \dots, w_n, w_{n+1})$  we one sentence from  $\Box\beta \in S$  gets forced by  $w_{n+1}$  and since these are finite, the process will stop at some point, and thus the path will be finite.

**Claim 2:** For all  $\beta \in S(\alpha)$  and for all  $(w_0, \dots, w_n) \in W_T$  we have:

$$\models_{(w_0, \dots, w_n)}^{\mathcal{K}_T} \beta \Leftrightarrow \models_{w_n}^{\mathcal{K}} \beta$$

This proof is done by induction on the complexity of  $\beta$ . We will only look at the case  $\beta = \Box\gamma$ . So we have by the induction hypothesis:

$$\begin{aligned} \models_{w_0} \Box\gamma &\Rightarrow \forall v (w_0 R v \Rightarrow \models_v \gamma) \\ &\Rightarrow \forall v ((w_0, \dots, w_n, v) \in W_T \Rightarrow \models_v \gamma) \\ &\Rightarrow \forall v ((w_0, \dots, w_n, v) \in W_T \Rightarrow \models_{(w_0, \dots, w_n, v)}^{\mathcal{K}_T} \gamma) \end{aligned}$$

And the last line is the same as  $\models_{(w_0, \dots, w_n)} \Box\gamma$  For the other way, we will use

## 4. General Results on GL

contraposition:

$$\begin{aligned} \not\models_{w_0} \Box \gamma &\Rightarrow \exists v (w_0 R v \ \& \ \not\models_v \gamma) \\ &\Rightarrow \exists v ((w_0, \dots, w_n, v) \in W_T \ \& \ \not\models_v \gamma) \\ &\Rightarrow \exists v ((w_0, \dots, w_n, v) \in W_T \ \& \ \not\models_{(w_0, \dots, w_n, v)}^{\mathcal{K}_T} \gamma) \end{aligned}$$

And the last line is the same as  $\not\models_{(w_0, \dots, w_n)} \Box \gamma$ .

The theorem now follows since we have:

$$\not\models_{w_0} \alpha \Rightarrow \not\models_{[w_0]}^{\mathcal{K}_T} \alpha$$

⊢

From this theorem there follows a number of interesting corollaries:

**Corollary 4.1** **GL** is decidable, i.e there is an effective procedure for each modal formula  $\alpha$  that determines if  $\vdash_{\mathbf{GL}} \alpha$  or  $\not\vdash_{\mathbf{GL}} \alpha$ .

{cor:Nec}

**Corollary 4.2** For all formulas  $\alpha$  we have:

$$\vdash_{\mathbf{GL}} \alpha \Leftrightarrow \vdash_{\mathbf{GL}} \Box \alpha$$

From this theorem, we can also find the minimum element of a Kripke model of **GL**. This leads to the following definition:

**Definition 4.8** Let  $\mathcal{K} = \langle W, R, \phi \rangle$  be a Kripke model. A *pointed Kripke model* is a pair  $\langle \mathcal{K}, w_0 \rangle$  where  $w_0$  is a node of  $W$ . In the rest of this project we will have that  $w_0$  is the minimum node of the tree of **GL**. We will often just define a pointed Kripke as  $\mathcal{K} = \langle W, R, \phi, w_0 \rangle$ . ⊢

### 4.2. The Continuity Theorem

In this section the continuity theorem and a corollary of it will be stated and proven. This section applies to all Kripke model, and thus is not only about **GL**.

We will first need to definitions:

**Definition 4.9** We define  $d(\alpha)$  in the following way:  $d(p) = d(\perp) = 0$ ,  $d(\alpha \rightarrow \beta) = \max(d(\alpha), d(\beta))$  and  $d(\Box(\alpha)) = d(\alpha) + 1$ . So  $d(\alpha)$  is the maximal number of nested occurrences of  $\Box$  in  $\alpha$ . We call  $d(\alpha)$  the modal degree of  $\alpha$ . ⊢

**Definition 4.10** Let  $R$  be a relation on a set  $W$ . For each  $i \in \omega$  define  $R^i$  as follows:  $R^0$  is the identity relation on  $W$ .  $R^{i+1} = \{ \langle w, v \rangle : \exists v' (w R^i v' \wedge v' R v) \}$ . This  $R^1 = R$  and  $w R^n v$  if and only if  $\exists v_0, \dots, v_n (w = v_0 R \dots R v_n = v)$ . ⊢

{thm:conti}

**Theorem 4.3** (The Continuity Theorem) Let  $\mathcal{K} = \langle W, R, \phi \rangle$  and  $\mathcal{K}' = \langle V, S, \phi' \rangle$  be models and let  $w \in W$ . Let  $P \subseteq \Phi$  be a set of proportional letters. Suppose that  $d(\alpha) = n$ , that all proportional letters that occur in  $\alpha$  are in  $P$ ,  $\{v : \exists i \leq n \, wR^i v\}$ ,  $S = \{\langle v, v' \rangle : x, y \in V \wedge vRv'\}$ , and  $p \in \phi'(v)$  if and only if  $p \in \phi(v)$  for all  $v \in V$  and all proportional letters in  $P$ . Then  $\models_w^\mathcal{K} \alpha$  if and only if  $\models_w^{\mathcal{K}'} \alpha$ .

*Proof.* We will show that for all subformulas  $\beta$  of  $\alpha$ , if we for some  $i$  have that  $wR^i v$  and  $d(\beta) + i \leq n$  such that  $i \leq n$  and  $v \in V$ , then  $\models_v^\mathcal{K} \beta$  if and only if  $\models_v^{\mathcal{K}'} \beta$ . Since  $wR^0 w$  and  $d(\alpha) = 0$  the theorem will follow.

The proof will be induction on the complexity of  $\beta$ . The cases where  $\beta$  is  $\perp$  or a  $p \in P$  is trivial. If  $\beta$  is  $\gamma \rightarrow \sigma$  then  $d(\gamma), d(\sigma) \leq d(\beta)$  and the result follows the induction hypothesis. If  $\beta$  is  $\Diamond \gamma$ ,  $wR^i v$  and  $d(\beta) + i \leq n$ . Then  $v \in V$  and  $d(\beta) = d(\gamma) + 1$ . If  $vRv'$  then  $wR^{i+1}v'$ ,  $d(\gamma) + i + 1 \leq n$ ,  $v' \in V$  and therefore  $vSv'$  and by the induction hypothesis we get  $\models_{v'}^\mathcal{K} \gamma$  if and only if  $\models_{v'}^{\mathcal{K}'} \gamma$ , since we have that  $S \subseteq R$ ,  $vRv'$  if  $vSv'$ . But this means that  $\models_v^\mathcal{K} \beta$  if and only if for all  $v'$  such that  $vRv'$  we have that  $\models_{v'}^\mathcal{K} \gamma$ ; if and only if for all  $v'$  such that  $vSv'$ :  $\models_{v'}^\mathcal{K} \gamma$  if and only if by the induction hypothesis for all  $v'$  such that  $vSv'$ ;  $\models_{v'}^{\mathcal{K}'} \gamma$  if and only if  $\models_v^{\mathcal{K}'} \beta$ .  $\dashv$

From this theorem the following corollary is a immediate consequence and we will use it to prove parts of the main theorem of the next chapter:

{cor:conti}

**Corollary 4.3** Let  $\alpha$  be a formula. Let  $\mathcal{K} = \langle W, R, \phi \rangle$  and  $\mathcal{K}' = \langle V, R, \phi' \rangle$  be models, and  $p \in \phi(w)$  if and only if  $p \in \phi'(w)$  for all  $w \in W$  and all  $p$  contained in  $\alpha$ . Then  $\models_w^\mathcal{K} \alpha$  if and only if  $\models_w^{\mathcal{K}'} \alpha$ .

## 4.3. The Modal Logic **GLS**

In this section we will define the modal logic **GLS** (The "S" is for Solovay). This logic is an extension of **GL**, but it is not as well behaved as **GL**. It is this logic we will use in the statement of Solovay's second completeness theorem, so it is important for each to know a few basic results about it.

### Definition 4.11 Dobbelt tjek i Per

The modal logic **GLS** is the logic which has the following axioms

GL All theorems of **GL**.

Refl  $\Box p \rightarrow p$

and which sole rule of inference is modus ponens.

$\dashv$

## 4. General Results on GL

Another more cumbersome definition of this modal logic can be found in Appendix A.

It should be noted that this modal logic is not a *normal* one, since it do not have necessitation as a rule of inference. If a modal logic just extended **GL** with the axiom *Refl* then this logic would prove both  $\neg\Box\perp$  and  $\Box\perp$  and thus be inconsistent; this is why we have not included it as rule. But in some way it still has necessitation as a rule, since it contains all the theorems of **GL**.

There is no completeness theorem for **GLS** in the same vein as the finite tree theorem for **GL** (This theorem is found in (Lindström 1997)). But there is another similar result we will now state and prove.

**Definition 4.12** Let  $\alpha$  be any modal formula, let  $\mathcal{K} = \langle W, R, \phi, w_0 \rangle$  be a pointed Kripke model and suppose that  $w \in W$ . Then  $w$  is  $\alpha$ -*reflexive* in  $\mathcal{K}$  if  $\Box\beta \rightarrow \beta \in \phi(w)$  for all  $\Box\beta \in S(\alpha)$ .  $\mathcal{K}$  is  $\alpha$ -reflexive if  $w_0$  is  $\alpha$ -reflexive. Let  $\mathcal{F}$  be a class of Kripke models; a formula  $\alpha$  is *r-valid* in  $\mathcal{F}$  if  $\models^{\mathcal{K}} \alpha$  for all  $\alpha$ -reflexive  $\mathcal{K} \in \mathcal{F}$ .  $\dashv$

If we want to show that a formula  $\alpha$  is not r-valid in the class of finite tree Kripke models, it is enough to find a  $\mathcal{K}$  in this class and a  $\alpha$ -reflexive  $w \in W$  such that  $\alpha \notin \phi(w)$ .

The main theorem about this modal logic is the following, which gives a sort of model theory for **GLS**.

{thm:MainGLS}

**Theorem 4.4** Let  $\alpha$  be a modal formulas. Define the following set:  $S_{\Box}(\alpha) = \{\Box\beta : \Box\beta \text{ is a subformula of } \alpha\}$  the set of subformulas of  $\alpha$  that is boxed. Then the following to statements are equivalent:

1.  $\vdash_{\mathbf{GLS}} \alpha$
2.  $\vdash_{\mathbf{GL}} \bigwedge_{\Box\beta \in S_{\Box}(\alpha)} (\Box\beta \rightarrow \beta) \rightarrow \alpha$
3.  $\alpha$  is r-valid in the class of finite tree Kripke models

This theorem reduces the decidability of **GLS** to the decidability of **GL** and describes a model theory for **GLS**.

We have that (2)  $\Leftrightarrow$  (3) by the finite tree theorem, and that (2)  $\Rightarrow$  (1) is trivial.

The proof that (1)  $\Rightarrow$  (2) is non trivial, and for this proof we need the following lemma:

{lem:hvad}

**Lemma 4.1** Let  $\mathcal{K} = \langle W, R, \phi, w_0 \rangle$  and suppose that  $w_0 R w_1 R \dots R w_{n+1}$ . Let  $\vartheta_n$  be the following formulas:

$$(\Box p_1 \rightarrow p) \wedge \dots \wedge (\Box p_n \rightarrow p_n)$$

Then there is an  $i$ ;  $1 \leq i \leq n + 1$ , such that  $\vartheta_n \in \phi(w_i)$ , and thus  $\models^{\mathcal{K}} \vartheta_n$ .

*Proof.* The lemma follows from the fact that for each  $k$ ;  $1 \leq k \leq n$ , there is at most one  $j$ ;  $1 \leq j \leq n+1$ , such that  $\neg(\Box p_k \rightarrow p_k) \in \phi(w_j)$   $\dashv$

Further for the proof we will make use of the operation on Kripke models:

**Definition 4.13** Let  $\mathcal{K} = \langle W, R, \phi \rangle$  be a Kripke Model. The derived model  $\mathcal{K}'$  is defined as follows:  $W' = W \cup \{w'\}$ , where  $w' \notin W$ . We further have that  $R'$  is defined as  $wR'v$  iff  $WRv$  for  $w, v \in W$ ;  $w'$  is the new minimum element. Lastly we have that  $w \in \phi'(p)$  iff  $w \in \phi(p)$  for  $w \in W$ , where  $p$  is any atomic formula. Further we have that  $w' \in \phi'(p)$  iff  $w_0 \in \phi(p)$ .  $\dashv$

We can further define the notion of a sequences of successive derived models:  $\mathcal{K}^{(1)}, \mathcal{K}^{(2)}, \dots$  that have  $w_1, w_2, \dots$  as their minima. We define  $\mathcal{K}^{(1)} = \mathcal{K}$  and  $\mathcal{K}^{(n+1)} = (\mathcal{K}^{(n)})'$  and let  $w_n$  denote the minimum of  $\mathcal{K}^{(n)}$ . This can graphically been seen by adding a tail of  $n$  nodes below  $w_0$ , as seen in the following graphs:

#### 4. General Results on GL

---




Figure 4.1.: The visualization of the models  $\mathcal{K}$  (left) and  $\mathcal{K}^{(n)}$  (right)

The following lemma will explain a bit deeper how these derived models works, and will be used in the proof of theorem.

{lem:GLS}

**Lemma 4.2** Let  $\mathcal{K}$  be a  $\alpha$ -reflexive model,  $S(\alpha)$  the set of subformulas of  $\alpha$  and  $\mathcal{K}^{(n)}$  be the  $n$ -derived model. Then for all  $\beta \in S(\alpha)$  we have:

$$\models_{w_0} \beta \Leftrightarrow \models_{w_n} \beta$$

*Proof.* We will prove this by induction on  $n$ . The lemma will follow from the case  $n = 1$ , since we can use that result  $n$  times to get the lemma, and this will be proven by induction on complexity of  $\beta$ . The only non trivial case is the case where  $\beta = \Box\gamma$ , so this is the only case that will be shown:

Let  $\beta = \Box\gamma \in S(\alpha)$ .

( $\Rightarrow$ ) The following is clear for  $\Box\gamma \in S(\alpha)$ :

$$\models_{w_0} \Box\gamma \Rightarrow \models_{w_0} \gamma$$

This follows since our model is  $\alpha$ -sound. Therefore we have:  $\forall v > w_1 (\models_{w_1} \gamma)$  and thus  $\models_{w_1} \Box\gamma$

( $\Leftarrow$ ) Here we can make the following deduction:

$$\begin{aligned} \models_{w_1} \Box\gamma &\Rightarrow \forall v > w_1 (\models_v \gamma) \\ &\Rightarrow \forall v > w_0 (\models_v \gamma) \\ &\Rightarrow \models_{w_0} \Box\gamma \end{aligned}$$



⊥

We can now return to our proof of the main theorem of this section by proving

*Proof of 4.4.* We will just have to show that (1)  $\Rightarrow$  (2). Assume that  $\alpha$  is not FT-r-valid, i.e we have that there is an  $\alpha$ -reflexive  $\mathcal{K} = \langle W, R, \phi, w_0 \rangle \in \text{FT}$  such that  $\models^{\mathcal{K}} \neg \alpha$ . Our goal is to show that  $\not\models_{\mathbf{GLS}} \alpha$ .

If we have that  $\vdash_{\mathbf{GLS}} \alpha$  there are formulas  $\gamma_i$ ,  $i < n$  such that if let  $\sigma$  be the following formula

$$\bigwedge \{ \Box \gamma_i \rightarrow \gamma_i : i < n \} \rightarrow \alpha$$

is provable in **GL** and therefor  $\sigma$  is FT-valid. The theorem follows if we can show that  $\sigma$  is not FT-valid. We will look at the derived model  $\mathcal{K}^{n+1}$ , and by 4.1 there is a  $j \leq n$  such that

$$w_{j+1} \in \phi^{n+1}(\bigwedge \{ \Box \gamma_i \rightarrow \gamma_i : i < n \})$$

Since we have that  $w_{j+1} \in \phi^{n+1} \neg \alpha$  and thus  $a_{j+1} \in \phi(\neg \sigma)$  and thus  $\sigma$  is not FT-valid ⊥

Since the models used in this proof is finite we get the following corollary:

**Corollary 4.4** **GLS** is decidable

In the chapter 6 it will be shown that **GL** axiomatise the provable schemata in **PRA** and **GLS** will axiomatise the *true* schemata of arithmetics; i.e truth in the standard model  $\mathcal{N} = \langle \omega, +, \cdot \rangle$  of arithmetics.



## 5. Fixed Point Theorem

In this section we will prove the so called *fixed point theorem* for **GL**. This theorem says that for some formulae  $\alpha$  of **GL** exists fixed points  $\sigma$ , i.e. that for these formulae:

$$\vdash_{\mathbf{GL}} \sigma \leftrightarrow \alpha(\sigma)$$

This theorem gives is one of the most striking applications of modal logic in the study of provability in arithmetic; it says that self-reference is not necessary in the proofs of Gödel's incompleteness theorems. There has been a lot of different ways to prove this theorem; Boolos list three different ways in his book (George. S Boolos 1993) and Per Lindström has another one in his paper (Lindström 1996). Here we will follow the proof from (Reidhaar-Olson 1990).

This proof of the theorem is a semantical proof; i.e. it makes use of the Kripke semantics of the modal logic **GL**, and it further gives a simple procedure for "calculating" fixed points.

We will end this chapter with calculating some fixed points. This will be possible since our the proof of the fixed point theorem contains a algorithm, which can be used to calculate these. In the next chapter we will use Solovay's completeness theorems and the fixed point theorem to conclude a few things about arithmetic with the help of modal logic. This is the *power* of the fixed point theorem; it makes it possible to study the proof predicate with the help of modal logic.

### 5.1. Prerequisite Results and Definitions

We will start of by make stating some definitions and results that are crucial in the proof of the fixed point theorem. We will also in this section define the special formulae for which the fixed point theorem holds.

**Definition 5.1** We abbreviate  $\Box\alpha \wedge \alpha$  as  $\Box\alpha$  for every  $\alpha$  in our language.  $\dashv$

$\Box$  is called "strong box" and for a given model  $\mathcal{K}$  we have that  $\vdash_w^{\mathcal{K}} \Box \alpha$  states that  $\alpha$  is both true and provable in node  $w$ . There are a few theorem about  $\Box$  in the modal logic **K4** and logics which extends this, that will be important in the following. These results are syntactical. We will show these now, and more results can be found in (Smorynski 1985) and George. S Boolos 1993

## 5. Fixed Point Theorem

{seet}

**Theorem 5.1** Let  $\alpha$  be a modal formula then:

1.  $\vdash_{\mathbf{K4}} (\Box\Box\alpha \leftrightarrow \Box\alpha) \leftrightarrow \Box\Box\alpha$
2.  $\vdash_{\mathbf{K4}} \Box\alpha \leftrightarrow \Box\Box\alpha$

*Proof.* We will prove (1). We have that  $\vdash_{\mathbf{K4}} \Box\alpha \rightarrow \Box\Box\alpha$  by the axiom 4, and thus since it is also a normal modal logic we have:

$$\vdash_{\mathbf{K4}} ((\Box\Box\alpha \wedge \Box\alpha) \leftrightarrow \Box\alpha) \leftrightarrow \Box(\Box\alpha \wedge \alpha)$$

And (1) is exactly this.

The proof of (2) is similar.  $\dashv$

{GLsæt}

**Theorem 5.2** Suppose that  $\mathbf{L}$  extends  $\mathbf{K4}$ , and we have that  $\vdash_{\mathbf{L}} \Box\alpha \rightarrow \beta$ . Then  $\vdash_{\mathbf{L}} \Box\alpha \rightarrow \Box\beta$  and  $\vdash_{\mathbf{L}} \Box\alpha \rightarrow \Box\Box\alpha$

This theorem clearly holds for  $\mathbf{GL}$ , and we will use it later in this chapter.

*Proof.* Since  $\mathbf{L}$  extends  $\mathbf{K4}$  we have that  $\vdash_{\mathbf{L}} \Box\Box\alpha \rightarrow \Box\alpha$  and thus by theorem 5.1 we have that  $\vdash_{\mathbf{L}} \Box\alpha \rightarrow \Box\beta$  and by the definition of  $\Box$  we get that  $\vdash_{\mathbf{L}} \Box\alpha \rightarrow \Box\beta$   $\dashv$

Having shown a few results about the syntactical properties of  $\Box$  we will now turn to some of the sementical properties about the relationship between  $\Box$  and  $\Diamond$ .

{rem:acc}

**Remark 5.1** By our semantic of modal logic, we have that  $\models_w \Box\alpha$  is true iff  $\models_v \alpha$  for all  $v \in \{w\} \cup \text{acc}(w)$ , where  $\text{acc}(w)$  is the collection of notes that are accessible from  $w$ , i.e:  $\text{acc} = \{v \in W : wRv\}$

This remark gives us an easy way to determine if  $\Box\alpha$  is true.

Another useful definition that we can make, since we for every Kripke model  $\mathcal{K} = \langle W, R, \phi \rangle$  of  $\mathbf{GL}$  have that  $W$  is a finite set is the following:

**Definition 5.2** Let  $\mathcal{K} = \langle W, R, \phi \rangle$  be a Kripke model of  $\mathbf{GL}$ . The *rank* of the notes  $w \in W$  is defined in the following way:  $\text{rank}(w) = 0$  iff there is no world  $v$  such that  $wRv$ . Otherwise we have that  $\text{rank}(w) = 1 + \max\{\text{rank}(v) : wRv\}$ .  $\dashv$

So  $\text{rank}(w)$  simple states how many worlds that can be seen from the world  $w$ . Since  $W$  is finite and that  $R$  is irreflexive we have that the for each  $w \in W$  the rank of  $w$  is unique.

We will need the following three lemmas in our proof of the Fixed Point Theorem. All three of these lemmas are sementical in nature.

{lem:acc}

**Lemma 5.1** Given any Kripke model  $\mathcal{K} = \langle W, R, \phi \rangle$  of **GL**,  $w \in W$  and sentence  $\alpha$ , we have that if  $\models_w \Box \alpha$  then  $\models_v \Box \alpha$  for any  $v \in \text{acc}(w)$ . Further we have that  $\models_v \Box \alpha$  for all  $v \in \text{acc}(w)$ .

*Proof.* Assume that we have  $\models_w \Box \alpha$ ,  $w$  sees  $v$  and  $v$  sees  $v'$ . Since  $R$  is transitive we have that  $w$  is also connected to  $v'$  and thus we have that  $\models_{v'} \alpha$ . Since  $v'$  was chosen at random we have that  $\models_v \Box \alpha$ . We also have that  $\models_v \alpha$  and thus we have  $\models_v \Box \alpha$ .  $\dashv$

{lem:con}

**Lemma 5.2** Given any Kripke model  $\mathcal{K} = \langle W, R, \phi \rangle$ ,  $w \in W$  and sentence  $\alpha$ , if  $\not\models_w \Box \alpha$  then there is "notes"  $v$  seen by  $w$  such that  $\models_v \Box \alpha$  and  $\not\models_v \alpha$ .

*Proof.* Assume that  $\not\models_w \Box \alpha$  then there is a notes  $v$  connected to  $w$  such that  $\not\models_v \alpha$ . Let  $v$  be the notes with the least rank with this property and suppose that  $vRv'$ . Since  $v'$  is of less rank than  $v$  we have that  $\models_{v'} \alpha$ . Now since  $v'$  was chosen arbitrarily we have that  $\models_v \Box \alpha$  and the lemma is proven.  $\dashv$

The next lemma will be crucial in the proof of the fixed-point theorem; since it enable us to substitute two formulae into another formula.

{lem:sem}

**Lemma 5.3 (Semantic Substitution Lemma)** For any sentences  $\alpha, \beta$  and  $\gamma$  we have that the following formula is valid in all models of **GL**:

$$\Box(\beta \leftrightarrow \gamma) \rightarrow (\alpha(\beta) \leftrightarrow \alpha(\gamma))$$

The meaning of  $\alpha(\beta)$  is that we replace every occurrences of  $p$  in  $\alpha$  with  $\beta$ ; the meaning of  $\alpha(\gamma)$  is similar.

*Proof.* We start of by fixing  $\beta$  and  $\gamma$ . The proof will be by induction on the complexity of  $\alpha$ . We will only proof the part where  $\alpha$  is  $\Box \sigma$ .

So suppose that  $\alpha$  is  $\Box \sigma$ , where  $\Box(\beta \leftrightarrow \gamma) \rightarrow (\sigma(\beta) \leftrightarrow \sigma(\gamma))$  is valid. Let  $\mathcal{K}$  be any model of **GL** and let  $w \in W$ . Suppose that  $\models_w \Box(\beta \leftrightarrow \gamma)$ . Let  $v$  be any world seen by  $w$ . Then by lemma 5.1 we have  $\models_v \Box(\beta \leftrightarrow \gamma)$ . Since  $\Box(\beta \leftrightarrow \gamma) \rightarrow (\sigma(\beta) \leftrightarrow \sigma(\gamma))$  is valid we get that  $\models_v \sigma(\beta) \leftrightarrow \sigma(\gamma)$  and since  $v$  was chosen arbitrary we get  $\models_w \Box(\sigma(\beta) \leftrightarrow \sigma(\gamma))$ . By proposition 1.2 and weak completeness we can conclude  $\models_w \Box \sigma(\beta) \leftrightarrow \sigma(\gamma)$ .  $\dashv$

We are now almost ready to state state and prove the fixed-point theorem. We will just need the next two definitions:

**Definition 5.3** A formula  $\alpha$  is called *modalized in  $p$*  if every occurrence of  $p$  in  $\alpha$  is under the scope of  $\Box$ .  $\dashv$

## 5. Fixed Point Theorem

---

That  $\alpha$  is modalized in  $p$  will be a sufficient condition for  $\alpha$  having a fixed point, but it is not a necessary condition for this.  $\Box p \vee p$  is one example of a formula that has a fixed point, even though  $p$  is not modalized in this formula. This will be showed later.

We will also need the following definition, which in some way is a generalization of being modalized in  $p$ :

**Definition 5.4** A sentence  $\alpha$  is said to be *n-decomposable* iff for some sequence  $q_1, \dots, q_n$  consisting of distinct sentence letters that do not occur in  $\alpha$  we have some sentence  $\beta(q_1, \dots, q_n)$  that do not contain  $p$  and another sequence of distinct sentences  $\gamma_1(p), \dots, \gamma_n(p)$ , which each contains  $p$  that we have

$$\alpha = \beta(\gamma_1(p), \dots, \gamma_n(p))$$

.

⊢

The following remark will be important in the proof of the fixed point theorem.

**Remark 5.2** It should be noted that if  $\alpha$  is modalized in  $p$  that we then have that  $\alpha$  is *n-decomposable* for some  $n$ .

### 5.2. Proof of the Fixed Point Theorem

We can now state the fixed point theorem. But we will still need a few lemmas in our proof. These will be stated and proven when needed. The theorem we will start of by showing can be seen as a uniqueness theorem, we will after this theorem state and prove a theorem that can be seen as a existences theorem. Lastly we will prove a corollary that will show why the name fixed point theorem is appropriate; this is not exactly from the fixed point theorem itself.

{thm:Fixed}

**Theorem 5.3** If  $\alpha$  is modalized in  $p$ , then there exists a formula  $\sigma$  in which the only sentence letters that occurs are these other than  $p$  that occur in  $\alpha$ , and such that:

$$\Box(p \leftrightarrow \alpha) \rightarrow (p \leftrightarrow \sigma)$$

The formula  $\sigma$  is called a *fixed-point* of  $\alpha$ .

We will later on proof that  $\vdash_{\mathbf{GL}} \sigma \leftrightarrow \alpha(\sigma)$  for such a  $\sigma$  and thus the name fixed point is appropriate.

*Proof.* We will prove this by showing that if  $\alpha$  is *n-decomposable* then it has a fixed point. We will show this by induction on  $n$ .

**Base case:** Suppose that  $\alpha$  is 0-decomposable. Then we have that  $p$  dose not occur in  $\alpha$  and it can thus itself be the sentence  $\beta$ .

**Induction step:** Assume that every sentence that is  $n$ -decomposable has a fixed point. We now have to show that every sentence that is  $(n+1)$ -decomposable also has a fixed point. To show this we will assume the following:

$$\alpha(P) = \beta(\Box\gamma_1(p), \dots, \Box\gamma_{n+1}(p))$$

Further for each  $i$  let:

$$\alpha_i(p) = \beta(\Box\gamma_1(p), \dots, \Box\gamma_{i-1}(p), \top, \Box\gamma_{i+1}(p), \dots, \Box\gamma_{n+1}(p))$$

Thus we have that for each  $i$  that  $\alpha(p)$  is  $n$ -decomposable, so it has a fixed point, that we call  $\sigma_i$ . Lastly we define:

$$\sigma = \beta(\Box\gamma_1(\sigma_1), \dots, \Box\gamma_{n+1}(\sigma_{n+1}))$$

Our goal is to show that  $\sigma$  is a fixed point of  $\alpha$ .

{lem:fix}

**Lemma 5.4** For each  $i$  we have that:

$$\vdash_{\mathbf{GL}} \Box(p \leftrightarrow \alpha) \rightarrow \Box(\Box\gamma_i(p) \leftrightarrow \gamma_i(\sigma_i))$$

*Proof.* Since we have that  $\mathbf{GL}$  is complete, we just ave to show that for any model  $\mathcal{K} = \langle W, R, \phi \rangle$  and any  $w \in W$  that:

$$\mathcal{K} \models \Box(p \leftrightarrow \alpha) \rightarrow \Box(\Box\gamma_i(p) \leftrightarrow \Box\gamma_i(\sigma_i)) \quad (5.1) \quad \{\text{eqn:1}\}$$

So we will start of by fixing  $i$ ,  $\mathcal{K}$  and  $w \in W$ . We will show 5.1 by assuming  $\models_w \Box(p \leftrightarrow \alpha)$  and then deduce:  $\models_w \Box(\Box\gamma_i(p) \leftrightarrow \Box\gamma_i(\sigma_i))$ ; this is equivalent to  $\models_v \Box\gamma_i(p) \leftrightarrow \Box\gamma_i(\sigma_i)$  for all  $v \in \{w\} \sup \text{acc}(w)$  by remark 5.1. So let  $v \in \{w\} \cup \text{acc}(w)$  and assume that  $\models \Box\gamma_i(p)$ , i.e  $\models_v \Box\gamma_i(p) \leftrightarrow \top$ . By lemma 5.1 we have that for any  $v' \in \text{acc}(v)$  that  $\models_{v'} \Box\gamma_i(p)$  and thus  $\models_{v'} (\Box\gamma_i(p) \leftrightarrow \top)$ . This means that we have:

$$\models_v \Box(\Box\gamma_i(p) \leftrightarrow \top)$$

And thus by lemma 5.3 we get that  $\models_v \alpha_i \leftrightarrow \alpha$  and since our  $v$  was chosen arbitrarily we have that  $\models_w \Box(\alpha_i \leftrightarrow \alpha)$  and thus by lemma 5.1 we get that:  $\models_v \Box(\alpha_i \leftrightarrow \alpha)$ . Since we have assumed that  $\models_w \Box(p \leftrightarrow \alpha)$  we again have by lemma 5.1 that  $\models_v \Box(p \leftrightarrow \alpha)$ , and hence we have  $\models_v \Box(p \leftrightarrow \alpha_i)$ . Since our logic is complete and we have assumed by the induction hypothesis that  $\alpha_i$  has a fixed point  $\sigma_i$  we have that  $\models_v (p \leftrightarrow \gamma_i)$ , and thus, since  $v$  was chosen arbitrarily we have that  $\models_w \Box(p \leftrightarrow \gamma_i)$ . So by using 5.1 again we get that  $\models_v \Box(p \leftrightarrow \gamma_i)$ . We will now use

## 5. Fixed Point Theorem

lemma 5.3 again and get that:

$$\models_v \gamma_i(p) \leftrightarrow \gamma_i(\sigma_i) \quad (5.2) \quad \{\text{eqn:sub1}\}$$

and

$$\models_v \Box \gamma_i(p) \leftrightarrow \Box \gamma_i(\sigma_i) \quad (5.3) \quad \{\text{eqn:sub2}\}$$

Notice that these two holds for any  $v \in \{w\} \cup \text{acc}(w)$  such that  $\models_v \Box \gamma_i(p)$ . Further by 5.3 we can deduce  $\models_v \Box \gamma_i(p) \rightarrow \Box \gamma_i(\sigma_i)$

For the next step of the prove of this lemma we will assume that  $\not\models_v \Box \gamma_i(p)$ . This means by lemma 5.2 that there is some world  $v'$  where  $v' \in \text{acc}(v)$ , such that  $\not\models_{v'} \gamma_i(p)$  and  $\models_{v'} \Box \gamma_i(p)$ . 5.2 holds for  $v'$  since  $v' \in \{w\} \cup \text{acc}(w)$  and thus we have  $\models_{v'} \gamma_i(p) \leftrightarrow \gamma_i(\sigma_i)$ . This gives that  $\not\models_{v'} \gamma_i(\sigma_i)$  and thus since  $v R v'$  we have that  $\not\models_v \Box \gamma_i(\sigma_i)$ . By contraposition we then get:  $\models_v \Box \gamma_i(\sigma_i) \rightarrow \Box \gamma_i(p)$ , and thus we have shown that:

$$\models_v \Box \gamma_i(p) \leftrightarrow \Box \gamma_i(\sigma_i)$$

We have now shown the lemma.  $\dashv$

We now go back and finish our proof of the fixed point theorem. Suppose that  $\mathcal{K}$  is a model and that  $w \in W$  such that  $\models_w \Box(p \leftrightarrow \alpha)$ . By lemma 5.4 and completeness we get  $\models_w \Box(\Box \gamma_i(p) \leftrightarrow \Box \gamma_i(\sigma_i))$ . By using lemma 5.3  $(n+1)$  times we can deduce that:

$$\models_w \beta(\Box \gamma_i(p), \dots, \Box \gamma_{n+1}(p)) \leftrightarrow \beta(\Box \gamma_1(\sigma_1), \dots, \Box \gamma_{n+1}(\sigma_{n+1}))$$

i.e  $\models_w \alpha \leftrightarrow \sigma$ .

Since we have  $\models_w p \leftrightarrow \alpha$  we get  $\models_w p \leftrightarrow \sigma$ , we can obtain  $\models_w \Box(p \leftrightarrow \alpha) \rightarrow (p \leftrightarrow \sigma)$ . Since our  $\mathcal{K}$  and  $w$  was chosen at random we have that  $\Box(p \leftrightarrow \alpha) \rightarrow (p \leftrightarrow \sigma)$  is valid. By completeness we then have:  $\vdash_{\mathbf{GL}} \Box(p \leftrightarrow \alpha) \rightarrow (p \leftrightarrow \sigma)$   $\dashv$

We can use the algorithm from this prove to find some fixed points of given formulae.

**Example 5.1** Let  $\alpha(p) = \Box \neg p$ . We will calculate the fixed point for  $\alpha$ . Let  $\gamma_1(p) = p$  and  $\beta(q_1) = q_1$ . We then have that  $\alpha(p) = \beta(\Box \gamma_1(p))$  and  $\alpha_1(p) = \beta(\top) = \top$ . The fixed point of  $\sigma_1$  of  $\alpha_1$  is then  $\top$ , since  $p$  does not occur in  $\alpha_1$ . We can thus conclude that the fixed point  $\sigma$  of  $\alpha$  is the following:

$$\beta(\Box \gamma_1(\top)) = \beta(\Box \neg \top) = \Box \neg \top$$

this formula is equivalent to  $\Box \perp$ .  $\dashv$



**Example 5.2** Let  $\alpha(p) = \Box p \rightarrow \Box \neg p$ . To find the fixed point, let  $\gamma_1(p) = p, \gamma_2(p) = \neg p$  and  $\beta(q_1, q_2) = q_1 \rightarrow q_2$ . Then clearly we have:

$$\alpha(p) = \beta(\Box \gamma_1(p), \Box \gamma_2(p))$$

$$\alpha_1(p) = \beta(\top, \Box \gamma_2(p)) = \top \rightarrow \Box \neg p$$

and

$$\alpha_2(p) = \beta(\Box \gamma_1(p), \top) = \Box p \rightarrow \top$$

It is clear that  $\alpha_1$  is equivalent to  $\Box \neg p$  and that  $\alpha_2$  is equivalent to  $\top$ ; thus they have fixed points  $\sigma_1 = \Box \perp$  and  $\sigma_2 = \top$ . So the fixed point  $\sigma$  of  $\alpha$  is the following:

$$\beta(\Box \gamma_1(\Box \perp), \Box \gamma_2(\top)) = \Box \Box \perp \rightarrow \Box \neg \top$$

or equivalent:  $\Box \Box \perp \rightarrow \Box \perp$ . ⊣

The following result follows from the fixed-point theorem.

{thm:exi}

**Theorem 5.4** Let  $\alpha(p)$  be modalized in  $p$ , and let  $\sigma$  be a fixed-point of  $\alpha$ . Then:

$$\vdash_{\mathbf{GL}} \Box(p \leftrightarrow \sigma) \rightarrow (p \leftrightarrow \alpha)$$

*Proof.* Suppose that  $\mathcal{K} = \langle W, R, \phi \rangle$  is a finite transitive and irreflexive model in which  $\Box(p \leftrightarrow \sigma) \rightarrow (p \leftrightarrow \alpha)$  is invalid. This means that for some  $w \in W$  of least rank that we have:  $\models_w \Box(p \leftrightarrow \sigma)$  and thus  $\models_w p \leftrightarrow \sigma$  and  $\not\models_w p \leftrightarrow \alpha$ . If  $wRv$  then  $\models_v \Box(p \leftrightarrow \sigma)$  and since  $x$  of lower rank than  $w$  we also have  $\models_v p \leftrightarrow \alpha$ . Let  $\varphi'$  be like  $\varphi$  expect that  $p \in \phi'(w)$  if and only if not  $p \in \phi(w)$ . Set  $\mathcal{K}' = \langle W, R, \phi' \rangle$ , and this model is clearly transitive and irreflexive.

In the rest of this prove we will use corollary to the continuity theorem; i.e corollary 4.3.

The formula  $\alpha$  is a truth-functional compound of closed formulas  $\Box \beta$  and propositional letters  $q$  such that each  $q$  is not  $p$ . We have that  $\models_w^\mathcal{K} \Box \beta$  if and only if  $\models_v^\mathcal{K} \beta$  for all  $v$  such that  $wRv$ , if and only if  $\models_v^{\mathcal{K}'} \beta$  for all  $v$  such that  $wRv$  (by continuity), if and only if  $\models_w^{\mathcal{K}'} \Box \beta$ . We further have by the definition of  $\mathcal{K}'$  that  $\models_w^\mathcal{K} \alpha$  if and only iff  $\models_w^{\mathcal{K}'} \alpha$  and  $\models_w^\mathcal{K} p$  if and only if not  $\models_w^{\mathcal{K}'} p$ . Thus we have  $\models_w^{\mathcal{K}'} p \leftrightarrow \alpha$  and by the continuity theorem we again get  $\models_v^{\mathcal{K}'} p \leftrightarrow \alpha$  for all  $v$  such that  $wRv$ . But this means that we get  $\models_w^{\mathcal{K}'} \Box(p \leftrightarrow \alpha)$ .

Since  $\sigma$  does not contain  $p$  we get by the conti them that  $\models_w^\mathcal{K} \sigma$  if and only if  $\models_w^{\mathcal{K}'} \sigma$ . Since we have that  $\models_w^\mathcal{K} p$  if and only if not  $\models_w^{\mathcal{K}'} p$  we get  $\not\models_w^{\mathcal{K}'} p \leftrightarrow \sigma$  and thus  $\Box(p \leftrightarrow \sigma) \rightarrow (p \leftrightarrow \sigma)$  is invalid.

By soundness and completeness of  $\mathbf{GL}$  we thus get that if  $\vdash_{\mathbf{GL}} \Box(p \leftrightarrow \alpha) \rightarrow (p \leftrightarrow \sigma)$  then  $\vdash_{\mathbf{GL}} \Box(p \leftrightarrow \sigma) \rightarrow (p \leftrightarrow \alpha)$ . ⊣

## 5. Fixed Point Theorem

---

From this we can prove the following corollary that show that the name fixed-point is appropriate:

**Corollary 5.1** Let  $\alpha(p)$  be modalized in  $p$  and let  $\sigma$  be a fixed-point of  $\alpha$ . Then:

$$\vdash_G \sigma \leftrightarrow \alpha(\sigma)$$

*Proof.* Since we have assumed uniform substitution and by theorem 5.4 the result of substituting  $\sigma$  for  $p$  in  $\Box(p \leftrightarrow \sigma) \rightarrow (p \leftrightarrow \alpha)$  is a theorem of **GL**. This means that  $\vdash_{\mathbf{GL}} \Box(\sigma \leftrightarrow \sigma) \rightarrow (\sigma \leftrightarrow \alpha(\sigma))$ .  $\Box(\sigma \leftrightarrow \sigma)$  is obviously a theorem of **GL** so we get:

$$\vdash_{\mathbf{GL}} \sigma \leftrightarrow \alpha(\sigma)$$

⊢

We can also state (by theorem 5.4 and 5.3) the fixed point theorem in the following form:

{cor:Fixed}

**Corollary 5.2** For every modal formula  $\alpha$  modalized in  $p$ , there is a modal formula  $\sigma$  only containing propositional letters contained in  $\alpha$  not containing  $p$  such that:

$$\vdash_{\mathbf{GL}} \Box(p \leftrightarrow \alpha) \leftrightarrow \Box(p \leftrightarrow \sigma)$$

### 5.3. Calculation of Fixed Points

The algorithm in the proof of the fixed theorem can be used to calculate specific fixed points for some given formula, that fulfills the conditions of the theorem.

When we have shown Solovay's completeness theorems, the fixed point theorem will then make it possible for us to determine the fixed point of a wide range of sentences of arithmetic. This is important because this way of finding these points are often a lot easier than finding the points only with the help of the arithmetical methods.

Below are a few example of how we can use this algorithm.

**Example 5.3** Let  $\alpha(p) = \Box \neg p$ . We will calculate the fixed point for  $\alpha$ . Let  $\gamma_1(p) = p$  and  $\beta(q_1) = q_1$ . We then have that  $\alpha(p) = \beta(\Box \gamma_1(p))$  and  $\alpha_1(p) = \beta(\top) = \top$ . The fixed point of  $\sigma_1$  of  $\alpha_1$  is then  $\top$ , since  $p$  does not occur in  $\alpha_1$ . We can thus conclude that the fixed point  $\sigma$  of  $\alpha$  is the following:

$$\beta(\Box \gamma_1(\top)) = \beta(\Box \neg \top) = \Box \neg \top$$

this formula is equivalent to  $\Box \perp$ .

⊢

**Example 5.4** Let  $\alpha(p) = \Box p \rightarrow \Box \neg p$ . To find the fixed point, let  $\gamma_1(p) = p, \gamma_2(p) = \neg p$  and  $\beta(q_1, q_2) = q_1 \rightarrow q_2$ . Then clearly we have:

$$\alpha(p) = \beta(\Box \gamma_1(p), \Box \gamma_2(p))$$

$$\alpha_1(p) = \beta(\top, \Box \gamma_2(p)) = \top \rightarrow \Box \neg p$$

and

$$\alpha_2(p) = \beta(\Box \gamma_1(p), \top) = \Box p \rightarrow \top$$

It is clear that  $\alpha_1$  is equivalent to  $\Box \neg p$  and that  $\alpha_2$  is equivalent to  $\top$ ; thus they have fixed points  $\sigma_1 = \Box \perp$  and  $\sigma_2 = \top$ . So the fixed point  $\sigma$  of  $\alpha$  is the following:

$$\beta(\Box \gamma_1(\Box \perp), \Box \gamma_2(\top)) = \Box \Box \perp \rightarrow \Box \neg \top$$

or equivalent:  $\Box \Box \perp \rightarrow \Box \perp$ . ⊣

In figure 5.1 we can see some more fixed points. formula on the left is the formula  $\alpha(p)$  and the formula on the right is the formula  $\sigma$ . Line (3) and (6) has already been shown.

|   |  |
|---|--|
| 1. $\neg \Box p$  | $\neg \Box \perp$  |
| 2. $\Box p$   | $\top$   |
| 3. $\Box \neg p$  | $\Box \perp$   |
| 4. $\neg \Box \neg p$   | $\neg \Box \perp$  |
| 5. $\neg \Box \Box \neg p$  | $\neg \Box \Box \perp$   |
| 6. $\Box p \rightarrow \Box \neg p$   | $\Box \Box \perp \rightarrow \Box \perp$                                   |
| 7. $\Box(\neg p \rightarrow \Box \perp) \rightarrow \Box(p \rightarrow \Box \perp)$ | $\Box \Box \Box \perp \rightarrow \Box \Box \perp$                         |
| 8. $\Box p \rightarrow q$   | $\Box q \rightarrow q$   |
| 9. $\Box(p \rightarrow q)$  | $\Box q$   |
| 10. $\Box p \wedge q$   | $\Box q \wedge q$  |
| 11. $\Box(p \wedge q)$  | $\Box q \wedge q$  |
| 12. $q \vee \Box p$   | $\top$   |
| 13. $\neg \Box(q \rightarrow p)$  | $\Diamond q$   |
| 14. $\Box(p \rightarrow q) \rightarrow \Box \neg p$                                 | $\Box(\Box \perp \rightarrow q) \rightarrow \Box \perp$                    |
| 15. $q \wedge (\Box(p \rightarrow q) \rightarrow \Box \neg p)$                      | $q \wedge \Box \neg q$   |
| 16. $\Diamond p \rightarrow (q \wedge \neg \Box(p \rightarrow q))$                  | $\Diamond \top \rightarrow (q \wedge \neg \Box(\Box \perp \rightarrow q))$ |
| 17. $\Box(\Box(p \wedge q) \wedge \Box(p \wedge r))$                                | $\Box(\Box q \wedge \Box r)$   |
| 18. $\Box p \vee p$   | $\top$   |

## 5. Fixed Point Theorem

{Fig}

Figure 5.1.: A Table of Fixed Points

We will now show that  $\Box p \vee p$  has the fixed point from the table. For this end we will need the following theorem: We will start of by showing that no  $\beta$  modalized in  $p$  is equivalent to  $\Box p \vee p$ . This will the following two propositions show:

**Proposition 5.1** Suppose that  $\beta$  is modalized in  $p$ . Then we have:

$$\vdash_{\mathbf{GL}} p \rightarrow \beta \Rightarrow \vdash_{\mathbf{GL}} \beta$$

*Proof.* Suppose that  $\not\vdash_{\mathbf{GL}} \beta$ . This means that for some finite transitive and ir-reflexive model  $\mathcal{K} = \langle W, R, \phi \rangle$  and some  $w \in W$  that:  $\not\models_w \beta$ . Let  $\phi'$  be such that  $w \in \phi'(p)$  and otherwise it is just  $\phi$  and let  $\mathcal{K}' = \langle W, R, \phi' \rangle$ . We have that  $\beta$  is a truth-functional compound of formulae  $\Box \gamma$  and propositional letters that are not  $p$ . By continuity we have that  $\not\models_w^{\mathcal{K}'} \beta$  but we have that  $\models_w^{\mathcal{K}'} p$ . Hence we have that  $\models_w^{\mathcal{K}'} p \rightarrow \beta$  and the propositions follows.  $\dashv$

The next proposition shows that no  $\beta$  equivalent to  $\Box p \vee p$  is modalized in  $p$ .

**Proposition 5.2** For no  $\beta$  modalized in  $p$ , we have that:

$$\vdash_{\mathbf{GL}} \beta \leftrightarrow (\Box p \vee p)$$

*Proof.* If there was such a  $\beta$  we would have that  $\vdash_{\mathbf{GL}} p \rightarrow \beta$  and thus from the just shown proposition we would have that  $\vdash_{\mathbf{GL}} \beta$  and  $\vdash_{\mathbf{GL}} \Box p \vee p$ . By substitution we would then have that  $\vdash_{\mathbf{GL}} \Box \perp \vee \perp$  and this is clearly false.  $\dashv$

So  $\Box p \vee p$  is not equivalent to any formula modalized in  $p$ , but we can show that it has a fixed point any way:

**Proposition 5.3**

$$\vdash_{\mathbf{GL}} \Box(p \leftrightarrow \Box p \vee p) \leftrightarrow \Box(p \leftrightarrow \top)$$

*Proof.* The right to left implication is trivial. The other way will be shown by the following deduction:

$$\begin{aligned} \vdash_{\mathbf{GL}} \Box(p \leftrightarrow \Box p \vee p) &\rightarrow \Box(\Box p \rightarrow p) \\ &\rightarrow \Box \Box(\Box p \rightarrow p) \text{ By theorem 5.1} \\ &\rightarrow \Box(\Box(\Box p \rightarrow p) \wedge (\Box p \rightarrow p)) \text{ By definition of } \Box \\ &\rightarrow \Box(\Box p \wedge (\Box p \rightarrow p)) \\ &\rightarrow \Box p \\ &\rightarrow (p \leftrightarrow \top) \end{aligned}$$

$\dashv$

We will after the proof of Solovay's Completeness Theorems come back to this table, and make some conclusions about the fixed points.



## 6. Solovays Completeness Theorems

In this section we will prove Solovay's completeness theorems. The proof of these theorems follows a technique invented by Robert Solovay, which today is known as a *Solovay construction*. This technique is a way of embedding Kripke models into arithmetic. The first theorem show that the  $\Box$ -operator of the logic **GL** behaves like the proof predicate  $\text{Pr}$  from **PRA**. The second theorem shows that the modal logic **GLS**.

Further these theorems shows that the proof predicate of arithmetics can be axiomatized, by the axioms of **GL** and **GLS**.

### 6.1. Soundness

For each formula in  $\mathcal{L}_\Box$  we want to assign a sentence of  $\mathcal{L}_{\mathbf{PRA}}$ . This can be done in the following way.

**Definition 6.1** An interpretation of  $\mathcal{L}_\Box$  in **PRA** is a function that to each formula  $\alpha$  of  $\mathcal{L}_\Box$  assigns a sentence  $\alpha^*$  of **PRA** which satisfies the following requirements:

1. For atomic  $p$ ,  $p^*$  is a formula of the language of arithmetic
2.  $(\perp)^* = "0 = 1"$
3.  $(\alpha \rightarrow \beta)^* = "\alpha^* \rightarrow \beta^*"$
4.  $(\Box\alpha)^* = "\text{Pr}(\ulcorner \alpha^* \urcorner)"$

We will further say that a modal formula  $\alpha$  is **PRA**-valid if, in every interpretation  $^*$ ,  $\alpha^*$  is a theorem of **PRA**. —

The goal of the current section is to prove the that the set of **PRA**-valid formulas are the theorems of **GL**. I.e we want to prove the following bi-implication:

$$\mathbf{GL} \vdash \alpha \Leftrightarrow \forall^* (\mathbf{PRA} \vdash \alpha^*)$$

This formula says that **GL** completely captures what **PRA** can say about its own provability. This result can be expanded to other fragments of arithmetics. So the modal logic **GL** is the modal logic that captures what a lot of different fragments

## 6. Solovays Completeness Theorems

---

of arithmetics can say about its own provability. Later on in the project it will be shown for which fragments this is the case.

In 6.3 we will show that **GLS** is the modal logic of provability in truth. This is called Solovay's second completeness theorem.

We will start with proving the " $\Rightarrow$ " implication; i.e that every theorem of **GL** is **PRA**-valid. The other way will be a bit harder to prove, and that is the part that is known as Solovay's first completeness theorem.

**Theorem 6.1 (Soundness)** For all modal sentences  $\varphi$  we have that :

$$\mathbf{GL} \vdash \varphi \Rightarrow \forall^*(\mathbf{PRA} \vdash \varphi^*)$$

*Proof.* The proof will be done as an induction proof on the number of axioms and rules of inference use in a **GL**-proof of a formula  $\alpha$ . The proof will be done by looking at the last rule or axiom schema used in the proof of  $\alpha$ .

The case of A1 and R1 are clear. We further have that the theorems of **PRA** are closed under modus ponens so A2 is also clear.

Assume that the last step of the proof of  $\alpha$  is an instance of R2. We have that  $\text{Pr}(x)$  is  $\Sigma_1$  formula, so it is equivalent to some formula of the form  $\exists y R(x, y)$ , where  $R$  is a primitive recursive predicate. We know that if a  $\Sigma_1$  sentences is true it is provably, so this shows the case of R2.

The axiom A3 can be derived from the others, so this case is redundant.

We will now just have to show the case of A4. This case follows by Löb's theorem.  $\dashv$

## 6.2. The First Theorem

Having shown soundness, we will now show the completeness theorem:

**Theorem 6.2 (Solovay's first completeness theorem)** For all modal sentences  $\varphi$  we have that:

$$\forall^*(\mathbf{PRA} \vdash \alpha^*) \Rightarrow \mathbf{GL} \vdash \alpha$$

This theorem will be shown by contraposition. So we want to show the following: "If  $\mathbf{GL} \not\vdash \alpha$  then we have one  $*$  such that  $\mathbf{PRA} \not\vdash \alpha^*$ ". The start of the proof is the following: If  $\mathbf{GL} \not\vdash \alpha$  then  $\mathbf{GL} \not\vdash \alpha$  and thus there is a finite pointed Kripke model  $\mathcal{K} = \langle W, R, \phi, w_0 \rangle$  in which we have  $w_0 \notin \phi(\alpha)$ . The goal is then to find an interpretation  $*$  such that  $\mathbf{PRA} \not\vdash \alpha^*$ .

The construction of this  $*$  is rather complex, and will take the rest of this subsection to prove. So for the rest of this subsection fix a sentence  $\varphi$  such that  $\mathbf{GL} \not\vdash \varphi$  and let  $\mathcal{K} = \langle W, R, \phi, w_0 \rangle$  be a pointed Kripke model such that  $w_0 \in \phi(\alpha)$ .



We will assume without loss of generality that  $W = \{0, \dots, n\}$  for some finite  $n$  and that  $w_0 = 1$ .  $R$  can be extended by setting  $0Ri$  for each  $i$  in  $W$ . It shall be noted that 0 is not part of our Kripke model, but we will in section refchap:second create a model where it is a part of the model.

Further we will first intuitively define a function  $\varphi : \omega \rightarrow \{0, \dots, n\}$  in the following way: Set  $\varphi(0) = 0$ . Further we define  $\varphi(x+1)$  in the following way: If  $x+1$  is the code of a proof that  $\lim_{k \rightarrow \infty}(k) \neq z$  for some  $z$  accessible to  $\varphi(x)$  we set  $\varphi(x+1) = z$  otherwise we have that  $\varphi(x+1) = \varphi(x)$ .

The way this function works can be explained intuitively by the following quote:

Imagine a refugee who is admitted from one country to another only if he/she provides a proof not to stay there forever. If the refugee is also never allowed to go to one of the previously visited countries, he/she must eventually stop somewhere. So, an honest refugee will never be able to leave his/her country of origin. (S. N. Artemov and Beklemishev n.d.)

Before we can give a formal definition of the function  $\varphi$ , we will have to introduce some notation. First of we will let  $\psi$  be an arbitrary partial recursive function with the following  $\Sigma_1$  graph:  $\tau v_0 v_1$ . From this graph we can obtain the following  $\Sigma_2$  formula:

$$\exists v_0 \forall v_1 > v_0 : \tau v_1 v$$

Which says that  $\psi$  has limit  $v$ . We will abbreviate this as  $L_\psi = v$ . We will use this notation to define  $\varphi$  in the following way:

$$\varphi(v_0) = v_1 \leftrightarrow \begin{cases} (v_0 = \bar{0} \wedge v_1 = \bar{0}) \vee \\ (v_0 > \bar{0} \wedge \text{Prov}(v_0, \ulcorner L \neq v_1 \urcorner) \wedge \varphi(v_0 - \bar{1}) \bar{R} v_2) \vee \\ (v_0 > \bar{0} \wedge \forall v_2 \leq v_0 \neg (\text{Prov}(v_0, \ulcorner L \neq v_2 \urcorner) \wedge \bar{\varphi}(v_0 - \bar{1}) \bar{R} v_2) \wedge \\ v_1 = \bar{\varphi}(v_0 - \bar{1})) \end{cases}$$

We will define the graph  $\tau v_0 v_1 = \exists v_3 \chi v_3 v_0 v_1$  of the partial recursive function  $\varphi$ . Since we have that the graph of  $\varphi$  is  $\Sigma_1$  we have that the graph  $\tau$  is also  $\Sigma_1$  and thus the graph  $\chi$  is  $\Delta_0$ . We will define a formula  $\Phi(\chi)$  as the disjunction of the following three formulas:

1.  $v_0 = \bar{0} \wedge v_1 = \bar{0}$
2.  $v_0 > \bar{0} \wedge \text{Prov}(v_0, \ulcorner L_\tau \neq v_1 \urcorner) \wedge \exists v_4 (\tau(v_0 - \bar{1}, v_4) \wedge v_4 \bar{R} v_1)$
3.  $v_0 > \bar{0} \wedge \exists v_3 v_4 \forall v_2 \leq v_0 \neq (\text{Prov}(v_0, \ulcorner L_\tau \neq v_2 \urcorner) \wedge \chi(v_3, v_0 - \bar{1}, v_4) \wedge v_4 \bar{R} v_1) \wedge \tau(v_0 - \bar{1}, v_1)$

## 6. Solovays Completeness Theorems

---

Thus we have that the set  $\Phi(\chi)$  can be seen as a primitive recursive function of  $\ulcorner \chi \urcorner$ ; i.e we have:

$$\ulcorner \Phi(\chi) \urcorner = \vartheta(\ulcorner \chi \urcorner)$$

Since primitive functions are recursive we have by the Recursion Theorem that we can pick an  $n$  such that  $\varphi_{\vartheta(n)} = \varphi_n$ . Now set  $\varphi = \varphi_n$ . This definition of  $\varphi$  is rather involved. But we have defined  $\varphi$  by its own graph, and avoided the imperant circularity by using the recursion theorem to not rely on the graph.

We further define the relation  $\bar{R}$  by listing all pairs  $(x, y) \in R$  (We can do this since the set  $R$  is finite) and then define:

$$v_0 \bar{R} v_1 : \bigvee_{(x,y) \in R} (v_0 = \bar{x} \wedge v_1 = \bar{y})$$

Thus we have that  $\varphi$  have the following properties:

1.  $\varphi(0) = 0$
2. If  $x + 1$  proves that  $L \neq \bar{z}$  and we have that  $\varphi(x) R z$ , then  $\varphi(x + 1) = z$
3. Else we have that  $\varphi(x + 1) = \varphi(x)$

Further it is a total function since it is defined by recursion, and this can be proven in **PRA**.

**Proposition 6.1** Let  $\psi$  be the  $\Sigma_1$ -formula that defines the graph of  $\varphi$ . Then:

$$\mathbf{PRA} \vdash \forall v_0 \exists! v_1 \psi v_0 v_1$$

*Proof.* The uniqueness of  $v_1$  is given by the definition of  $\varphi$ . The existence of a value is given by induction on  $v_0$  in the  $\Sigma_1$  formula  $\exists v_1 \psi v_0 v_1$  in **PRA**, and thus this induction is possible in **PRA**.

**Base step:** When  $v_0 = 0$  we have that  $\psi(0) = 0$  by definition of  $\varphi$ ; i.e  $\exists v_1 \psi v_0 v_1$ , where  $v_1 = 0$ .

**Induction step:** Assume that  $\exists v_1 \psi v_0 v_1$  holds for  $v_0 = n$ , i.e we have a  $v_1 = m$  such that  $\varphi(n) = m$ . Look at  $\varphi(n + 1)$ , then if  $n + 1$  proves that  $L_\psi \neq \bar{z}$  and we have that  $\varphi(n) R z$ , then we will have that  $\varphi(n + 1) = z$ , i.e  $v_1 = z$  otherwise we will have that  $\varphi(n + 1) = \varphi(n) = m$ , and the induction is done.  $\dashv$

Further we will expand the language with a new function constant (This can be done, since  $\psi$  is the graph of a total function)  $\bar{\varphi}$  with the following defining axiom:

$$\bar{\varphi}(v_0) = v_1 \leftrightarrow \psi(v_0) = v_1$$

We will now prove and state a few lemmas about the function  $\varphi$  and the limit of this function  $L\tau = L$ . These will build up the proof of Solovay's First Completeness Theorem. Thus we will use the function  $\varphi$  and the limit  $L$  to deduce the theorem

{lem:2}

**Lemma 6.1** The following three statements holds:

$$1. \mathbf{PRA} \vdash \forall v_0 (\bar{\varphi}v_0 \leq \bar{n})$$

2. For all  $x \in \omega$  we have that:

$$\mathbf{PRA} \vdash \forall v_0 (\bar{\varphi}v_0 = \bar{x} \rightarrow \forall v_1 > v_0 (\bar{\varphi}v_1 = \bar{x} \vee \bar{x}\bar{R}\bar{\varphi}v_1))$$

$$3. \mathbf{PRA} \vdash \exists v_0 v_1 \forall v_2 > v_0 (\bar{\varphi}v_2 = v_1).$$

Where (3) just means that  $\mathbf{PRA} \vdash \exists v_1 (L = v_1)$ .

*Proof.* We will prove each part separately

1. We will prove this part by induction.

**Base case:**  $\varphi(0) = 0 \leq n$  is clearly true.

**Induction step:** Assume that  $\varphi(x) \leq n$  is true. Then we have that  $\varphi(x+1)$  is in the range of  $\varphi$  and this  $\leq n$  or we have that  $\varphi(x+1) \leq n$  so the induction is complete.

2. We will start of by write the formula we have to prove as the following equivalent formula:

$$\forall v_1 \forall v_0 (\bar{\varphi}v_0 = \bar{x} \rightarrow \varphi(v_0 + v_1 + 1) = \bar{x} \vee \bar{x}\bar{R}\bar{\varphi}(v_0 + v_1 + 1))$$

To prove this we will use induction on  $v_1$ . This is an induction on a  $\Pi_1$  formula; which is a possible induction for  $\mathbf{PRA}$  by [Ref]

**Base case:** if  $v_1 = 0$  then clearly we have that  $\varphi(v_0 + 1) = \bar{x}$  or  $\bar{x}\bar{R}\bar{\varphi}(v_0 + 1)$  by the definition of  $\varphi$ .

**Induction step:** Assume that it holds for  $v_1 = n - 1$ , i.e that we for all  $v_0$  have that:

$$\varphi(v_0 + \bar{n}) = \bar{x} \text{ or } \bar{x}\bar{R}\bar{\varphi}(v_0 + \bar{n})$$

We will split the rest of this proof up in two parts; one where the first disjunct is true, and one where the second is true.

**The first disjunct is true:.** Assume that both  $v_0 + n + 1$  codes a proof that  $L \neq z$  and  $\varphi(v_0 + \bar{n})Rz$  is true. This means that  $\varphi(v_0 + \bar{n} + 1) = z$ . But this is just that  $\bar{x}\bar{R}\bar{\varphi}(v_0 + \bar{n} + 1)$ , and the lemma is true in this case. If these

## 6. Solovays Completeness Theorems

assumptions are not true, we will then have that  $\varphi(v_0 + \bar{n} + 1) = \varphi(v_0 + \bar{n})$ . But then we have that  $\varphi(v_0 + \bar{n} + 1) = \bar{x}$ , and the lemma holds.

**The second disjunct is true:** We will again start of by assuming that  $L \neq z$  and  $\varphi(v_0 + n)Rz$ ; i.e  $\varphi(v_0 + n + 1) = z$ . We then get that  $\varphi(v_0 + n)R\varphi(v_0 + n + 1)$  and by the transitivity of  $R$  we get:  $x\bar{R}\varphi(v_0 + n + 1)$ . If we otherwise have that  $\varphi(v_0 + n) = \varphi(v_0 + n + 1)$  we clearly have that  $\bar{x}\bar{R}\varphi(v_0 + n + 1)$ .

3. Here we will first prove the following:

$$\forall v_0(\exists v_1(\bar{F}v_1 = v_0) \rightarrow \exists v_1(L = v_1))$$

This is clearly true for  $v_0 > n$ . For  $v_0 \leq n$  we will use induction on the converse of  $R$ . By 2) from this lemma it holds for maximal nodes  $y$  in  $W$ . If  $y$  is not a maximal node and  $\exists v_1(\bar{\varphi}(v_1) = \bar{x})$  then again by 2) we either get that  $L = \bar{x}$  or that for some  $y$  such that  $xRy$ :  $\exists v_1(\bar{\varphi}(v_1) = \bar{y})$ , and the induction hypothesis gives us that  $\exists v_1(L = v_1)$ . But since the set  $W$  is finite we will end up with getting:

$$\mathbf{PRA} \vdash \exists v_1(\bar{\varphi}(v_1) = \bar{0}) \rightarrow \exists v_1(L = v_1)$$

But we have that  $\mathbf{PRA} \vdash \bar{\varphi}(\bar{0}) = \bar{0}$  so clearly:  $\mathbf{PRA} \vdash \exists v_1(L = v_1)$

⊢

The induction in the above proof was not done inside of **PRA**, both was done as metamathematical induction outside of **PRA**. This done because the formula  $\exists v_1(L = v_1)$  is  $\Sigma_2$  and thus we can not use induction on this formula inside of **PRA**.

**Corollary 6.1**  $\mathbf{PRA} \vdash L \leq \bar{n}$ , i.e  $\mathbf{PRA} \vdash \bigvee_{x \leq n} L = \bar{x}$

*Proof.* By lemma 6.1 1) and 3) and the following implication:

$$\mathbf{PRA} \vdash v \leq \bar{n} \rightarrow \bigvee_{x \leq n} v = \bar{x}$$

The corollary follows.

⊢

{lem:4}

**Lemma 6.2** For all  $x, y \leq n$  we have that:

1.  $L = \bar{x} \wedge \bar{x}\bar{R}\bar{y} \rightarrow \text{Con}_{\mathbf{PRA}+L=\bar{y}}$
2.  $\mathbf{PRA} \vdash L = \bar{x} \wedge \bar{x} \neq \bar{y} \wedge \neg(\bar{x}\bar{R}\bar{y}) \rightarrow \neg\text{Con}_{\mathbf{PRA}+L=\bar{y}}$

3.  $\mathbf{PRA} \vdash L = \bar{x} \wedge \bar{x} > \bar{0} \rightarrow \text{Pr}(\ulcorner L \neq \bar{x} \urcorner)$

*Proof.* We will prove each statement separately:

1) Let  $xRy$  and assume for contradiction that  $L = \bar{x} \wedge \text{Pr}(\ulcorner L \neq \bar{y} \urcorner)$ . Since we have that  $L = \bar{x}$  we can chose a  $v_0$  such that  $\forall v_2 (v_2 > v_0 \rightarrow \bar{F}v_2 = \bar{x})$  and we can also chose  $v_1 + \bar{1} > v_0$  such that  $\text{Prov}(v_1 + \bar{1}, \ulcorner L \neq \bar{y} \urcorner)$  But we also have the following:

$$\mathbf{PRA} \vdash \text{Prov}(v_1 + \bar{1}, \ulcorner L \neq \bar{y} \urcorner) \wedge \bar{F}v_1 = \overline{x_1, \dots, x_n} \wedge \bar{x}\bar{R}\bar{y} \rightarrow \bar{F}(v_1 + \bar{1}) = \bar{y}$$

This contradicts with  $\forall v_2 > v_0 (\bar{F}v_2 = \bar{x})$  which came from the assumption that  $L = \bar{x}$  so we can conclude:

$$\mathbf{PRA} \vdash L = \bar{x} \wedge \bar{x}\bar{R}\bar{y} \rightarrow \neg \text{Pr}(\ulcorner L \neq \bar{y} \urcorner)$$

Where we have that  $\neg \text{Pr}(L \neq \bar{y})$  is equivalent to  $\text{Con}_{\mathbf{PRA}+L=\bar{y}}$  and 1) follows.

2) By [ref??] We have the following deduction:

$$\mathbf{PRA} \vdash L = \bar{x} \rightarrow \exists v_0 (\bar{F}v_0 = \bar{x}) \quad (6.1)$$

$$\rightarrow \text{Pr}(\ulcorner \exists v_0 (\bar{F}v_0 = \bar{x}) \urcorner) \quad (6.2) \quad \{\text{eq:21}\}$$

Further we have from lemma 6.1.2 and [D2, lob?] that:

$$\mathbf{PRA} \vdash \forall v_0 (\bar{F}v_0 = \bar{x} \rightarrow (L = \bar{x} \vee \bar{x}\bar{R}L)) \quad (6.3)$$

$$\mathbf{PRA} \vdash (\ulcorner \forall v_0 (\bar{F}v_0 = \bar{x} \rightarrow (L = \bar{x} \vee \bar{x}\bar{R}L)) \urcorner) \quad (6.4) \quad \{\text{eq:22}\}$$

From 6.2 and 6.4 we have the following:

$$\mathbf{PRA} \vdash L = \bar{x} \rightarrow \text{Pr}(\ulcorner L = \bar{x} \vee \bar{x}\bar{R}L \urcorner) \quad (6.5) \quad \{\text{eq:23}\}$$

asd?

Which gives us [Why?]

$$\mathbf{PRA} \vdash \bar{x} \neq \bar{y} \wedge \neg(\bar{x}\bar{R}\bar{y}) \rightarrow \text{Pr}(\ulcorner \bar{x} \neq \bar{y} \wedge \neg(\bar{x}\bar{R}\bar{y}) \urcorner)$$

This with 6.5 gives us:

$$\mathbf{PRA} \vdash L = \bar{x} \wedge \bar{x} \neq \bar{y} \wedge \neg \bar{x}\bar{R}\bar{y} \rightarrow \text{Pr}(\ulcorner L = \bar{x} \vee \bar{x}\bar{R}L \urcorner) \wedge \text{Pr}(\ulcorner \bar{x} \neq \bar{y} \wedge \neg \bar{x}\bar{R}\bar{y} \urcorner)$$

From which we can deduce:

$$\mathbf{PRA} \vdash L\bar{x} \wedge \bar{x} \neq \bar{y} \wedge \neg \bar{x}\bar{R}\bar{y} \rightarrow \text{Pr}(\ulcorner L \neq \bar{y} \urcorner)$$

## 6. Solovays Completeness Theorems

Which gives us 2)

3) From the least number principle we have that:

$$\mathbf{PRA} \vdash L = \bar{x} \wedge \bar{x} > \bar{0} \rightarrow \exists v (\bar{F}(v + \bar{1}) = \bar{x} \wedge \bar{F}v \neq \bar{x})$$

By the definition of  $F$  we have for such a  $v$  the following:

$$\mathbf{PRA} \vdash \bar{F}(v + \bar{1}) = \bar{x} \bar{F}v \neq \bar{x} \rightarrow \text{Prov}(v + \bar{1}, \ulcorner L \neq \bar{x} \urcorner)$$

And thus we have the following:

$$\mathbf{PRA} \vdash L = \bar{x} \wedge \bar{x} > \bar{0} \rightarrow \text{Pr}(\ulcorner L \neq \bar{x} \urcorner)$$

⊥

Lemma 6.1 and 6.2 gives us the moss of the facts that we need about  $L$  and  $\varphi$ ; at least the facts about them that we can prove in **PRA**. We will also need the following result, which cannot be proven in **PRA**.

{lem:5}

**Lemma 6.3** The following two statements are true, but they cannot be proven in **PRA**.

1.  $L = \bar{0}$
2. For  $0 \leq x \leq n$  we have that  $\mathbf{PRA} + L = \bar{x}$  is consistent.

*Proof.* 1) By lemma 6.1.3 the limit  $L$  exists. If  $x > 0$  we have by lemma 6.2 that:

$$\begin{aligned} L = \bar{x} &\Rightarrow \mathbf{PRA} \vdash L \neq \bar{x} \\ &\Rightarrow L \neq \bar{x} \end{aligned}$$

Since **PRA** is sound. But this is a contradiction and we must conclude that  $L = 0$ .

2) Since we have that  $L = \bar{0}$  is true and we have that **PRA** is sound, we have that  $\mathbf{PRA} + L = \bar{0}$  is consistent. For  $x > 0$  we will apply lemma 6.2.1 and get:

$$\mathbf{PRA} \vdash L = \bar{0} \wedge \bar{0} \bar{R} \bar{x} \rightarrow \text{Con}_{\mathbf{PRA} + L = \bar{x}}$$

We have that the antecedent is true and hence that  $\text{Con}_{\mathbf{PRA} + L = \bar{x}}$  is true, which proves this part of the lemma. ⊥

We have now shown all the important basic properties that  $\varphi$  and  $L$  holds. In the next part of the proof we will simulate the Kripke model  $\mathcal{K} = \langle W, R, \phi \rangle$ , where  $1 \notin \phi(\alpha)$ . For this end we will let  $L = \bar{x}$ , for  $x > 0$ , assume the rule nodes of

$W = \{1, \dots, n\}$ . We will start of by defining the interpretation  $*$ . So for any  $p$  let:

$$p^* = \bigvee \{L = \bar{x} : 1 \leq x \leq n \text{ and } x \in \phi(p)\}$$

If this disjunction is empty, we will set it to be  $\bar{0} = \bar{1}$ . Further we are only interested the sentence  $\alpha$ ; i.e the set:

$$S(\alpha) = \{\beta : \beta \text{ is a subformula of } \alpha\}$$

We will not look at  $p \notin S(\alpha)$

The following lemma is the crucial result about this interpretation, and the theorem will follow easily from this result:

{lem:10}

**Lemma 6.4** Let  $1 \leq x \leq n$ . For any  $\beta$  and  $*$  as defined just above we have that:

1.  $x \in \phi(\beta) \Rightarrow \mathbf{PRA} \vdash L = \bar{x} \rightarrow \beta^*$
2.  $x \notin \phi(\beta) \Rightarrow \mathbf{PRA} \vdash L = \bar{x} \rightarrow \neg \beta^*$

*Proof.* We will use induction on complexity of  $\psi$ . If  $\beta = p$ , then since  $L = \bar{x}$  is a disjunct of  $p^*$  we get:

$$x \in \phi(\varphi) \Rightarrow \mathbf{PRA} \vdash L = \bar{x} \rightarrow p^*$$

Which proves 1, for  $\psi$  atomic. For 2 observe that if  $x \notin \phi(p)$ , then  $L = \bar{x}$  contradicts all the disjuncts of  $p^*$  and thus:

$$x \notin \phi(p) \rightarrow \mathbf{PRA} \vdash L = \bar{x} \rightarrow \neg p^*$$

The cases where  $\psi$  is  $\neg\gamma, \gamma \wedge \sigma, \gamma \vee \sigma$  and  $\gamma \rightarrow \sigma$  are trivial. So we will just look at the case where  $\psi = \Box\gamma$ . Here we make the following deductions:

$$\begin{aligned} x \in \phi(\Box\theta) &\Rightarrow \forall y (xRy \Rightarrow y \in \phi(\gamma)) \\ &\Rightarrow \forall (xR \Rightarrow \mathbf{PRA} \vdash L = \bar{y} \rightarrow \gamma^*) \\ &\Rightarrow \bigwedge_{xRy} (\mathbf{PRA} \vdash L = \bar{y} \rightarrow \gamma^*) \\ &\Rightarrow \mathbf{PRA} \vdash \bigvee_{xRy} L = \bar{y} \rightarrow \gamma^* \\ &\Rightarrow \mathbf{PRA} \vdash \text{Pr}(\bigvee_{xRy} L = \bar{y} \rightarrow \gamma^*) \end{aligned}$$

And by the last line we can get the following by using the axioms of **GL**

$$x \in \phi(\Box\gamma) \Rightarrow \mathbf{PRA} \vdash \text{Pr}(\bigvee_{xRy} L = \bar{y} \rightarrow \gamma^*) \quad (6.6) \quad \{\text{eq:1}\}$$

## 6. Solovays Completeness Theorems

---

We can now invoke lemma 6.3 2 and 3 and get:

$$\mathbf{PRA} \vdash L = \bar{x} \rightarrow \bigwedge_{\neg(xRz)} \text{Pr}(\ulcorner L \neq z \urcorner) \quad (6.7) \quad \{\text{eq:2}\}$$

and therefore we have that:

$$\{\text{eq:3}\} \quad \mathbf{PRA} \vdash L = \bar{x} \rightarrow \text{Pr}(\ulcorner \bigvee_{xRy} L = \bar{y} \urcorner) \quad (6.8)$$

Now by 6.6 and 6.8 we get that:

$$\begin{aligned} \vdash_x \Box\theta &\Rightarrow \mathbf{PRA} \vdash L = \bar{x} \rightarrow \text{Pr}(\ulcorner \gamma^* \urcorner) \\ &\Rightarrow \mathbf{PRA} \vdash L = \bar{x} \rightarrow (\Box\gamma)^* \end{aligned}$$

I.e we have proven part 1 of the lemma. Similarly we can do the following deduction:

$$\begin{aligned} x \notin \phi(\Box\gamma) &\Rightarrow \exists y(xRy \wedge y \notin \phi(\gamma)) \\ &\Rightarrow \exists y(xRy \wedge \mathbf{PRA} \vdash L = \bar{y} \rightarrow \neg\gamma^*) \\ &\Rightarrow \exists y(xRy \wedge \mathbf{PRA} \vdash \gamma^* \rightarrow L \neq \bar{y}) \\ &\Rightarrow \exists y(xRy \wedge \mathbf{PRA} \vdash \text{Pr}(\ulcorner \gamma^* \urcorner) \rightarrow \text{Pr}(\ulcorner L \neq \bar{y} \urcorner)) \end{aligned}$$

But by lemma 6.3 we get that if  $xRy$ :

$$\mathbf{PRA} \vdash L = \bar{x} \rightarrow \neg\text{Pr}(\ulcorner L \neq \bar{y} \urcorner)$$

all in all this gives us:

$$\mathbf{PRA} \vdash L = \bar{x} \rightarrow \neg\text{Pr}(\ulcorner \gamma^* \urcorner)$$

Which is just the following we where trying to show:

$$\mathbf{PRA} \vdash L = \bar{x} \rightarrow \neg(\Box\gamma)^*$$

+

We can now finally prove Solovay's first completeness theorem:

*Proof of Solovay's first completeness theorem.* By lemma 6.4 we have that

$$1 \notin \phi(\alpha) \Rightarrow \mathbf{PRA} \vdash L = \bar{1} \rightarrow \neg\alpha^*$$

But by lemma 6.3 we have that  $\mathbf{PRA} \vdash L = 1$  is consistent, from which it follows



that  $\mathbf{PRA} + \neg\varphi^*$  is consistent; so  $\varphi^*$  is not a theorem of  $\mathbf{PRA}$  and thus we have:  $\mathbf{PRA} \not\vdash \varphi^*$  and the theorem follows by contraposition.  $\dashv$

With this proof we can give a interpretation of the set  $W$  and relation  $R$  in a model of  $\mathbf{GL}$ . The set  $W$  consists of recursively axiomatized extensions of  $\mathbf{PRA}$ . And we have that  $w_1 R w_2$  if and only if  $w_1 \vdash \text{Con}(w_2)$ , i.e the theory  $w_1$  has the consistency of theory  $w_2$  as one of its theorems. It is clear that this relation is transitive and conversely well-founded.

## 6.3. The Second Theorem

{chap:second}

Solovay's Second Completeness Theorem is a strengthening of the first one, since it tells about trueness of a formulae  $\alpha^*$  in the standard model  $\mathcal{N} = \langle \omega, +, \cdot \rangle$  of arithmetics. The proof will follow that of the first theorem, but we will look at the modal logic  $\mathbf{GLS}$  instead of the modal logic  $\mathbf{GL}$ .

**Theorem 6.3** For all modal sentences  $\varphi$ , the following is equivalent:

1.  $\mathbf{GLS} \vdash \alpha$
2.  $\mathbf{GL} \vdash \bigwedge_{\square\beta \in S(\alpha)} (\square\beta \rightarrow \beta) \rightarrow \alpha$
3.  $\alpha$  is true in all  $\alpha$ -reflexive FT Kripke models
4.  $\forall^*(\alpha^* \text{ is true})$

Some parts of this theorem has already been proven. We have  $(1) \Leftrightarrow (2) \Leftrightarrow (3)$  by theorem 4.4. The implication  $(1) \Rightarrow (4)$  is clear, so we just have to prove  $(4) \Rightarrow (3)$

For proving  $(4) \Rightarrow (3)$  we will again make use of contraposition. Let  $\mathcal{K} = \langle (1, \dots, n), R, \phi \rangle$  be given and let 1 be the root. Further let  $\alpha$  be such that  $1 \notin \phi(\alpha)$ . We will assume that  $\mathcal{K}$  is  $\alpha$ -reflexive. This means that we have:  $1 \in \phi(\square\beta \rightarrow \beta)$  for all  $\beta \in S(\alpha)$

We will set  $0Rw$  for all  $w \in W$ . We will now create a new model  $\mathcal{K}'$  where we have added 0, so in this proof it is part of the Kripke model we will look at. We will create this new model in the following way:

$$W' = \{0, 1, \dots, n\}$$

$$R' \text{ extends } R \text{ by assuming that } 0R'x \text{ for all } x \in W$$

$$\alpha_0 = 0$$

$$\phi' \text{ extends } \phi \text{ by putting } 0 \in \phi'(p) \text{ iff } 1 \in \phi(p) \text{ for all } p \in S(\varphi)$$

## 6. Solovays Completeness Theorems

---

We will abuse notation and let  $R$  denote  $R'$  and  $\phi$  denote  $\phi'$ .

**Lemma 6.5** For all  $\beta \in S(\alpha)$  we have that:

$$0 \in \phi(\beta) \text{ iff } 1 \in \phi(\beta)$$

*Proof.* Kig på intro af GLS. ⊥

We will now define a function *varphi* in the same way as before.

$$\begin{aligned} \varphi(0) &= 0 \\ \varphi(x+1) &= \begin{cases} y & \text{Prov}(\bar{x} + \bar{1}, \ulcorner L \neq \bar{y} \urcorner) \wedge xRy \\ \varphi(x) & \text{else} \end{cases} \end{aligned}$$

All the Lemmas about  $\varphi$  and  $L$  still holds, since the function  $\varphi$  is only determined by the frame  $\langle W, R \rangle$  and not the Kripke model that we are looking at. But the behavior of  $\phi$  has changed, since we must added the node 0 and we can only use sub formulas of  $\alpha$ . So we define:

$$p^* = \bigvee \{L = \bar{x} : 0 \leq x \leq n \wedge x \in \phi(p)\}$$

For  $p \in S(\alpha)$  and let  $p^*$  be random for all  $p$ 's that is not a subformula of  $\alpha$ . We now prove and state the following lemma is analogies to lemma 6.3:

**Lemma 6.6** Let  $0 \leq x \leq n$ . For any  $\beta \in S(\alpha)$  and  $*$  as defined above we have:

1.  $x \in \phi(\beta) \Rightarrow \mathbf{PRA} \vdash L = \bar{x} \rightarrow \beta^*$
2.  $x \notin \phi(\beta) \Rightarrow \mathbf{PRA} \vdash L = \bar{x} \rightarrow \neg\beta^*$

*Proof.* For  $0 < x$  the proof is identical to the proof of [ref?]. So we will only prove the case where  $x = 0$ . The is again an induction on the complexity of  $\beta$ . We will only prove the cases where  $\beta = \Box\gamma$ .

Let  $\beta = \Box\gamma$ . We then have:

$$\begin{aligned} 0 \in \phi(\Box\gamma) &\Rightarrow \forall x(1 \leq x \leq n \Rightarrow x \in \phi(\gamma)) \\ &\Rightarrow \forall x(1 \leq x \leq n \Rightarrow \mathbf{PRA} \vdash L = \bar{x} \rightarrow \gamma^*) \end{aligned}$$

Since  $x > 1$  and this case of the lemma has been proven, when we proved the lemma for the first completeness theorem. We can also make the following deduction by

the induction hypothesis:

$$\begin{aligned} 0 \in \phi(\Box\gamma) &\Rightarrow 1 \in (\gamma) \\ &\Rightarrow 0 \in \gamma) \\ &\Rightarrow \mathbf{PRA} \vdash L = \bar{0} \rightarrow \gamma^* \end{aligned}$$

By combining these two we get:

$$\begin{aligned} 0 \in \phi(\Box\gamma) &\Rightarrow \bigwedge_{x \leq n} (\mathbf{PRA} \vdash L = \bar{x} \rightarrow \gamma^*) \\ &\Rightarrow \mathbf{PRA} \vdash (\bigvee_{x \leq n} L = \bar{x}) \rightarrow \gamma^* \\ &= \mathbf{PRA} \vdash \Pr(\ulcorner \bigvee L = \bar{x} \urcorner) \rightarrow \Pr(\ulcorner \gamma^* \urcorner) \end{aligned}$$

But by corollary [ref?] we have that  $\mathbf{PRA} \vdash \bigvee L = \bar{x}$  and thus  $\mathbf{PRA} \vdash \Pr(\ulcorner \bigvee L = \bar{x} \urcorner)$  so all in all we have:

$$\begin{aligned} 0 \in (\Box\gamma) &\Rightarrow \mathbf{PRA} \vdash \Pr(\ulcorner \gamma^* \urcorner) \\ &\Rightarrow \mathbf{PRA} \vdash L = \bar{0} \rightarrow \Pr(\ulcorner \gamma^* \urcorner) \end{aligned}$$

This proves (1). The proof of (2) is a bit easier. We have:

$$\begin{aligned} 0 \notin \phi(\Box\gamma) &\Rightarrow \exists x(1 \leq x \leq n \wedge x \notin \phi(\gamma)) \\ &\Rightarrow \exists x(1 \leq x \leq n \wedge \mathbf{PRA} \vdash L = \bar{x} \rightarrow \neg\gamma^*) \\ &\Rightarrow \exists x(1 \leq x \leq n \wedge \mathbf{PRA} \vdash \gamma^* \rightarrow L \neq \bar{x}) \\ &\Rightarrow \mathbf{PRA} \vdash L = \bar{0} \rightarrow \neg\Pr(\ulcorner \gamma^* \urcorner) \end{aligned}$$

And thus by Lemma [Ref?] we have that  $\mathbf{PRA} \vdash L = \bar{0} \rightarrow \neg\Pr(\ulcorner L \neq \bar{x} \urcorner)$  for  $x > 0$   $\dashv$

We thus have that  $L = \bar{0}$  is true, and we can now finally prove the second completeness theorem:

*Proof of the Second Completeness theorem.* Assume that  $\alpha$  is false in  $\mathcal{K}$  i.e  $1 \notin \phi(\alpha)$ . Then by lemma [asd] we get  $0 \notin \phi(\alpha)$  and the just proven lemma gives:

$$\mathbf{PRA} \vdash L = \bar{0} \rightarrow \neg\alpha^*$$

Since we have that  $L = \bar{0}$  is true we then get  $\neg\alpha^*$  which just means that  $\alpha^*$  is false.  $\dashv$

### 6.3.1. Using the Second Completeness Theorem

We can use the second completeness theorem to prove the following theorem named after Rosser:


**Theorem 6.4** There is a arithmetical sentences  $F$  such that:

1.  $\mathbf{PRA} \not\vdash F$
2.  $\mathbf{PRA} \not\vdash \neg F$
3.  $\mathbf{PRA} \vdash \text{Con} \rightarrow \neg \text{Pr}(\ulcorner F \urcorner)$
4.  $\mathbf{PRA} \vdash \text{Con} \rightarrow \neg \text{Pr}(\ulcorner \neg F \urcorner)$

*Proof.* We define the following pointed Kripke model:

$$\mathcal{K} = \langle \{1, 2, 3\}, \{(1, 2), (1, 3)\}, \{(1, p), (2, p)\}, 1 \rangle$$

it can be visualized by the following graph: Where  $p$  is true in world 2. We will



further let  $\alpha$  be the following formula:

$$\neg \Box p \wedge \neg \Box \neg p \wedge \Box (\neg \Box \perp \rightarrow \neg \Box p \wedge \neg \Box \neg p)$$

We have that  $\alpha$  is true in  $\mathcal{K}$ . Since we have that:

1.  $1 \in \phi(\neg \Box p)$  since we have that  $1R3$  and that  $3 \notin \phi(p)$
2.  $1 \in \phi(\neg \Box \neg p)$  since we have that  $1R2$  and that  $2 \in \phi(p)$
3.  $1 \in \phi(\Box (\neg \Box \perp \rightarrow \neg \Box p \wedge \neg \Box \neg p))$  since we have that  $2, 3 \in \phi(\Box \perp)$

It can be shown that this model is  $\alpha$ -reflexive; the proof of this can be found in (Smorynski 1985) –

This result can be strengthened and it can be shown that the found formula is actually  $\Sigma_1$

## 6.4. Generalisations of the Completeness Theorems

In this section we have shown the Solovay's Completeness Theorems for **PRA**. But these theorems holds for a much wider class of fragments of Peano Arithmetics. It can be shown **Ref De joghn** that the theorem holds for theories  $T$  that fulfills the following to conditions:

1.  $T$  extends  $I\Delta_0 + \text{EXP}$
2. Let  $\alpha(x)$  be  $\Delta_0$  formula, then  $T$  does not prove any false sentences of the form  $\exists x\alpha(x)$

This proof does not use the Recursion theorem for creating the function  $\varphi$ , but instead uses the diagonalization lemma (i.e Lemma 2.1). The proof of this will not this will not be given here, but it can be found in **Ref artiklen**.

It is not known if Solovays theorems holds for weaker conditions than  $I\Delta_0 + \text{EXP}$ ; i.e we do not know if it holds for example for the theory  $I\Delta_0 + \Omega_1$ .

Solovay's theorems can also be proven for some fragments without the use of fixed points. The two different proof strategies seen so fair each uses fixed points. Either by using the recursion theorem or by using the digitalization theorem for a given arithmetic theory. Fedor Pakhomov has proven the theorems without the use of the fixed point lemma or the recursion theorem in (Pakhomov 2017). This newer proof might be able to give a specific lower bound on the strength of the fragments Solovays theorems holds for.

## 6.5. Solovays Completeness Theorems and Fixed Points

From table 5.1 and the arithmetical soundness theorem, we can conclude the following things:

1. From line 3, we have that  $\vdash_{\mathbf{GL}} \Box(p \leftrightarrow \Box\neg p) \leftrightarrow \Box(p \leftrightarrow \Box\perp)$ ; i.e a sentence  $F$  of **PRA** is equivalent to its own unprovability if and only if  $F$  is equivalent to the assertion that **PRA** is inconsistent. For let  $*$  be such that  $F = p^*$ , then we if  $F$  is such that  $\mathbf{PRA} \vdash F \leftrightarrow \text{Pr}(\ulcorner \neg F \urcorner)$  which is the same as  $\mathbf{PRA} \vdash (p \leftrightarrow \Box\neg p)^*$  and thus we have:  $\mathbf{PRA} \vdash \Box(p \leftrightarrow \Box\neg p)^*$  and hence  $\mathbf{PRA} \vdash \Box(p \leftrightarrow \Box\neg p)^*$ . But by the soundness theorem we get that:  $\mathbf{PRA} \vdash (\Box(p \leftrightarrow \Box\neg p) \leftrightarrow \Box(p \leftrightarrow \Box\perp))^*$ , and thus  $\mathbf{PRA} \vdash \Box(p \leftrightarrow \Box\perp)^*$  whence  $\mathbf{PRA} \vdash (p \leftrightarrow \Box\perp)^*$  and thus  $\mathbf{PRA} \vdash F \leftrightarrow \text{Pr}(\ulcorner \perp \urcorner)$ . Which shows that  $F$  is equivalent to the inconsistency of **PRA**.

## 6. Solovays Completeness Theorems

---

2. We can from line 1 and 6 infer that a sentence of **PRA** is equivalent to its own unprovability if and only if it is equivalent to the assertion that **PRA** is consistent and that a sentence of **PRA** that is equivalent to the assertion that it is unprovable if it is provable if and only if it is equivalent to the assertion that if the inconsistency of **PRA** is provable then **PRA** is provable.
3. From line 10 we can conclude that for arbitrary sentence  $F$  and  $G$  of **PRA** that:  $F$  is equivalent to the conduction of  $F$  is provable and  $G$  is true if and only if  $F$  is equivalent to the conduction of  $G$  is provable and true. Here we take  $*$  such that  $p^* = F$  and  $q^* = G$ .
4. From line 6 again we can conclude that a fixed point may have a bigger modal degree, than formula of which it is a fixed point: the formula  $\Box p \rightarrow \Box \neg p$  has modal degree 1 and the formula  $\Box \Box \perp \rightarrow \Box \perp$  has degree 2. We can conclude the same for line 7, where the fixed point has modal degree 3 and the formula it is a fixed point for has degree 2.
5. The modal formula  $\alpha$  has at most the same modal degree as the modal formula  $\sigma$ .
6. We can further see that line 1 gives that sentences of arithmetic that express their own unprovability are equivalent to the assertion arithmetic is consistent. Line 2 is the same as Löb's answer to Henkin's question, and line 4 gives that refutable sentences are those equivalent to their own consistency.

### 6.6. Concluding remarks: Implications of Solovay's Theorems

The clearest implication of the generalized versions of Solovay's Completeness Theorems is that the proof predicate of arithmetics can be axiomatized; everything there is to say about the predicate, can be deduced from the derivability conditions and Löb's theorem. This can be seen both as a positive and negative result. Positive since the proof predicate follows some simple rules and these rules does not alter even though you add more induction to the fragment of arithmetics you are working with. The last point can also be seen as a negative one; there is no way to differentiate the different fragments of arithmetics by looking at their proof predicate.

Further the proportional provability logic is not effected by Quines critique of modal logic as being unintelligible, since it has a very clear and unambiguous arith-

## 6.6. Concluding remarks: Implications of Solovay's Theorems

---

metric interpretation. This sets it apart from the more philosophical interpretations of modal logic, since these interpretations do not have a clear meaning.





## 7. Further Results in Provability Logic

{chap:Further}

In this chapter we will enrich the language  $\mathcal{L}_\square$  with both more modal operators and quantifiers. For these language a number of different results can be proven. We will also state some results about provability logic where we enrich our language  $\mathcal{L}$  with quantifiers. In the case with more modal operators the results will be positive; i.e there are versions of Solovay's completeness theorems for these kinds of modal logic. But in the case where we enrich our language with quantifiers our results will be negative; i.e we cant prove versions of Solovay's completeness theorems for these kinds of modal logic.

This chapter will skip the most of the proofs of these results, but the proofs can be found in (George. S Boolos 1993).

### 7.1. Multi-modal provability logic

We can extend our language  $\mathcal{L}_\square$  with more modality operators. We will start of by extending it with the modal operator:  $\Box$  and its dual  $\Diamond$ . In the next subsection it will be explained what we will mean with the  $\Box$ . Latter on we will extend the language with even more modal operators.

#### 7.1.1. The system GLB

**Definition 7.1** We will say that a theory  $\mathbf{T}$  is  $\omega$ -inconsistent iff for some formula  $\alpha(x)$ ,  $\mathbf{T} \vdash \exists x \alpha(x)$  and for every  $n \in \omega$  we have:  $\mathbf{T} \vdash \neg \alpha(n)$ .  $\mathbf{T}$  is called  $\omega$ -consistent iff it is not  $\omega$ -inconsistent.  $\dashv$

We have that if  $\mathbf{T}$  is  $\omega$ -consistent then  $\mathbf{T} \not\vdash \exists x x \neq x$  and thus  $\mathbf{T}$  is consistent. So  $\omega$ -consistency implies consistency; but the converse does not hold.

**Definition 7.2** A sentence  $\alpha$  is  $\omega$ -inconsistent in  $\mathbf{T}$  if the axioms of  $\mathbf{T}$  plus  $\alpha$  is  $\omega$ -inconsistent.  $\alpha$  is  $\omega$ -consistent iff it is not  $\omega$ -inconsistent.

We further say that a sentence  $\alpha$  is  $\omega$ -provable in  $\mathbf{T}$  iff  $\neg \alpha$  is  $\omega$ -inconsistent with  $\mathbf{T}$ .  $\dashv$

## 7. Further Results in Provability Logic

---

It is clear that if  $\alpha$  is provable in **T** then it is  $\omega$ -provable in **T**. We will introduce a new modal system **GLB**, where **B** stands for bimodal, where we add two new modal operators:  $\Box$  and  $\Diamond$  to our modal language  $\mathcal{L}_\Box$ . We will call this new language for  $\mathcal{L}_\Box$

**Definition 7.3** The language  $\mathcal{L}_\Box$  is the language  $\mathcal{L}_\Box$  extend with the modal operator  $\Box$ , where the syntax  $\Box$  is the same as that of  $\Box$ . The operator  $\Diamond$  is defined as the dual of  $\Box$   $\dashv$

We now define the system of **GLB**. We will later on discuss the semantics of **GLB**, but this is not of important right now.

**Definition 7.4** The axioms of **GLB** are all tautologies and all sentences of the following kind:

$$\text{A1 } \Box(\alpha \rightarrow \beta) \rightarrow (\Box\alpha \rightarrow \Box\beta)$$

$$\text{A2 } \Box(\alpha \rightarrow \beta) \rightarrow (\Box\alpha \rightarrow \Box\beta)$$

$$\text{A3 } \Box(\Box\alpha \rightarrow \alpha)\Box\alpha$$

$$\text{A4 } \Box(\Box\alpha \rightarrow \alpha)\Box\alpha$$

$$\text{A5 } \Box\alpha \rightarrow \Box\alpha$$

$$\text{A6 } \neg\Box\alpha \rightarrow \Box\neg\Box\alpha$$

MP Modus ponens

Nec  $\Box$ -necessitation; from  $\alpha$  infer  $\Box\alpha$ .

$\dashv$

It is clear that  $\Box$ -necessitation is a rule of **GLB**, since we have that if **GLB**  $\vdash \alpha$  then **GLB**  $\vdash \Box\alpha$  and by A5 we then have **GLB**  $\vdash \Box\alpha$ .

The next goal is to prove an arithmetical soundness theorem for **GLB**. For this end we need the following definition:

**Definition 7.5** We call the following rule for the  $\omega$ -rule: Infer  $\forall x\alpha(x)$  for all  $\alpha(\bar{n})$ , where  $n \in \omega$ . A sentence is provable under the  $\omega$ -rule in **T** if it belongs to all classes containing the axioms of **T** and closed under MP, Nec and the  $\omega$ -rule. We further say that a sentence  $F$  is provable in **PRA** by *one application of the  $\omega$ -rule* if for some formula  $G(x)$  we have that **PRA**  $\vdash G(\bar{n})$  for all  $n$  and **PRA**  $\vdash \forall xG(x) \rightarrow F$ .  $\dashv$

It can be shown that a sentence  $F$  is  $\omega$ -provable if and only if it is provable by one application of the  $\omega$ -rule. With this we can define the notion of an  $\omega$ -proof.

**Definition 7.6** A sentence  $\forall xG(x) \rightarrow F$  is called an  $\omega$ -proof of  $F$  if  $\mathbf{PRA} \vdash G(\bar{n})$  for all  $n$  and  $\mathbf{PRA} \vdash \forall xG(x) \rightarrow F$ .  $\dashv$

It is clear that if  $F$  has an  $\omega$ -proof then it is  $\omega$ -provable.

We will state one last definition in the same vein as the others above:

**Definition 7.7**  $\mathbf{PRA}^+$  is the theory extending  $\mathbf{PRA}$  with all sentences  $\forall xG(x)$  such that for every  $n \in \omega$  we have that  $\mathbf{PRA} \vdash G(\bar{n})$ .  $\dashv$

The following theorem can then be proven about the relationships between the above definitions:

{thm:omega}

**Theorem 7.1** The following are equivalent:

1.  $F$  is  $\omega$ -provable.
2.  $\mathbf{PRA}^+ \vdash F$ .
3.  $F$  is provable by one application of the  $\omega$ -rule.
4. There is an  $\omega$ -proof of  $F$ .

We need a few more definitions before we can state the arithmetical soundness theorem for **GLB**:

**Definition 7.8** Let  $\omega\text{Prov}(y, x)$  be the formula of  $\mathbf{PRA}$  that formalizes that  $y$  is the code of an  $\omega$ -proof of  $x$ , and let  $\omega\text{Pr}(x) = \exists y\omega\text{Prov}(y, x)$ .  $\dashv$

We then have that  $\omega\text{Pr}(x)$  by theorem 7.1 is provable coextensive with the formulas stating in  $\mathbf{PRA}$  the following properties: " $\omega$ -provable", "provable in  $\mathbf{PRA}^+$ " and "provable by one application of the  $\omega$ -rule". This means that for each sentence  $F$  in  $\mathbf{PRA}$  that we have:

$$\mathbf{PRA} \vdash \text{Pr}(\ulcorner F \urcorner) \rightarrow \omega\text{Pr}(\ulcorner F \urcorner)$$

With these results we can now extend the interpretation of  $\mathcal{L}_\square$  in  $\mathbf{PRA}$  to one of  $\mathcal{L}_\sqcup$ :

**Definition 7.9** An interpretation  $*$  of  $\mathcal{L}_\sqcup$  in  $\mathbf{PRA}$  is an extension of an interpretation  $\mathcal{L}_\square$  in  $\mathbf{PRA}$  where we add the following:

$$\sqcup(\alpha)^* = \omega\text{Pr}(\ulcorner \alpha^* \urcorner)$$

$\dashv$

## 7. Further Results in Provability Logic

---

We can now state the arithmetical soundness theorem for **GLB**:

**Theorem 7.2** (The arithmetical soundness theorem for **GLB**) For any modal formula  $\alpha$  and interpretation of  $\mathcal{L}_{\Box}$  in **PRA** we have that:

$$\vdash_{\mathbf{GLB}} \alpha \rightarrow \mathbf{PRA} \vdash \alpha^*$$

We omit the proof here. We can not prove the arithmetical completeness theorem for **GLB** without the help of another multi modal logic.

This is because there is no good Kripke semantics for the logic **GLB**. We will start of by defining a Kripke modal for the logic **GLB**; this definitions is a extension of a "normal" frame, but with two relations instead of one

**Definition 7.10** A frame for a modal logic with two modal operators is a triple  $\mathcal{H} = \langle W, R, R_1 \rangle$  where both  $R$  and  $R_1$  are relations on  $W$ . A model is a quadruple  $\mathcal{K} = \langle W, R, R_1, \phi \rangle$  where  $\phi$  is a valuation function on  $W$ . The truth of modal formula  $\alpha$  in a given node  $w \in W$  is defined in the obvious way, and the two main clauses of the definition is:

1.  $\vdash_w^{\mathcal{K}} \Box \alpha$  iff for all  $v \in W$  such that  $wRv$  we have  $\vdash_v^{\mathcal{K}} \alpha$
2.  $\vdash_w^{\mathcal{K}} \Box_1 \alpha$  iff for all  $v \in W$  such that  $wR_1v$  we have  $\vdash_v^{\mathcal{K}} \alpha$

⊥

The problem here for is that no matter what the relation  $R_1$  will be the empty relation  $W$  and therefor  $\Box_1 \perp$  will be valid and thus contradict that  $\vdash_{\mathbf{GLB}} \Box_1 \perp$  from the arithmetical soundness of **GLB**. Giorgie Dzhaparidze found a way around this by looking at another multi modal logic, which will be introduced in the next section. His result was improved by Konstantin Ignatiev .

### 7.1.2. The system **IDzh** and completeness of **GLB**

Since **GLB** does not have any good Kripke semantic we will at a multi modal logic that has such a semantic; we will call this modal logic for **IDzh** Ignatiev and Dzhaparidze.

**Definition 7.11** The language of **IDzh** is the same as **GLB**. Further the axioms of **IDzh** are all tautologies and the following modal formulas:

$$\text{A1 } \Box(\alpha \rightarrow \beta) \rightarrow (\Box\alpha \rightarrow \Box\beta)$$

$$\text{A2 } \Box_1(\alpha \rightarrow \beta) \rightarrow (\Box_1\alpha \rightarrow \Box_1\beta)$$

$$\text{A3 } \Box(\Box\alpha \rightarrow \alpha) \rightarrow \Box\alpha$$

A4  $\Box(\Box\alpha \rightarrow \alpha) \rightarrow \Box\alpha$

A5  $\Box\alpha \rightarrow \Box\Box\alpha$

A6  $\neg\alpha \rightarrow \Box\neg\Box\alpha$ .

MP Modus ponens

Nec  $\Box$ -necessitation

Nec<sub>1</sub>  $\Box$ -necessitation

Here we add the inference Nec<sub>1</sub> since this rule can not be proven from the other rules and axioms in this modal logic.  $\dashv$

The modal logic **IDzh** can be shown to be weaker than **GLB** and thus **IDzh**  $\subseteq$  **GLB**.

The multi modal logic **IDzh** has the following Kripke semantic for which a soundness and completeness theorem can be proven.

**Definition 7.12** An **IDzh**-model is a quadruple  $\mathcal{K} = \langle W, R, R_1, \phi \rangle$ , where  $W$  is a finite non-empty set,  $\phi$  is a valuation function  $W$ , and  $R$  and  $R_1$  are transitive irreflexive relation on  $W$  such that for all  $w, v_1, v_2 \in W$  we have that:

If  $wR_1v_1$  then  $wRv_2$  if and only if  $v_1Rv_2$

$\dashv$

This means that we can not have that  $wRv$  and  $wR_1v$  since then  $vRv$ ; contradicting the irreflexivity  $R$

For such models the following theorem can be proven:

**Theorem 7.3**  $\alpha$  is valid in all **IDzh**-models if and only if  $\vdash_{\mathbf{IDzh}} \alpha$ .

This theorem shows that **IDzh** has a natural Kripke semantic to which it is sound and complete.

We will need the following definitions to state the theorem from which the arithmetical completeness theorem for **GLB** follows:

**Definition 7.13** For any modal formula  $\alpha$  we define  $\Delta\alpha$  as the formula:

$$\alpha \wedge \Box\alpha \wedge \Box\Box\alpha \wedge \Box\Box\Box\alpha$$

We define the formula  $\Psi\alpha$  as:

$$\bigwedge_{\Box\beta \in S(\alpha)} \Delta(\Box\beta \rightarrow \Box\Box\beta)$$

## 7. Further Results in Provability Logic

+

It is clear that  $\vdash_{\mathbf{GLB}} \Psi\alpha$  To show completeness of **GLB** the following three statements should be proven to be equivalent:

1.  $\vdash_{\mathbf{IDzh}} \Psi\alpha \rightarrow \alpha$
2.  $\vdash_{\mathbf{GLB}} \alpha$
3.  $\mathbf{PRA} \vdash \alpha^*$  for all  $*$

We have that (1)  $\Rightarrow$  (2) since  $\mathbf{IDzh} \subset \mathbf{GLB}$  and the fact that  $\vdash_{\mathbf{GLB}} \Psi\alpha$ . That (2)  $\Leftrightarrow$  (3) is just the arithmetical soundness theorem for **GLB**. The proof of (3)  $\Rightarrow$  (1) will obviously prove the arithmetical completeness theorem for **GLB**. The proof of this can be found in (George. S Boolos 1993) and is beyond the scope of this project.

We can also look at the multi modal logic **GLSB** which is the modal logic that have all the theorems of **GLB** and all formulas:  $\Box\alpha \rightarrow \alpha$  as axioms, and which only have modus ponens as its rule of inference. A variant of Solovay's second completeness theorem can be shown for this multi modal logic.

### 7.1.3. The system GLP

We will end this section by introducing the system **GLP** The language  $\mathcal{L}_{[n]}$  of **GLP** is an extension of  $\mathcal{L}_{\Box}$ , where instead of  $\Box$  we write  $[0]$  and we further add a countable infinite amount of boxes:  $[1], [2], \dots$  representing provability in  $\mathbf{PRA}^+$ ,  $\mathbf{PRA}^{++}$  and so on, where  $\mathbf{PRA}^{++}$  is  $\mathbf{PRA}^+$  with all formulas of the form  $\forall x G(x)$  such that for every  $n \in \omega$  such that  $\mathbf{PRA}^+ \vdash G(\bar{n})$  added. We define their duals  $\langle 0 \rangle, \langle 1 \rangle, \langle 2 \rangle, \dots$  in the obvious way and these represent consistency,  $\omega$ -consistency,  $\omega$ - $\omega$ -consistency and so on. We can now define the system **GLP**:

**Definition 7.14** The axioms and inferences rules of **GLP** are the following:

$$\text{A1 } [n](\alpha \rightarrow \beta) \rightarrow ([n]\alpha \rightarrow [n]\beta)$$

$$\text{A2 } [n]([n]\alpha \rightarrow \alpha) \rightarrow [n]\alpha$$

$$\text{A3 } [n]\alpha \rightarrow [n+1]\alpha$$

$$\text{A4 } \neg[n]\alpha \rightarrow [n+1]\neg[n]\alpha$$

MP Modus Ponens

Nec<sub>0</sub>  $[0]$ -necessiation

⊢

Following Dzhaparidze the arithmetical completeness theorem can be proven for this system of multi modal logic. The proof follows that of the proof of the arithmetical completeness theorem for **GLB**.

Again we can look at the multi modal logic **GLSP** which has all theorems and sentences  $[n]\alpha \rightarrow \alpha$  as its axioms and which only have modus ponens as its inference rule. Again a variant of Solovay's second completeness theorem holds for this logic.

So Solovay's completeness theorems generalize nicely to multi modal logics; i.e the concept of  $\omega$ -provability (and beyond) can also be axiomatized.

The proofs of these results are rather involved, but they still follow the same idea as Solovay's original idea. This shows how deep the concept of embedding modal logic into arithmetic is. But this fundamental idea does not generalize to all extensions of **GL**, which will be discussed in the next section.

## 7.2. Quantified provability logic

There is another way to extend the logic **GL**; we can add quantifiers to the language. We will call this logic for quantified modal logic (QML). The first goal is to define what a formula is:

**Definition 7.15**  $\alpha$  is a formula of QML if and only if it can be obtained from a formula of first order logic by replacing occurrences of the negation sign " $\neg$ " with occurrences of  $\Box$ . ⊢

There is not a version of Solovay's completeness theorems for this logic; i.e it is not arithmetically complete with respect to any fragment of arithmetics. Furthermore this logic does not have the fixed point property and is not complete with respect to any class of Hintikka frames. These results can actually be shown to hold for an even more conservative extension of **GL**: one where we only allow one place predicates and where we do not permit nestings of the  $\Box$ -operator. Further information on this topic and proofs of these statements can be found in (George. S Boolos 1993), but this goes way beyond the scope of this project and are left out.





## 8. Bibliography

- Aristoteles (2021). *Metafysik*. Forlaget Klim.
- (1971). *Aristotle’s Categories and De interpretatione*. Clarendon Aristotle series. Oxford: Oxford University Press.
- Artemov, Sergei (2007). “Handbook of modal logic: 1st ed”. In: *Handbook of modal logic*. Ed. by P Blackburn, J van Benthem, and F Wolter. Vol. 3. Studies in logic and practical reasoning. Elsevier. Chap. 16.
- Artemov, Sergei N. and Lev D. Beklemishev (n.d.). “Provability Logic”. In: *Handbook of Philosophical Logic, 2nd Edition*. Handbook of Philosophical Logic. Dordrecht: Springer Netherlands, pp. 189–360.
- Barry Cooper, S (2004). *Computability Theory*. Studies in logic and the foundations of mathematics. Boca Raton: Chapman & Hall/CRC.
- Blackburn, Patrick (2002). *Modal logic*. Ed. by Maarten de Rijke and Yde Venema. Cambridge tracts in theoretical computer science ; 53. Cambridge: Cambridge University Press.
- Blackburn, Patrick, Johan van Benthem, and Frank Wolter, eds. (2007). *Handbook of modal logic: 1st ed*. Vol. 3. Studies in logic and practical reasoning. Amsterdam: Elsevier.
- Boolos, George and Giovanni Sambin (1991). “Provability: The Emergence of a Mathematical Modality”. In: *Studia logica* 50.1, pp. 1–23.
- Boolos, George S., John P. Burgess, and Richard C. Jeffrey (2007). *Computability and logic*. 5th ed. Cambridge: Cambridge University Press.
- Boolos, George. S (1979). *The unprovability of consistency : an essay in modal logic*. Cambridge: Cambridge University Press.
- (1993). *The logic of provability*. Cambridge: Cambridge University Press.
- Chang, C. C. and H Jerome Keisler (2012). *Model Theory*. 3. th. Mineola: Dover Publications, inc.
- Cohen, Paul (1966). *Set Theory and the Continuum Hypothesis*. W. A. Benjamin, Inc.
- Copeland, B. Jack (2020). “The Church-Turing Thesis”. In: *The Stanford Encyclopedia of Philosophy*. Ed. by Edward N. Zalta. Summer 2020. Metaphysics Research Lab, Stanford University.
- Cutland, N.J (1980). *Computability : an introduction to recursive function theory*. Cambridge: Cambridge University Press.

## 8. Bibliography

---

- Davis, Martin, ed. (1965). *The undecidable : basic papers on undecidable propositions, unsolvable problems and computable functions*. Hewlett, N. Y: Raven Press.
- Enderton, Herbert B. (2001). *A Mathematical Introduction to Logic*. Academic Press.
- Garson, James W (2006). *Modal Logic for Philosophers*. Cambridge: Cambridge University Press.
- Gödel, Kurt (1986a). “Eine Interpretation des intuitionistischen Aussagenkalküls”. In: *Collected works*. Ed. by Solomon Feferman. Oxford University Press, pp. 300–301.
- (1986b). “Über formal unentschiedbare Sätze der Principia Mathematica und verwandter System I”. In: *Collected works*. Ed. by Solomon Feferman. Oxford University Press, pp. 144–195.
- Goldblatt, Robert (2003). “Mathematical modal logic: A view of its evolution”. In: *Journal of applied logic* 1.
- Hájek, Petr. and P Pudlak (1998). *Metamathematics of First-Order Arithmetic*. Perspectives in Mathematical Logic. Berlin: Springer.
- Hendricks, Vincent Fella and Stig Andur Pedersen (2011). *Moderne elementær logik*. Ed. by Vincent Fella Hendricks and Stig Andur Pedersen. 2. revider. Kbh: Automatic Press.
- Hilbert, David (1926). “Über das Unendliche”. In: *Mathematische annalen* 95.1, pp. 161–190.
- (1930). *Grundlagen der Geometrie, 7. umgearb. und verm. Aufl.* Wissenschaft und Hypothese ; 7. Leipzig und Berlin.
- Hintikka, K. Jaakko J. (1962). *Knowledge and belief*. Ed. by Jaakko. Hintikka. Contemporary philosophy. Ithaca, N. J: Cornell Univ. Press.
- (1969). *Models for modalities : Selected essays*. Ed. by Jaakko. Hintikka. Dordrecht: D. Reidel.
- Hughes, G.E and M.J Cresswell (1971). *An introduction to modal logic*. Methuen.
- Japaridze, G and Dick de Jongh (1998). “Handbook of proof theory”. In: *Handbook of proof theory*. Ed. by Samuel R. Buus. Studies in logic and the foundations of mathematics. Amsterdam ; Elsevier. Chap. 7, pp. 475–546.
- Jørgensen, Thorvald (2021). “Project outside the course scope: Introduction to Modal Logic”. In: URL: <https://github.com/thorvald94/Modal-logic/blob/main/FagProjekt.pdf>.
- (2022). “Project outside the course scope: Gödel’s Incompleteness Theorems”. In: URL: <https://github.com/thorvald94/GodelIncompleteness/blob/main/project.pdf>.
- Kleene, S.C. (1952). *Introduction to metamathematics*. 8. repr. Bibliotheca mathematica ; vol. 1. Groningen: Wolters-Noordhoff.

- Kripke, Saul A (1963). “Semantical Analysis of Modal Logic I Normal Modal Propositional Calculi”. In: *Mathematical logic quarterly* 9.
- Lemmon, E. J. (1977). *An introduction to modal logic : the "Lemmon notes"*. Ed. by Dana. Scot. American philosophical quarterly. Monograph series ; monograph no. 11. Oxford: Basil Blackwell.
- Lindström, Per (1996). “Provability logic-a short introduction”. In: *Theoria (Lund, Sweden)* 62.1-2, pp. 19–61.
- (1997). *Aspects of Incompleteness*. Berlin/Heidelberg: Springer.
- Liu, Yang (2013). *Incompleteness result and provability logic*.
- Łukasiewicz, Jan. (1957). *Aristotle’s syllogistic from the standpoint of modern formal logic*. 2. ed. enl. Oxford: Clarendon.
- Mendelson, Elliott (1964). *Introduction to Mathematical Logic*. Van Norstrand.
- Owens, James. C (1973). “Diagonalization and the Recursion Theorem”. In: *Notre Dame Journal of Formal Logic* XIV.No 1, pp. 95–99.
- Pakhomov, Fedor (2017). “Solovay’s completeness without fixed points”. In: URL: <http://arxiv.org/licenses/nonexclusive-distrib/1.0>.
- Quine, W. V (1966). “Necessary Truth”. In: *The Ways of Paradox and other essays*. New York: Random House, pp. 48–56.
- R. Buss, Samuel, ed. (1998a). *Handbook of proof theory*. Studies in logic and the foundations of mathematics. Amsterdam ; Elsevier.
- (1998b). “Handbook of proof theory”. In: *Handbook of proof theory*. Ed. by Samuel R. Buus. Studies in logic and the foundations of mathematics. Amsterdam ; Elsevier. Chap. 1, pp. 79–147.
- Raatikainen, Panu (2022). “Gödel’s Incompleteness Theorems”. In: *The Stanford Encyclopedia of Philosophy*. Ed. by Edward N. Zalta. Spring 2022. Metaphysics Research Lab, Stanford University.
- Reidhaar-Olson, L (1990). “A new proof of the fixed-point theorem of provability logic”. In: *Notre Dame journal of formal logic* 31.1, pp. 37–43.
- Rosser, John Barkley (1936). “Extensions of some theorems of Gödel and Church”. In: *Journal of Symbolic Logic*.
- Segerberg, Krister (1971). *An Essay in Classical Modal Logic*. Filosofiska Studier, vol.13. Uppsala Universitet.
- Sipser, Michael (2013). *Introduction to the theory of computation*. 3. ed. Boston: Thomson Course Technology.
- Skolem, Th. (1923). *Begründung der elementaren Arithmetik durch die rekurrende Denkweise ohne Anwendung scheinbarer veränderlichen mit unendlichem Ausdehnungsbereich*. Videnskapsselskapets skrifter. Mat.-naturv. klasse ; 1923:6. Kristiania: Jacob Dybwad.
- Smiley, T J (1963). “The Logical Basis of Ethics”. In: *Acta Philosophica Fennica* 16, pp. 237–.

## 8. Bibliography

---

- Smorynski, Craig (1977a). “Beth’s theorem and self-referential sentences”. In: *Logic Colloquium 77*. Ed. by A Macintyre, L Pacholski, and J Paris. Studies in logic and the foundations of mathematics. Amsterdam, New York, Oxford: North.Holland Publishing Company, pp. 253–261.
- (1977b). “The Incompleteness Theorems”. In: *The Handbook of Mathematical logic*. Ed. by John Barwise. North-Holland Publishing Company.
  - (1985). *Self-Reference and modal logic*. Springer-verlag.
- Soare, Robert I (1987). *Recursively Enumerable Sets and Degrees : A Study of Computable Functions and Computably : Generated Sets*. Perspectives in Mathematical Logic. Berlin ; Springer-Verlag.
- (2016). *Turing Computability: Theory and Applications*. Theory and Applications of Computability. Berlin, Heidelberg: Springer Berlin / Heidelberg.
- Solovay, Robert M (1976). “Provability interpretations of modal logic”. In: *Israel journal of mathematics* 25.3-4.
- Urbaniak, Rafal and Pawel Pawlowski (2018). “Logics of (Formal and Informal) Provability”. In: *Introduction to Formal Philosophy*. Springer Undergraduate Texts in Philosophy. Springer International Publishing, pp. 191–237.
- Verbrugge, Rineke (L.C.) (2017). “Provability Logic”. In: *The Stanford Encyclopedia of Philosophy*. Ed. by Edward N. Zalta. Fall 2017. Metaphysics Research Lab, Stanford University.
- Whitehead, Alfred North. and Bertrand Russell (1973). *Principia mathematica*. 2. ed., reprinted. Cambridge: Cambridge University Press.

## A. Necessitation free definition of GL and GLS

There is another way to define the modal logic **GLS**. The definition of this section is the one Smorynski uses in Smorynski 1985

In this section we will take **K4** as our base logic and see **GL** as an extension of this. We will need the following notation to define this. Given a set  $\Gamma$ :

1.  $\Gamma \vdash_{\mathbf{MP}} \alpha$  if  $\alpha$  is derivable from  $\Gamma$  by only using  $R1$
2.  $\Gamma \vdash_{\mathbf{Nec}} \alpha$  if  $\alpha$  is derivable from  $\Gamma$  by using  $R1$  and  $R2$ .

We will further define the following set for any set  $\Gamma$ :

$$\Gamma^{\mathbf{Nec}} = \Gamma \cup \{\Box\beta : \beta \in \Gamma\}$$

This set allows us to avoid Nec by the axioms of **K4**, which the following lemma shows:

**Lemma A.1** Let  $\Gamma$  include all the instances of axioms of **K4**. Then:

$$\Gamma \vdash_{\mathbf{Nec}} \alpha \Leftrightarrow \Gamma^{\mathbf{Nec}} \vdash_{\mathbf{MP}} \alpha$$

*Proof.* The way  $\Leftarrow$  is trivial. So we will only show  $\Rightarrow$ . To prove this, it is enough to show that the set of theorems of  $\Gamma^{\mathbf{Nec}}$  is closed under  $R2$ ; i.e to show that:

$$\Gamma^{\mathbf{Nec}} \vdash \alpha \Rightarrow \Gamma^{\mathbf{Nec}} \vdash_{\mathbf{MP}} \Box\alpha$$

Let  $\Gamma^{\mathbf{Nec}} \vdash_{\mathbf{MP}} \alpha$ . Then we have a sequence  $\alpha_1, \dots, \alpha_n = \alpha$ , where for each  $1 \leq i \leq n$  we have that  $\alpha_i$  is one of the following three:

1. We have that it is an axiom  $\alpha_i \in \Gamma$
2. It is the necessitation  $\Box\beta_1$  of an axiom  $\beta_1 \in \Gamma$
3. It is a consequence of **MP** of  $\alpha_j, \alpha_k = \alpha_j \rightarrow \alpha_i$  where  $j, k < i$ .

We will show by induction on the length  $i$  of a 'initial' segment  $\alpha_1, \dots, \alpha_i$  that  $\Gamma^{\mathbf{Nec}} \vdash_{\mathbf{MP}} \Box\alpha_i$ .

## A. Necessitation free definition of **GL** and **GLS**

---

1. If we have that  $\alpha_i \in \Gamma$ . Then by definition we have that  $\Box\alpha_i \in \Gamma^{\mathbf{Nec}}$  and thus  $\Gamma^{\mathbf{Nec}} \vdash_{\mathbf{MP}} \Box\alpha_i$ .
2. If  $\alpha_i = \Box\beta_i$  for some  $\beta_i \in \Gamma$ , then we can make the following deduction:

$$\begin{array}{ll}
 \Gamma^{\mathbf{Nec}} \vdash_{\mathbf{MP}} \Box\beta_i & \text{Since } \Box\beta_i \in \Gamma^{\mathbf{Nec}} \\
 \vdash_{\mathbf{NP}} \Box\beta_i \rightarrow \Box\Box\beta_i & \text{By } \mathbf{4} \\
 \vdash_{\mathbf{MP}} \Box\Box\beta_i (= \Box\alpha_i) & \text{By } \mathbf{MP}
 \end{array}$$

Which shows the result.

3. If  $\alpha_i$  follows form  $\alpha_j, \alpha_k = \alpha_j \rightarrow \alpha_i$  by **MP** we have by the induction hypothesis that  $\Gamma^{\mathbf{Nec}} \vdash_{\mathbf{MP}} \Box\alpha_j$  and  $\Gamma^{\mathbf{Nec}} \vdash_{\mathbf{MP}} \Box(\alpha_j \rightarrow \alpha_i)$ . By **K** we also have that  $\Gamma^2 \vdash_{\mathbf{MP}} \Box\alpha_j \wedge \Box(\alpha_j \rightarrow \alpha_i) \rightarrow \Box\alpha_i$  and by propositional logic we get that:  $\Gamma^2 \vdash_{\mathbf{MP}} \Box\alpha_i$ .

This ends the induction and the lemma will follow.  $\dashv$

By this lemma we can get a **Nec**-free definition of **K4** and **GL** that we will call **K4<sup>Nec</sup>** and **GL<sup>Nec</sup>**. We can further by the deduction theorem from propositional logic get the following:

**Lemma A.2**  $\Gamma \vdash \alpha$  iff there is a finite set  $\{\gamma_0, \dots, \gamma_{n-1}\} \subseteq \Gamma$  such that:  $K \vdash \wedge \gamma_i \rightarrow \alpha$

From all this we can define **GLS** in an alternative way:

**Definition A.1** The system of modal logic **GLS** is defined as the system of propositional modal logic that have all the theorems of **GL<sup>nec</sup>** and addition of the axiom of reflexion:

*Refl*:  $\Box\alpha \rightarrow \alpha$

and which sole rule of inference is *modus ponens*.  $\dashv$