

Twisting cochains and twisted complexes

Simplicial methods in complex-analytic geometry

Tim Hosgood

24/07/19

Université d'Aix-Marseille

<https://thosgood.github.io>

Plan

History

Twisting cochains (OTT)

- The bicomplex

- The total complex

- Why this emphasis on the first index?

- Maurer-Cartan

Twisted complexes (BK)

- Pretriangulated vs. triangulated

- Generalisation of twisting cochains

Other fun things

History

- Edgar H Brown. “Twisted tensor products, I”. In: *Annals of Mathematics* 69.1 (1959), pp. 223–246.
- John C Moore. “Differential homological algebra”. In: *Actes du Congres International des Mathématiciens* 1 (1970), pp. 335–339.

- Domingo Toledo and Yue Lin L Tong. “A parametrix for δ and Riemann-Roch in Čech theory”. In: *Topology* 15.4 (1976), pp. 273–301.
- Domingo Toledo and Yue Lin L Tong. “Duality and Intersection Theory in Complex Manifolds. I”. In: *Mathematische Annalen* 237 (1978), pp. 41–77.
- Nigel R O’Brian, Domingo Toledo, and Yue Lin L Tong. “The Trace Map and Characteristic Classes for Coherent Sheaves”. In: *American Journal of Mathematics* 103.2 (1981), pp. 225–252.

- A I Bondal and M M Kapranov. “Enhanced Triangulated Categories”. In: *Math. USSR Sbornik* 70.1 (1991), pp. 1–15.
- Giovanni Faonte. *Simplicial nerve of an A-infinity category*. 2015. arXiv: 1312.2127 [math.AT].

Twisting cochains (OTT)

Definition (Stein spaces)

A complex-analytic¹ manifold Y is said to be *Stein* if it is

1. *holomorphically convex*; and
2. *holomorphically separable*.

¹analytic = \mathcal{O}_Y is holomorphic functions, Y has the \mathbb{C}^n -induced topology;
algebraic = \mathcal{O}_Y is algebraic functions, Y has the Zariski topology.

²*Locally finite, Stein, and trivialising* (for the bundles in question).

Nice spaces

Definition (Stein spaces)

A complex-analytic¹ manifold Y is said to be *Stein* if it is

1. *holomorphically convex*; and
2. *holomorphically separable*.

Motto

Stein things are nice.

¹analytic = \mathcal{O}_Y is holomorphic functions, Y has the \mathbb{C}^n -induced topology;
algebraic = \mathcal{O}_Y is algebraic functions, Y has the Zariski topology.

²*Locally finite, Stein, and trivialising* (for the bundles in question).

Nice spaces

Definition (Stein spaces)

A complex-analytic¹ manifold Y is said to be *Stein* if it is

1. *holomorphically convex*; and
2. *holomorphically separable*.

Motto

Stein things are nice.

Throughout, X is a complex-analytic manifold with a nice² cover $\mathcal{U} = \{U_\alpha\}_{\alpha \in I}$.

¹analytic = \mathcal{O}_Y is holomorphic functions, Y has the \mathbb{C}^n -induced topology;
algebraic = \mathcal{O}_Y is algebraic functions, Y has the Zariski topology.

²*Locally finite*, *Stein*, and *trivialising* (for the bundles in question).

Endomorphisms of bounded-graded modules

Let $V = \{V_\alpha^\bullet\}$ be a collection of *bounded-graded* \mathcal{O}_{U_α} -modules:

$$V_\alpha^\bullet = \bigoplus_{q \in \mathbb{N}} V_\alpha^q \quad \text{such that } V_\alpha^q \text{ is zero for all but finitely many } q.$$

Endomorphisms of bounded-graded modules

Let $V = \{V_\alpha^\bullet\}$ be a collection of *bounded-graded* \mathcal{O}_{U_α} -modules:

$$V_\alpha^\bullet = \bigoplus_{q \in \mathbb{N}} V_\alpha^q \quad \text{such that } V_\alpha^q \text{ is zero for all but finitely many } q.$$

Think of a bounded chain complex of vector bundles, but without the information of a differential.

Endomorphisms of bounded-graded modules

Let $V = \{V_\alpha^\bullet\}$ be a collection of *bounded-graded* \mathcal{O}_{U_α} -modules:

$$V_\alpha^\bullet = \bigoplus_{q \in \mathbb{N}} V_\alpha^q \quad \text{such that } V_\alpha^q \text{ is zero for all but finitely many } q.$$

Think of a bounded chain complex of vector bundles, but without the information of a differential.

Definition (Endomorphisms)

The collection of *degree- q endomorphisms* $\text{End}^q(V)$ of V is, over each $U_{\alpha_0 \dots \alpha_p}$, given by

$$\text{End}^q(V)|_{U_{\alpha_0 \dots \alpha_p}} = \bigoplus_{i \in \mathbb{Z}} \text{Hom}(V_{\alpha_p}^i|_{U_{\alpha_0 \dots \alpha_p}}, V_{\alpha_0}^{i+q}|_{U_{\alpha_0 \dots \alpha_p}}).$$

Endomorphisms of bounded-graded modules

Let $V = \{V_\alpha^\bullet\}$ be a collection of *bounded-graded* \mathcal{O}_{U_α} -modules:

$$V_\alpha^\bullet = \bigoplus_{q \in \mathbb{N}} V_\alpha^q \quad \text{such that } V_\alpha^q \text{ is zero for all but finitely many } q.$$

Think of a bounded chain complex of vector bundles, but without the information of a differential.

Definition (Endomorphisms)

The collection of *degree- q endomorphisms* $\text{End}^q(V)$ of V is, over each $U_{\alpha_0 \dots \alpha_p}$, given by

$$\text{End}^q(V)|_{U_{\alpha_0 \dots \alpha_p}} = \bigoplus_{i \in \mathbb{Z}} \text{Hom}(V_{\alpha_p}^i|_{U_{\alpha_0 \dots \alpha_p}}, V_{\alpha_0}^{i+q}|_{U_{\alpha_0 \dots \alpha_p}}).$$

Warning

The maps are from the α_p part to the α_0 part.

The deleted Čech complex

Definition (Deleted Čech complex)

Define the chain complex $(\hat{\mathcal{C}}^\bullet(\mathcal{U}, \text{End}^\circ(V)), \hat{\delta})$ by

$$\hat{\mathcal{C}}^p(\mathcal{U}, \text{End}^q(V)) = \bigoplus_{(\alpha_0, \dots, \alpha_p)} \text{End}^q(V)|_{U_{\alpha_0 \dots \alpha_p}}$$

(where $\text{End}^q(V)|_{U_{\alpha_0 \dots \alpha_p}} = 0$ if $U_{\alpha_0 \dots \alpha_p} = \emptyset$) with the **deleted Čech differential**

$$\hat{\delta}: \hat{\mathcal{C}}^p(\mathcal{U}, \text{End}^q(V)) \rightarrow \hat{\mathcal{C}}^{p+1}(\mathcal{U}, \text{End}^q(V))$$

$$(\hat{\delta}c)_{\alpha_0 \dots \alpha_{p+1}} = \sum_{i=1}^p (-1)^i c_{\alpha_0 \dots \hat{\alpha}_i \dots \alpha_{p+1}}.$$

A notational note

We use $\hat{\mathcal{C}}$ and $\hat{\delta}$ for the *deleted* Čech objects and \mathcal{C} and δ for the ‘full’ Čech objects.

Further structure

- If V has a differential then this gives us a *bicomplex*.

Further structure

- If V has a differential then this gives us a *bicomplex*.
- There is a natural multiplication structure given by composition:

$$(c^{p,q} \cdot \tilde{c}^{\tilde{p},\tilde{q}})_{\alpha_0 \dots \alpha_{p+\tilde{p}}} = (-1)^{q\tilde{p}} c_{\alpha_0 \dots \alpha_p}^{p,q} \tilde{c}_{\alpha_p \dots \alpha_{p+\tilde{p}}}^{\tilde{p},\tilde{q}}.$$

Further structure

- If V has a differential then this gives us a *bicomplex*.
- There is a natural multiplication structure given by composition:

$$(c^{p,q} \cdot \tilde{c}^{\tilde{p},\tilde{q}})_{\alpha_0 \dots \alpha_{p+\tilde{p}}} = (-1)^{q\tilde{p}} c_{\alpha_0 \dots \alpha_p}^{p,q} \tilde{c}_{\alpha_p \dots \alpha_{p+\tilde{p}}}^{\tilde{p},\tilde{q}}.$$

- We could define the same complex for an arbitrary bounded graded vector bundle, i.e.

$$\hat{\mathcal{C}}^p(\mathcal{U}, V^q) = \bigoplus_{(\alpha_0, \dots, \alpha_p)} V_{\alpha_0}^q$$

but where the deleted Čech differential only omits the *first* index (but includes the $(p+1)$ th).

A brief interlude on bundles

A holomorphic vector bundle E on X is described exactly by its *transition maps* $g_{\alpha\beta} \in \mathrm{GL}(n, \mathbb{C})$, which describe the change in trivialisation from over U_β to over U_α .

A brief interlude on bundles

A holomorphic vector bundle E on X is described exactly by its *transition maps* $g_{\alpha\beta} \in \mathrm{GL}(n, \mathbb{C})$, which describe the change in trivialisation from over U_β to over U_α .

These transition maps satisfy two conditions:

A brief interlude on bundles

A holomorphic vector bundle E on X is described exactly by its *transition maps* $g_{\alpha\beta} \in \mathrm{GL}(n, \mathbb{C})$, which describe the change in trivialisation from over U_β to over U_α .

These transition maps satisfy two conditions:

1. $g_{\alpha\beta}g_{\beta\gamma} = g_{\alpha\gamma}$ (the cocycle condition); and

A brief interlude on bundles

A holomorphic vector bundle E on X is described exactly by its *transition maps* $g_{\alpha\beta} \in \mathrm{GL}(n, \mathbb{C})$, which describe the change in trivialisation from over U_β to over U_α .

These transition maps satisfy two conditions:

1. $g_{\alpha\beta}g_{\beta\gamma} = g_{\alpha\gamma}$ (the *cocycle condition*); and
2. $g_{\alpha\alpha} = \mathrm{id}$ (the *invertibility condition*).

A brief interlude on bundles

A holomorphic vector bundle E on X is described exactly by its *transition maps* $g_{\alpha\beta} \in \mathrm{GL}(n, \mathbb{C})$, which describe the change in trivialisation from over U_β to over U_α .

These transition maps satisfy two conditions:

1. $g_{\alpha\beta}g_{\beta\gamma} = g_{\alpha\gamma}$ (the *cocycle condition*); and
2. $g_{\alpha\alpha} = \mathrm{id}$ (the *invertibility condition*).

Note that these are maps from $E|_{U_{\alpha p}}$ to $E|_{U_{\alpha_0}}$ in the specific case where $p = 1$.

Rewriting the cocycle condition

Thinking of $g_{\alpha\beta}$ as an element of $\mathcal{C}^1(\mathcal{U}, E)$, we see that

$$\begin{aligned}(\hat{\delta}g)_{\alpha\beta\gamma} &= -g_{\alpha\gamma} \\ (g \cdot g)_{\alpha\beta\gamma} &= g_{\alpha\beta}g_{\beta\gamma}.\end{aligned}$$

Rewriting the cocycle condition

Thinking of $g_{\alpha\beta}$ as an element of $\mathcal{C}^1(\mathcal{U}, E)$, we see that

$$\begin{aligned}(\hat{\delta}g)_{\alpha\beta\gamma} &= -g_{\alpha\gamma} \\ (g \cdot g)_{\alpha\beta\gamma} &= g_{\alpha\beta}g_{\beta\gamma}.\end{aligned}$$

This means that we can rewrite the cocycle condition as

$$\hat{\delta}g + g \cdot g = 0,$$

which looks like the *Maurer-Cartan equation* (an observation to which we will later return).

Twisting cochains

Definition (Twisting cochains)

A (holomorphic) twisting cochain over V is a formal sum

$$a = \bigoplus_{k \in \mathbb{N}} a^{k, 1-k}$$

where $a^{k, 1-k} \in \hat{\mathcal{C}}^k(\mathcal{U}, \text{End}^{1-k}(V))$ such that

1. $\hat{\delta}a + a \cdot a = 0$; and
2. $a_{\alpha\alpha}^{1,0} = \text{id}$.

Twisting cochains

Definition (Twisting cochains)

A (holomorphic) twisting cochain over V is a formal sum

$$a = \bigoplus_{k \in \mathbb{N}} a^{k, 1-k}$$

where $a^{k, 1-k} \in \hat{\mathcal{C}}^k(\mathcal{U}, \text{End}^{1-k}(V))$ such that

1. $\hat{\delta}a + a \cdot a = 0$; and
2. $a_{\alpha\alpha}^{1,0} = \text{id}$.

The invertibility condition “should” really be weakened by asking only that $a_{\alpha\alpha}^{1,0}$ be *homotopic* to the identity.

Twisting cochains (cont.)

Warning

The multiplication is **not** simply component-wise: it is given by taking all possible combinations, i.e.

$$(a \cdot b)^{p,s} = \bigoplus_{\substack{q+q'=p \\ t+t'=s}} a^{q,t} \cdot b^{q',t'}.$$

Twisting cochains (cont.)

Warning

The multiplication is **not** simply component-wise: it is given by taking all possible combinations, i.e.

$$(a \cdot b)^{p,s} = \bigoplus_{\substack{q+q'=p \\ t+t'=s}} a^{q,t} \cdot b^{q',t'}.$$

- It might be the case that all but finitely many of the $a^{k,1-k}$ are zero, but **never** $a^{1,0}$, since it has to be the identity on $\alpha\alpha$.

Twisting cochains (cont.)

Warning

The multiplication is **not** simply component-wise: it is given by taking all possible combinations, i.e.

$$(a \cdot b)^{p,s} = \bigoplus_{\substack{q+q'=p \\ t+t'=s}} a^{q,t} \cdot b^{q',t'}.$$

- It might be the case that all but finitely many of the $a^{k,1-k}$ are zero, but **never** $a^{1,0}$, since it has to be the identity on $\alpha\alpha$.
- If V has a differential then a is an element of total degree 1.

Twisting cochains (cont.)

Warning

The multiplication is **not** simply component-wise: it is given by taking all possible combinations, i.e.

$$(a \cdot b)^{p,s} = \bigoplus_{\substack{q+q'=p \\ t+t'=s}} a^{q,t} \cdot b^{q',t'}.$$

- It might be the case that all but finitely many of the $a^{k,1-k}$ are zero, but **never** $a^{1,0}$, since it has to be the identity on $\alpha\alpha$.
- If V has a differential then a is an element of total degree 1.
- We haven't said *when* twisting cochains exist, but under pretty mild assumptions they always do (by an inductive construction).

Unpacking the definition

$(k = 0) \rightsquigarrow a_{\alpha}^{0,1} \cdot a_{\alpha}^{0,1} = 0$, which tells us that $a_{\alpha}^{0,1}$ is a differential on V_{α}^{\bullet} .

Unpacking the definition

$(k = 0) \rightsquigarrow a_{\alpha}^{0,1} \cdot a_{\alpha}^{0,1} = 0$, which tells us that $a_{\alpha}^{0,1}$ is a *differential on V_{α}^{\bullet}* .

$(k = 1) \rightsquigarrow a_{\alpha}^{0,1} \cdot a_{\alpha\beta}^{1,0} = a_{\alpha\beta}^{1,0} \cdot a_{\beta}^{0,1}$, which tells us that we have a *chain map of chain complexes*

$$a_{\alpha\beta}^{1,0}: (V_{\beta}^{\bullet}|U_{\alpha\beta}, a_{\beta}^{0,1}) \rightarrow (V_{\alpha}^{\bullet}|U_{\alpha\beta}, a_{\alpha}^{0,1})$$

Unpacking the definition

$(k = 0) \rightsquigarrow a_{\alpha}^{0,1} \cdot a_{\alpha}^{0,1} = 0$, which tells us that $a_{\alpha}^{0,1}$ is a *differential on V_{α}^{\bullet}* .

$(k = 1) \rightsquigarrow a_{\alpha}^{0,1} \cdot a_{\alpha\beta}^{1,0} = a_{\alpha\beta}^{1,0} \cdot a_{\beta}^{0,1}$, which tells us that we have a *chain map of chain complexes*

$$a_{\alpha\beta}^{1,0}: (V_{\beta}^{\bullet} | U_{\alpha\beta}, a_{\beta}^{0,1}) \rightarrow (V_{\alpha}^{\bullet} | U_{\alpha\beta}, a_{\alpha}^{0,1})$$

$(k = 2) \rightsquigarrow -a_{\alpha\gamma}^{1,0} + a_{\alpha\beta}^{1,0} \cdot a_{\beta\gamma}^{1,0} = a_{\alpha}^{0,1} \cdot a_{\alpha\beta\gamma}^{2,-1} + a_{\alpha\beta\gamma}^{2,-1} \cdot a_{\gamma}^{0,1}$, which says that $a_{\alpha\beta\gamma}^{2,-1}$ witnesses a *chain homotopy* between $a_{\alpha\gamma}^{1,0}$ and $a_{\alpha\beta}^{1,0} \cdot a_{\beta\gamma}^{1,0}$. On $\alpha\beta\alpha$ and $\beta\alpha\beta$ this tells us that $a_{\alpha\beta}^{1,0}$ and $a_{\beta\alpha}^{1,0}$ are *chain homotopic inverses*, i.e. *quasi-isomorphism*.

Unpacking the definition (cont.)

$(k \geq 3) \rightsquigarrow$ some sort of 'higher homotopic gluings', whatever this might mean.

Unpacking the definition (cont.)

$(k \geq 3) \rightsquigarrow$ some sort of ‘higher homotopic gluings’, whatever this might mean.

This is one of the things that we want to formalise!

Unpacking the definition (cont.)

$(k \geq 3) \rightsquigarrow$ some sort of ‘higher homotopic gluings’, whatever this might mean.

This is one of the things that we want to formalise!

Extra-curricular

By taking (internal) homology we obtain something strict: a complex of *coherent sheaves* $H^\bullet(a)$. This is because quasi-isomorphisms become strict isomorphisms in homology.

We can use this fact to construct twisting cochains that resolve coherent sheaves by taking *local* resolutions by vector bundles.

The total differential

Lemma

For any $a \in \text{Tot}^1 \hat{\mathcal{C}}^\bullet(\mathcal{U}, \text{End}^\circ(V))$, the map

$$\begin{aligned} D_a : \text{Tot}^r \hat{\mathcal{C}}^\bullet(\mathcal{U}, V^\circ) &\rightarrow \text{Tot}^{r+1} \hat{\mathcal{C}}^\bullet(\mathcal{U}, V^\circ) \\ c &\mapsto \hat{\delta}c + c \cdot a \end{aligned}$$

defines a differential (i.e. squares to zero) if and only if a is a twisting cochain.

Proof.

(Tedious) definition chasing. □

The total differential (cont.)

We can actually define twisting cochains in a different way using this lemma (but we won't do so today).

The total differential (cont.)

We can actually define twisting cochains in a different way using this lemma (but we won't do so today).

But this approach lets us think of a twisting cochain as a *first-order perturbation of the deleted Čech differential*.

Examples

Example

Look at the most trivial example: let V be an *ungraded* vector bundle, and $a = a^{0,1} + a^{1,0}$, where $a_{\alpha}^{0,1} = \text{id}_{V_{\alpha}}$, and the $a^{1,0}$ are the transition maps. Then

$$(D_a c)_{\alpha_0 \dots \alpha_{p+1}} = a_{\alpha_0 \alpha_1}^{1,0} c_{\alpha_1 \dots \alpha_{p+1}} + \sum_{i=1}^{p+1} (-1)^i c_{\alpha_0 \dots \hat{\alpha}_i \dots \alpha_{p+1}}.$$

Examples

Example

Look at the most trivial example: let V be an *ungraded* vector bundle, and $a = a^{0,1} + a^{1,0}$, where $a_{\alpha}^{0,1} = \text{id}_{V_{\alpha}}$, and the $a^{1,0}$ are the transition maps. Then

$$(D_a c)_{\alpha_0 \dots \alpha_{p+1}} = a_{\alpha_0 \alpha_1}^{1,0} c_{\alpha_1 \dots \alpha_{p+1}} + \sum_{i=1}^{p+1} (-1)^i c_{\alpha_0 \dots \hat{\alpha}_i \dots \alpha_{p+1}}.$$

Note that we couldn't use the full Čech differential on $\hat{\mathcal{C}}^{\bullet}(\mathcal{U}, V^{\circ})$ because everything has to lie over U_{α_0} , but this total differential solves that problem — recall that $a_{\alpha_0 \alpha_1}^{1,0}$ is a (quasi-)isomorphism.

Examples

Example

Look at the most trivial example: let V be an *ungraded* vector bundle, and $a = a^{0,1} + a^{1,0}$, where $a_{\alpha}^{0,1} = \text{id}_{V_{\alpha}}$, and the $a^{1,0}$ are the transition maps. Then

$$(D_a c)_{\alpha_0 \dots \alpha_{p+1}} = a_{\alpha_0 \alpha_1}^{1,0} c_{\alpha_1 \dots \alpha_{p+1}} + \sum_{i=1}^{p+1} (-1)^i c_{\alpha_0 \dots \hat{\alpha}_i \dots \alpha_{p+1}}.$$

Note that we couldn't use the full Čech differential on $\hat{\mathcal{C}}^{\bullet}(U, V^{\circ})$ because everything has to lie over U_{α_0} , but this total differential solves that problem — recall that $a_{\alpha_0 \alpha_1}^{1,0}$ is a (quasi-)isomorphism.

A spectral-sequence argument shows that, in fact, D_a here really is 'the same as' the full Čech differential.

Examples (cont.)

Example

Now look at a slightly-less trivial example: let V^\bullet consist of complexes $(V_\alpha^\bullet, d_\alpha)$ of vector bundles, and $a = a^{0,1} + a^{1,0}$, where $a_\alpha^{0,1} = d_\alpha$, and the $a^{1,0}$ are the transition maps. Then

$$\begin{aligned}(D_a c)_{\alpha_0 \dots \alpha_{p+1}} &= (-1)^p a_{\alpha_0}^{0,1} c_{\alpha_0 \dots \alpha_p} + a_{\alpha_0 \alpha_1}^{1,0} c_{\alpha_1 \dots \alpha_{p+1}} \\ &\quad + \sum_{i=1}^{p+1} (-1)^i c_{\alpha_0 \dots \hat{\alpha}_i \dots \alpha_{p+1}}.\end{aligned}$$

Examples (cont.)

Example

Now look at a slightly-less trivial example: let V^\bullet consist of complexes $(V_\alpha^\bullet, d_\alpha)$ of vector bundles, and $a = a^{0,1} + a^{1,0}$, where $a_\alpha^{0,1} = d_\alpha$, and the $a^{1,0}$ are the transition maps. Then

$$\begin{aligned}(D_a c)_{\alpha_0 \dots \alpha_{p+1}} &= (-1)^p a_{\alpha_0}^{0,1} c_{\alpha_0 \dots \alpha_p} + a_{\alpha_0 \alpha_1}^{1,0} c_{\alpha_1 \dots \alpha_{p+1}} \\ &\quad + \sum_{i=1}^{p+1} (-1)^i c_{\alpha_0 \dots \hat{\alpha}_i \dots \alpha_{p+1}}.\end{aligned}$$

Identifying the second and third terms with the full Čech differential, as above, gives the usual total differential of the Čech bicomplex:

$$D_a = \check{\delta} \pm d_V.$$

Twisted complexes (BK)

Other fun things
