

# class09

Thrisha Praveen

Reading in the CSV and looking at it closer:

```
pdb_stats <- read.csv("/Users/thrishapraveen/Downloads/BIMM143/class09/Data Export Summary.csv")
pdb_stats
```

	Molecular.Type	X.ray	EM	NMR	Multiple.methods	Neutron	Other
1	Protein (only)	169,563	16,774	12,578	208	81	32
2	Protein/Oligosaccharide	9,939	2,839	34	8	2	0
3	Protein/NA	8,801	5,062	286	7	0	0
4	Nucleic acid (only)	2,890	151	1,521	14	3	1
5	Other	170	10	33	0	0	0
6	Oligosaccharide (only)	11	0	6	1	0	4
	Total						
1		199,236					
2		12,822					
3		14,156					
4		4,580					
5		213					
6		22					

```
str(pdb_stats)
```

```
'data.frame': 6 obs. of 8 variables:
 $ Molecular.Type : chr "Protein (only)" "Protein/Oligosaccharide" "Protein/NA" "Nucleic acid (only)" "Other" "Oligosaccharide (only)"
 $ X.ray          : chr "169,563" "9,939" "8,801" "2,890" ...
 $ EM            : chr "16,774" "2,839" "5,062" "151" ...
 $ NMR           : chr "12,578" "34" "286" "1,521" ...
 $ Multiple.methods: int 208 8 7 14 0 1
 $ Neutron       : int 81 2 0 3 0 0
 $ Other         : int 32 0 0 1 0 4
 $ Total         : chr "199,236" "12,822" "14,156" "4,580" ...
```

Q1: What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.  $\text{sum(X.ray)} + \text{sum(EM)} = 191,374 + 24,836 = 216,210$  / total =  $216,210 / 231,029 = 0.9359 = 93.59\%$

Q2: What proportion of structures in the PDB are protein? If we're looking at protein(only), then it's  $199,236 / 231,029 = 0.862$  If we're looking at all the protein columns, then it's  $(199,236 + 12,822 + 14,156) / 231,029 = 0.979$

Q3: Type HIV in the PDB website search box on the home page and determine how many HIV-1 protease structures are in the current PDB? 231,029 HIV-1 protease structures

## 2. Visualizing the HIV-1 protease structure

Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure? In PDB files, hydrogen atoms aren't explicitly stored. Thus, for water molecules, only the oxygen atom is stored/visualized.

Q5: There is a critical "conserved" water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have? I didn't understand how to tell what is a conserved water molecule, but the one that seems to play a role in the binding site is HOH 308.

Q6: Generate and save a figure clearly showing the two distinct chains of HIV-protease along with the ligand. You might also consider showing the catalytic residues ASP 25 in each chain and the critical water (we recommend "Ball & Stick" for these side-chains). Add this figure to your Quarto document.


```
library(knitr)
include_graphics("/Users/thrishapraveen/Downloads/BIMM143/class09/1HSG (1).png")
```



```
include_graphics("/Users/thrishapraveen/Downloads/BIMM143/class09/1HSG(1.5).png")
```



```
include_graphics("/Users/thrishapraveen/Downloads/BIMM143/class09/1HSG (2).png")
```



### 3. Introduction to Bio3D in R

```
library(bio3d)  
pdb <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
pdb
```

```
Call: read.pdb(file = "1hsg")
```

```
Total Models#: 1
```

```
Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)
```

```
Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 172 (residues: 128)
```

```
Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
```

```
Protein sequence:
```

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
VNIIGRNLLTQIGCTLNF
```

```
+ attr: atom, xyz, seqres, helix, sheet,
      calpha, remark, call
```

```
Q7: How many amino acid residues are there in this pdb object? 198
```

```
Q8: Name one of the two non-protein residues? HOH
```

```
Q9: How many protein chains are in this structure? 2
```

```
attributes(pdb)
```

```
$names
```

```
[1] "atom" "xyz" "seqres" "helix" "sheet" "calpha" "remark" "call"
```

```
$class
```

```
[1] "pdb" "sse"
```

```
head(pdb$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40

```

5 ATOM      5      CB <NA>   PRO      A      1      <NA> 30.508 37.541 6.342 1 37.87
6 ATOM      6      CG <NA>   PRO      A      1      <NA> 29.296 37.591 7.162 1 38.40
  segid elesy charge
1  <NA>      N  <NA>
2  <NA>      C  <NA>
3  <NA>      C  <NA>
4  <NA>      O  <NA>
5  <NA>      C  <NA>
6  <NA>      C  <NA>

```

Predicting functional motions of a single structure

```
adk <- read.pdb("6s36")
```

Note: Accessing on-line PDB file

PDB has ALT records, taking A only, rm.alt=TRUE

```
adk
```

Call: read.pdb(file = "6s36")

Total Models#: 1

Total Atoms#: 1898, XYZs#: 5694 Chains#: 1 (values: A)

Protein Atoms#: 1654 (residues/Calpha atoms#: 214)

Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)

Non-protein/nucleic Atoms#: 244 (residues: 244)

Non-protein/nucleic resid values: [ CL (3), HOH (238), MG (2), NA (1) ]

Protein sequence:

```

MRIILLGAPGAGKGTQAQFIMEKYGIPQISTGDMRLRAAVKSGSELGKQAKDIMDAGKLV
TDELVIALVKERIAQEDCRNGFLLDGFPRTPQADAMKEAGINVDYVLEFDVPDELIVDKI
VGRRVHAPSGRVYHVKFNPPKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQMTAPLIG
YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG

```

```


+ attr: atom, xyz, seqres, helix, sheet,
      calpha, remark, call

```

```
# Perform flexibility prediction
m <- nma(adk)
```

```
Building Hessian...      Done in 0.014 seconds.
Diagonalizing Hessian... Done in 0.278 seconds.
```

```
plot(m)
```



```
mktrj(m, file="adk_m7.pdb")
```

4. Comparative structure analysis of Adenylate Kinase >Q10. Which of the packages above is found only on BioConductor and not CRAN? msa

Q11. Which of the above packages is not found on BioConductor or CRAN?:  
bio3d-view

Q12. True or False? Functions from the devtools package can be used to install packages from GitHub and BitBucket? TRUE

Search and retrieve ADK structures



```
library(bio3d)
aa <- get.seq("lake_A")
```

Warning in get.seq("lake\_A"): Removing existing file: seqs.fasta

Fetching... Please wait. Done.

```
aa
```

```

      1      .      .      .      .      .      60
pdb|1AKE|A  MRIILLGAPGAGKGTQAQFIMEKYGIPQISTGDMRLAAVKSSELGKQAKDIMDAGKLV
      1      .      .      .      .      .      60

      61      .      .      .      .      .      120
pdb|1AKE|A  DELVIALVKERIAQEDCRNGFLLDGFPRTIPQADAMKEAGINVVDYVLEFDVPDELIVDRI
      61      .      .      .      .      .      120

     121      .      .      .      .      .      180
pdb|1AKE|A  VGRRVHAPSGRVYHVKNPPKVEGKDDVTGEELTRKDDQEETVRKRLVEYHQMTPALIG
     121      .      .      .      .      .      180

     181      .      .      .      214
pdb|1AKE|A  YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG
     181      .      .      .      214
```

Call:

```
read.fasta(file = outfile)
```

Class:

```
fasta
```

Alignment dimensions:

```
1 sequence rows; 214 position columns (214 non-gap, 0 gap)
```

```
+ attr: id, ali, call
```

Q13. How many amino acids are in this sequence, i.e. how long is this sequence?  
214

```
# Blast or hmmer search
#b <- blast.pdb(aa)
```

```
# Plot a summary of search results
#hits <- plot(b)
```

```
# List out some 'top hits'
#head(hits$ pdb.id)
```

```
hits <- NULL
hits$ pdb.id <- c('1AKE_A','6S36_A','6RZE_A','3HPR_A','1E4V_A','5EJE_A','1E4Y_A','3X2S_A','6HPR_A')
```

```
# Download related PDB files
files <- get.pdb(hits$ pdb.id, path="pdb", split=TRUE, gzip=TRUE)
```

```
Warning in get.pdb(hits$ pdb.id, path = "pdb", split = TRUE, gzip = TRUE):
pdb/1AKE.pdb.gz exists. Skipping download
```

```
Warning in get.pdb(hits$ pdb.id, path = "pdb", split = TRUE, gzip = TRUE):
pdb/6S36.pdb.gz exists. Skipping download
```

```
Warning in get.pdb(hits$ pdb.id, path = "pdb", split = TRUE, gzip = TRUE):
pdb/6RZE.pdb.gz exists. Skipping download
```

```
Warning in get.pdb(hits$ pdb.id, path = "pdb", split = TRUE, gzip = TRUE):
pdb/3HPR.pdb.gz exists. Skipping download
```

```
Warning in get.pdb(hits$ pdb.id, path = "pdb", split = TRUE, gzip = TRUE):
pdb/1E4V.pdb.gz exists. Skipping download
```

```
Warning in get.pdb(hits$ pdb.id, path = "pdb", split = TRUE, gzip = TRUE):
pdb/5EJE.pdb.gz exists. Skipping download
```

```
Warning in get.pdb(hits$ pdb.id, path = "pdb", split = TRUE, gzip = TRUE):
pdb/1E4Y.pdb.gz exists. Skipping download
```

```
Warning in get.pdb(hits$ pdb.id, path = "pdb", split = TRUE, gzip = TRUE):
pdb/3X2S.pdb.gz exists. Skipping download
```


Warning in get.pdb(hits\$pdb.id, path = "pdbs", split = TRUE, gzip = TRUE):  
pdbs/6HAP.pdb.gz exists. Skipping download

Warning in get.pdb(hits\$pdb.id, path = "pdbs", split = TRUE, gzip = TRUE):  
pdbs/6HAM.pdb.gz exists. Skipping download

Warning in get.pdb(hits\$pdb.id, path = "pdbs", split = TRUE, gzip = TRUE):  
pdbs/4K46.pdb.gz exists. Skipping download

Warning in get.pdb(hits\$pdb.id, path = "pdbs", split = TRUE, gzip = TRUE):  
pdbs/3GMT.pdb.gz exists. Skipping download

Warning in get.pdb(hits\$pdb.id, path = "pdbs", split = TRUE, gzip = TRUE):  
pdbs/4PZL.pdb.gz exists. Skipping download



```
|=====| 92%
|
|=====| 100%
```

Align and superpose structures

```
# Align related PDBs
pddb <- pddbaln(files, fit = TRUE, exefile="msa")
```

Reading PDB files:

```
pddb/split_chain/1AKE_A.pdb
pddb/split_chain/6S36_A.pdb
pddb/split_chain/6RZE_A.pdb
pddb/split_chain/3HPR_A.pdb
pddb/split_chain/1E4V_A.pdb
pddb/split_chain/5EJE_A.pdb
pddb/split_chain/1E4Y_A.pdb
pddb/split_chain/3X2S_A.pdb
pddb/split_chain/6HAP_A.pdb
pddb/split_chain/6HAM_A.pdb
pddb/split_chain/4K46_A.pdb
pddb/split_chain/3GMT_A.pdb
pddb/split_chain/4PZL_A.pdb
  PDB has ALT records, taking A only, rm.alt=TRUE
.   PDB has ALT records, taking A only, rm.alt=TRUE
.   PDB has ALT records, taking A only, rm.alt=TRUE
.   PDB has ALT records, taking A only, rm.alt=TRUE
..  PDB has ALT records, taking A only, rm.alt=TRUE
.... PDB has ALT records, taking A only, rm.alt=TRUE
.   PDB has ALT records, taking A only, rm.alt=TRUE
...
```

Extracting sequences

```
pddb/seq: 1   name: pddb/split_chain/1AKE_A.pdb
  PDB has ALT records, taking A only, rm.alt=TRUE
pddb/seq: 2   name: pddb/split_chain/6S36_A.pdb
  PDB has ALT records, taking A only, rm.alt=TRUE
pddb/seq: 3   name: pddb/split_chain/6RZE_A.pdb
  PDB has ALT records, taking A only, rm.alt=TRUE
pddb/seq: 4   name: pddb/split_chain/3HPR_A.pdb
  PDB has ALT records, taking A only, rm.alt=TRUE
```

```

pdb/seq: 5   name: pdbs/split_chain/1E4V_A.pdb
pdb/seq: 6   name: pdbs/split_chain/5EJE_A.pdb
           PDB has ALT records, taking A only, rm.alt=TRUE
pdb/seq: 7   name: pdbs/split_chain/1E4Y_A.pdb
pdb/seq: 8   name: pdbs/split_chain/3X2S_A.pdb
pdb/seq: 9   name: pdbs/split_chain/6HAP_A.pdb
pdb/seq: 10  name: pdbs/split_chain/6HAM_A.pdb
           PDB has ALT records, taking A only, rm.alt=TRUE
pdb/seq: 11  name: pdbs/split_chain/4K46_A.pdb
           PDB has ALT records, taking A only, rm.alt=TRUE
pdb/seq: 12  name: pdbs/split_chain/3GMT_A.pdb
pdb/seq: 13  name: pdbs/split_chain/4PZL_A.pdb


```

```

# Vector containing PDB codes for figure axis
ids <- basename.pdb(pdb$id)

# Draw schematic alignment
#plot(pdb, labels=ids)
library(knitr)
include_graphics("/Users/thrishapraveen/Downloads/BIMM143/class09/seq_align.png")

```



My Sequence Alignment looks different than the one on the Class 9 page. Some of the sequences there are missing here. This image wasn't letting me render, so I attached a screenshot instead.

Annotate collected PDB structures

```
anno <- pdb.annotate(ids)
unique(anno$source)
```

```
[1] "Escherichia coli"
[2] "Escherichia coli K-12"
[3] "Escherichia coli O139:H28 str. E24377A"
[4] "Escherichia coli str. K-12 substr. MDS42"
[5] "Photobacterium profundum"
[6] "Burkholderia pseudomallei 1710b"
[7] "Francisella tularensis subsp. tularensis SCHU S4"
```

**[6] “Vibrio cholerae O1 biovar El Tor str. N16961” is missing and I’m not sure why**

```
anno
```

	structureId	chainId	macromoleculeType	chainLength	experimentalTechnique
1AKE_A	1AKE	A	Protein	214	X-ray
6S36_A	6S36	A	Protein	214	X-ray
6RZE_A	6RZE	A	Protein	214	X-ray
3HPR_A	3HPR	A	Protein	214	X-ray
1E4V_A	1E4V	A	Protein	214	X-ray
5EJE_A	5EJE	A	Protein	214	X-ray
1E4Y_A	1E4Y	A	Protein	214	X-ray
3X2S_A	3X2S	A	Protein	214	X-ray
6HAP_A	6HAP	A	Protein	214	X-ray
6HAM_A	6HAM	A	Protein	214	X-ray
4K46_A	4K46	A	Protein	214	X-ray
3GMT_A	3GMT	A	Protein	230	X-ray
4PZL_A	4PZL	A	Protein	242	X-ray
	resolution	scopDomain	pfam		
1AKE_A	2.00	Adenylate kinase	Adenylate kinase, active site lid (ADK_lid)		
6S36_A	1.60	<NA>	Adenylate kinase (ADK)		
6RZE_A	1.69	<NA>	Adenylate kinase, active site lid (ADK_lid)		

3HPR_A	2.00	<NA>	Adenylate kinase, active site lid (ADK_lid)
1E4V_A	1.85	Adenylate kinase	Adenylate kinase (ADK)
5EJE_A	1.90	<NA>	Adenylate kinase, active site lid (ADK_lid)
1E4Y_A	1.85	Adenylate kinase	Adenylate kinase, active site lid (ADK_lid)
3X2S_A	2.80	<NA>	Adenylate kinase (ADK)
6HAP_A	2.70	<NA>	Adenylate kinase (ADK)
6HAM_A	2.55	<NA>	Adenylate kinase, active site lid (ADK_lid)
4K46_A	2.01	<NA>	Adenylate kinase (ADK)
3GMT_A	2.10	<NA>	Adenylate kinase (ADK)
4PZL_A	2.10	<NA>	Adenylate kinase (ADK)

#### ligandId

1AKE_A	AP5
6S36_A	CL (3),NA,MG (2)
6RZE_A	NA (3),CL (2)
3HPR_A	AP5
1E4V_A	AP5
5EJE_A	AP5,CO
1E4Y_A	AP5
3X2S_A	JPY (2),AP5,MG
6HAP_A	AP5
6HAM_A	AP5
4K46_A	ADP,AMP,PO4
3GMT_A	SO4 (2)
4PZL_A	CA,FMT,GOL

#### ligandName

1AKE_A	BIS(ADENOSINE)-5'-PENTAPHOSPHATE
6S36_A	CHLORIDE ION (3),SODIUM ION,MAGNESIUM ION (2)
6RZE_A	SODIUM ION (3),CHLORIDE ION (2)
3HPR_A	BIS(ADENOSINE)-5'-PENTAPHOSPHATE
1E4V_A	BIS(ADENOSINE)-5'-PENTAPHOSPHATE
5EJE_A	BIS(ADENOSINE)-5'-PENTAPHOSPHATE,COBALT (II) ION
1E4Y_A	BIS(ADENOSINE)-5'-PENTAPHOSPHATE
3X2S_A	N-(pyren-1-ylmethyl)acetamide (2),BIS(ADENOSINE)-5'-PENTAPHOSPHATE,MAGNESIUM ION
6HAP_A	BIS(ADENOSINE)-5'-PENTAPHOSPHATE
6HAM_A	BIS(ADENOSINE)-5'-PENTAPHOSPHATE
4K46_A	ADENOSINE-5'-DIPHOSPHATE,ADENOSINE MONOPHOSPHATE,PHOSPHATE ION
3GMT_A	SULFATE ION (2)
4PZL_A	CALCIUM ION,FORMIC ACID,GLYCEROL

#### source

1AKE_A	Escherichia coli
6S36_A	Escherichia coli
6RZE_A	Escherichia coli
3HPR_A	Escherichia coli K-12

1E4V_A	Escherichia coli
5EJE_A	Escherichia coli 0139:H28 str. E24377A
1E4Y_A	Escherichia coli
3X2S_A	Escherichia coli str. K-12 substr. MDS42
6HAP_A	Escherichia coli 0139:H28 str. E24377A
6HAM_A	Escherichia coli K-12
4K46_A	Photobacterium profundum
3GMT_A	Burkholderia pseudomallei 1710b
4PZL_A	Francisella tularensis subsp. tularensis SCHU S4

1AKE_A	STRUCTURE OF THE COMPLEX BETWEEN ADENYLATE KINASE FROM ESCHERICHIA COLI AND THE INHIBIT
6S36_A	
6RZE_A	
3HPR_A	
1E4V_A	
5EJE_A	
1E4Y_A	
3X2S_A	
6HAP_A	
6HAM_A	
4K46_A	
3GMT_A	
4PZL_A	

		citation	rObserved	rFree
1AKE_A	Muller, C.W., et al. J Mol Biol (1992)	0.19600	NA	
6S36_A	Rogne, P., et al. Biochemistry (2019)	0.16320	0.23560	
6RZE_A	Rogne, P., et al. Biochemistry (2019)	0.18650	0.23500	
3HPR_A	Schrank, T.P., et al. Proc Natl Acad Sci U S A (2009)	0.21000	0.24320	
1E4V_A	Muller, C.W., et al. Proteins (1993)	0.19600	NA	
5EJE_A	Kovermann, M., et al. Proc Natl Acad Sci U S A (2017)	0.18890	0.23580	
1E4Y_A	Muller, C.W., et al. Proteins (1993)	0.17800	NA	
3X2S_A	Fujii, A., et al. Bioconj Chem (2015)	0.20700	0.25600	
6HAP_A	Kantaev, R., et al. J Phys Chem B (2018)	0.22630	0.27760	
6HAM_A	Kantaev, R., et al. J Phys Chem B (2018)	0.20511	0.24325	
4K46_A	Cho, Y.-J., et al. To be published	0.17000	0.22290	
3GMT_A	Buchko, G.W., et al. Biochem Biophys Res Commun (2010)	0.23800	0.29500	
4PZL_A	Tan, K., et al. To be published	0.19360	0.23680	

	rWork	spaceGroup
1AKE_A	0.19600	P 21 2 21
6S36_A	0.15940	C 1 2 1
6RZE_A	0.18190	C 1 2 1
3HPR_A	0.20620	P 21 21 2
1E4V_A	0.19600	P 21 2 21



```

5EJE_A 0.18630 P 21 2 21
1E4Y_A 0.17800 P 1 21 1
3X2S_A 0.20700 P 21 21 21
6HAP_A 0.22370 I 2 2 2
6HAM_A 0.20311 P 43
4K46_A 0.16730 P 21 21 21
3GMT_A 0.23500 P 1 21 1
4PZL_A 0.19130 P 32

```

Principal component analysis

```

# Perform PCA
pc.xray <- pca(pdbx)
plot(pc.xray)

```




Figure 9: Results of PCA on Adenylate kinase X-ray structures. Each dot represents one PDB structure.

The values are a little off compared to the Class 9 page.

```

# Calculate RMSD
rd <- rmsd(pdbx)

```

Warning in rmsd(pdb): No indices provided, using the 204 non NA positions

```
# Structure-based clustering
hc.rd <- hclust(dist(rd))
grps.rd <- cutree(hc.rd, k=3)

plot(pc.xray, 1:2, col="grey50", bg=grps.rd, pch=21, cex=1)
```




Figure 10: Projection of Adenylate kinase X-ray structures. Each dot represents one PDB structure.