

NEUTRAL ISM, LYMAN-ALPHA AND LYMAN-CONTINUUM IN NEARBY STARBURST HARO 11¹

T. EMIL RIVERA-THORSEN^{2, 3} AND GÖRAN ÖSTLIN^{2, 3}

¹Based on observations Cosmic Origins Spectrograph on the Hubble Space Telescope, program XXXXXX, PI YYYYYYYYYY

²Department of Astronomy, Oskar Klein Centre, Stockholm University, AlbaNova University Centre, SE-106 91 Stockholm, Sweden

³Oscar Klein Centre for Cosmoparticle Physics, Department of Astronomy, Stockholm University, Stockholm, Sweden

ABSTRACT

Something something something.

1. INTRODUCTION AND OBSERVATIONS

About Haro 11: Some of the same stuff as in Rivera-Thorsen et al. (in prep.) could be mentioned, with a different emphasis. Star formation rate should be mentioned from Adamo et al. (2010). Population from same and from Micheva et al. (2010). Knots named by Vader et al. (1993), and also mentioned by Kunth et al. (2003). It's an LBA (Hoopes et al. 2007), and an LAE analog (Hayes et al. 2007; Leitet et al. 2011). Knot C a strong Ly α emitter (Hayes et al. 2007), and also a LyC emleaker

[Bergvall et al. (2006); Leitet2011]. Kinematics, it's a merger (Östlin et al. 1999, 2001, 2015). More weight on the Lyman- α escape thing, and LyC. Mention Ly α escape and the history here, too? References:

2. ANALYSIS

Bla bla bla

3. DISCUSSION AND CONCLUSIONS

Bla bla bla

REFERENCES

- Adamo, A., Östlin, G., Zackrisson, E., et al. 2010, MNRAS, 407, 870
- Bergvall, N., Zackrisson, E., Andersson, B.-G., et al. 2006, A&A, 448, 513
- Hayes, M., Östlin, G., Atek, H., et al. 2007, MNRAS, 382, 1465
- Hoopes, C. G., Heckman, T. M., Salim, S., et al. 2007, ApJS, 173, 441
- Kunth, D., Leitherer, C., Mas-Hesse, J. M., Östlin, G., & Petrosian, A. 2003, ApJ, 597, 263
- Leitet, E., Bergvall, N., Piskunov, N., & Andersson, B.-G. 2011, A&A, 532, A107
- Micheva, G., Zackrisson, E., Östlin, G., Bergvall, N., & Pursimo, T. 2010, MNRAS, 405, 1203
- Östlin, G., Amram, P., Bergvall, N., et al. 2001, A&A, 374, 800
- Östlin, G., Amram, P., Masegosa, J., Bergvall, N., & Boulesteix, J. 1999, A&AS, 137, 419
- Östlin, G., Marquart, T., Cumming, R. J., et al. 2015, A&A, 583, A55
- Sandberg, A., Östlin, G., Hayes, M., et al. 2013, A&A, 552, A95
- Vader, J. P., Frogel, J. A., Terndrup, D. M., & Heisler, C. A. 1993, AJ, 106, 1743




Figure 1. Approximate coverage of the COS aperture. The aperture, which was centered on the supercluster in knot C, has a diameter of $2.5''$




Figure 2. Full spectrum in restframe wavelength of Haro 11 C. Redshift adopted from [Sandberg et al. \(2013\)](#). Orange lines denote Milky Way absorption, while blue lines denote absorption internal to the target.




Figure 3. The Si II (**upper**) and Si IV (**lower**) profiles included in this study.




Figure 4. **Upper panel:** Ly α profile of Haro 11 C, in approximate units of the surrounding continuum level. Full line is the measured values smoothed by a 5 px. flat kernel; surrounding shading encloses the $\pm 1\sigma$ error band. **Middle panel:** Black steps show the averaged, LIS line profile, smoothed by a 5px kernel. Surrounding gray shading denotes the $\pm 1\sigma$ confidence band. **Lower panel:** Same as middle panel, but for the Si IV transitions.