Proofs without words I

Exercises in METAPOST

Toby Thurston

March 2021 —



Contents

| Geometry and Algebra | 3 |
|---|----|
| Trigonometry, Calculus, & Analytic Geometry | 28 |

Geometry and Algebra

| The Pythagorean theorem I | 4 |
|--|----|
| The Pythagorean theorem II | 5 |
| The Pythagorean theorem III | 6 |
| The Pythagorean theorem IV | 7 |
| The Pythagorean theorem V \ldots | 8 |
| The Pythagorean theorem VI | 9 |
| A Pythagorean theorem: $aa' = bb' + cc'$ | 10 |
| The rolling circle squares itself | 11 |
| On trisecting an angle | 12 |
| Trisection in an infinite number of steps | 13 |
| Trisection of a line segment | 14 |
| The vertex angles of a star sum to 180° | 15 |
| Viviani's theorem I | 16 |
| Viviani's theorem II | 17 |
| A theorem about right angles | 18 |
| Area and the projection theorem of a right triangle | 19 |
| Chords and tangents of equal length | 20 |
| Completing the square | 21 |
| Algebraic areas I | 22 |
| Algebraic areas II | 23 |
| Sum of squares identity | 24 |
| Polygonal numbers | 25 |
| The volume of a frustrum of a square pyramid | 26 |
| The volume of a homisphere via Cavalieri's Principle | 27 |

The Pythagorean theorem I





— adapted from the Chou pei san ching

The Pythagorean theorem II





Behold!

— Bhāskara (12th century)

The Pythagorean theorem III



— based on Euclid's proof

The Pythagorean theorem IV



— H. E. Dudeney (1917)

The Pythagorean theorem \boldsymbol{V}



— James A. Garfield (1876)

The Pythagorean theorem VI



— Michael Hardy

A Pythagorean theorem: aa' = bb' + cc'





$$\frac{x}{b'} = \frac{b}{a} \implies \frac{x}{b} = \frac{b'}{a} \implies ax = bb';$$

$$\frac{y}{c'} = \frac{c}{a} \implies \frac{y}{c} = \frac{c'}{a} \implies ay = cc';$$

$$\therefore aa' = a(x + y) = bb' + cc'.$$

— Enzo R. Gentile

The rolling circle squares itself



— Thomas Elsner

On trisecting an angle



— Rufus Isaacs

Trisection in an infinite number of steps



 $\frac{1}{3} = \frac{1}{2} - \frac{1}{4} + \frac{1}{8} - \frac{1}{16} + \cdots$

— Eric Kincanon

Trisection of a line segment









 $\overline{AF} = \frac{1}{3} \cdot \overline{AB}$

— Scott Cobel

The vertex angles of a star sum to $180\ensuremath{^\circ}$



— Fouad Nakhli

Viviani's theorem I

The perpendiculars to the sides from a point on the boundary or within an equilateral triangle add up to the height of the triangle.



This shows a particular example, with C'GQ collinear, rather than the general case

— Samuel Wolf

Viviani's theorem II

The perpendiculars to the sides from a point on the boundary or within an equilateral triangle add up to the height of the triangle.



— Ken-Ichiroh Kawasaki

A theorem about right angles

The internal bisector of the right angle of a right triangle bisects the square on the hypotenuse



— Roland H. Eddy

Area and the projection theorem of a right triangle



— Sidney H. Kung

Chords and tangents of equal length

If circle C_1 passes through the center O of circle C_2 , the length of the common chord \overline{PQ} is equal to the tangent segment \overline{PR} .



— Roland H. Eddy

Completing the square

$$x^2 + ax = (x + a/2)^2 - (a/2)^2$$



— Charles D. Gallant

Algebraic areas I

$$(a+b)^2 + (a-b)^2 = 2(a^2 + b^2)$$



— Shirley Wakin

Algebraic areas II

$$(a+b+c)^{2} + (a+b-c)^{2} + (a-b+c)^{2} + (a-b-c)^{2} = (2a)^{2} + (2b)^{2} + (2c)^{2}$$





— Sam Pooley and K. Ann Drude

Sum of squares identity

$$(a^2 + b^2)(c^2 + d^2) = (ab + bc)^2 + (bd - ac)^2$$



— Diophantus of Alexandria

Polygonal numbers

The
$$k^{\text{th}}$$
 n-gonal number is $1 + (k-1)(n-1) + \frac{1}{2}(k-2)(k-1)(n-2)$



— Dave Logothetti

The volume of a frustrum of a square pyramid



$$V_{P_1} = \frac{h}{b-a} \cdot V_{P_2} = \frac{h}{b-a} \cdot \frac{1}{3} \left(b^3 - a^3 \right) = \frac{h}{3} \left(a^2 + ab + b^2 \right)$$

— The Moscow Papyrus, c. 1850 BCE

The volume of a hemisphere via Cavalieri's Principle



$$V_S = V_P = \frac{1}{3}r^2 \cdot 2\pi r = \frac{2}{3}\pi r^3$$


— Sidney H. Kung

Trigonometry, Calculus, & Analytic Geometry

| Sine of the sum | | | | | | | | | | | | | | | | | 2 | C |
|-----------------|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|---|---|
| Juie of the sum | | | | | | | | | | | | | | | | | _ | |

Sine of the sum

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$
 for $\alpha + \beta < \pi$



$$c = a\cos\beta + b\cos\alpha$$

$$r = 1/2 \implies \sin\gamma = \frac{c/2}{1/2} = c, \ \sin\alpha = a, \ \sin\beta = b$$

$$\sin(\alpha + \beta) = \sin(\pi - (\alpha + \beta)) = \sin\gamma = \sin\alpha\cos\beta + \sin\beta\cos\alpha$$

— Sidney H. Kung