

Diff Eq Notes

Marcell J. Vazquez-Chanlatte September 3, 2012

Contents

1	Initial Definitions	2
2	Operator Notation	2
3	Stability	2
3.1	Definition	2
4	Initial Value Problems	3
5	Seperable DE	3
5.1	Definition	3
5.1.1	Form	3
5.1.2	Solution	3
6	Homogenous Equations	3
6.1	Form	3
6.2	Technique for Solving:	3
7	Exact Equations	3
7.1	Form	3
7.2	Technique for Solving:	3
8	Autonomous Equations	4
8.1	Form	4
8.2	Anaylsis	4
8.2.1	Differentiation Fields	4
9	Linear Diff Equations	4
9.1	First Order Linear Eq	4
9.1.1	Form	4
9.1.2	Solving using Integration Factors	4
9.1.3	Bernoulli's equations	4
9.1.4	Picard Iteration	5
9.1.5	Lipsichitz Condition	5
9.1.6	Uniform Convergence (U.C.)	5
9.1.7	Existence Theorem	6
9.1.8	Uniqueness Theorm	6
9.2	Second Order Linear Eq	6
9.2.1	The Wronskian:	6
9.2.2	Existence Theorem	7
9.2.3	Uniqueness	7
9.2.4	Second Order Linear Homogenous Diff Eq (S.O.L.H.D.E)	7
9.2.5	Second Order Linear Inhomogenous Diff Eq (S.O.L.I.D.E)	8
9.2.6	Foulrier Transform	9
9.2.7	Strum Comparison Theorem	9
9.3	General Linear Diff Eq and Variation of Parameters	9

9.3.1	Form	9
9.3.2	Equation:	10
9.3.3	Theorem for L.D.E	10
10	First Order Systems of Linear Diff Equations with Const Coefficients	10
10.1	Form	10
10.1.1	Example 1st Order	10
10.2	Solution	10
10.3	Classification	10
10.3.1	Theorem: Change of coordinates	11

1 Initial Definitions

- Definition:
 - DE is an equation that describes the properties of an unknown function(s)
- Ordinary DE:
 - describes functions of 1 variable
- Partial DE:
 - describes multivariable functions
- Notation:
 - independent variable: y
 - dependent variable: t

2 Operator Notation

Definition: $\frac{d^n}{dt^n} = D^n \rightarrow f^{(n)} = D^n(f)$

3 Stability

3.1 Definition

1. Stability \equiv a system in which long term behavior does not depend within some variation on initial conditions
2. For Linear D.E. y is Stable iff:

$$\lim_{t \rightarrow \infty} y_h = 0 \pm \epsilon \quad (1)$$

3. For S.O.L.E

(a) Stable iff either of the equivalent statements is true:

- i. $Re(r) < 0$
- ii. $a, b, c > 0$ or $a, b, c < 0$

4 Initial Value Problems

$$IVP = \begin{cases} DE \\ y = C_1 \\ \vdots \\ y^{(n)} = C_n \end{cases} \quad (2)$$

5 Seperable DE

5.1 Definition

5.1.1 Form

$$\frac{dy}{dt} = f(y)g(t) \quad (3)$$

5.1.2 Solution

1. Let DE be $\frac{dy}{dt} = f(y)g(t)$
2. Separating terms: $\frac{dt}{f(y)} \frac{dy}{dt} = \frac{dt}{f(y)} f(y)g(t) \implies \frac{dy}{f(y)} = g(t)dt$
 - (a) If $f(y)^{-1}$ is not defined from some t, solve outside of that range and consider those t separately
3. Integrate both sides: $\int \frac{dy}{f(y)} = \int g(t)dt$
4. If possible solve for $y(t)$

6 Homogenous Equations

6.1 Form

$$\frac{dy}{dt} = f\left(\frac{y}{t}\right) \quad (4)$$

6.2 Technique for Solving:

1. Let $v = \frac{y}{t} \implies y = vt$
2. Change of variable: $y' = v't + v \implies v't + v = f(v)$
3. Moving v to left side: $t \frac{dv}{dt} = f(v) - v = g(v)$
4. Separating: $\frac{dv}{g(v)} = \frac{dt}{t}$
5. Solve using Seperable D.E. techniques

7 Exact Equations

7.1 Form

$$\frac{\partial}{\partial t} \Psi(f(t), y(t)) = \frac{\partial \Psi}{\partial f} \frac{df}{dt} + \frac{\partial \Psi}{\partial y} \frac{dy}{dt} \quad (5)$$

7.2 Technique for Solving:

1. Suppose DE is of the form: $M(x, y) + N(x, y)y_x = 0$
2. If $M_y = N_x$, then DE is an Exact Eq, solve for $\Psi(x, y) = \Psi(f(x), y(x))$

8 Autonomous Equations

8.1 Form

$$y' = f(y) \quad (6)$$

8.2 Analysis

8.2.1 Differentiation Fields

9 Linear Diff Equations

Definition: For an operator, L , the DE: $L(y) = 0$ is linear iff:

- $L(y_1 + y_2) = L(y_1) + L(y_2)$
- $L(cy) = cL(y)$

9.1 First Order Linear Eq

9.1.1 Form

$$y' + p(t)y = f(t) \quad (7)$$

9.1.2 Solving using Integration Factors

1. Let μ be a mult factor s.t. $\mu y' + \mu' y = g(t) \implies [\mu(t)y(t)]' = g(t)$
2. Thus $\mu' = \mu p(t) \implies \frac{d\mu}{\mu} = p(t)dt \implies \mu = e^{\int p(t)dt}$
3. Therefore

$$y(t)e^{\int p(t)dt} = \int g(t)dt \quad (8)$$

□

9.1.3 Bernoulli's equations

- Form

$$y' + p(t)y = q(t)y^n, n \in \mathbb{Z} \quad (9)$$

- Solution

1. Let $v = y^{1-n} \implies v' = (1-n)y^{-n}y'$
2. Thus $y' = \frac{v'}{1-n}$ and $y = y^n v$
3. Substituting in Bernoulli equation: $\frac{v'}{1-n}y^n + p(t)y^n v = q(t)y^n$
4. Moving into standard form:

$$v' + (1-n)p(t)v = (1-n)q(t) \quad (10)$$

5. Solve using Integration Factors □

9.1.4 Picard Iteration

- Integral Equations

Suppose f is continuous, then a function $y = \Phi(t)$ solves the IVP iff $y = \Phi(t)$ solves the corresponding integral equation:

$$y(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds \quad (11)$$

- Idea

1. Let $f(t) = \frac{dy}{dt}$
2. Construct a sequence of functions $\{g_n(t) : n \geq 0, n \in \mathbb{Z}\}$ that converges to soln:
 - (a) $y_0(t) = y_0$
 - (b) $y_{n+1}(t) = y_0 + \int_{t_0}^t f(s, y_n(s)) ds$

9.1.5 Lipschitz Condition

1. For $f(t, y) \in \mathbb{R}$, f is Lipschitz iff $\exists L \in \mathbb{R}$:

$$|f(t_1, y_1) - f(t_2, y_2)| \leq L \cdot |(y_1 - y_2)| \quad (12)$$

2. If $\Delta y \neq 0$ then this can be thought of as:

$$\left| \frac{\Delta f}{\Delta y} \right| \leq L \quad (13)$$

3. Lemma: if f_y is bounded then f is Lipschitz

9.1.6 Uniform Convergence (U.C.)

- Definition:

A sequence of functions $\{f_n(t) : n \geq 0; n \in \mathbb{Z}\}$ defined on the interval I uniformly converges to $f(t)$ iff $\forall t \in I, \exists N \in \mathbb{Z}$ s.t. $|f_n(t) - f(t)| < \epsilon$ everywhere on $I \forall n > N$

- Theorem: Given $f_n(t)$ is continuous on I , if $\lim_{n \rightarrow \infty} f_n(t) \rightarrow f(t)$ with U.C, then:

1. f is continuous
2. If f_n is differentiable, then f is differentiable and f'_n U.C. to f'
3. The limit is commutative with respect to integration

$$\lim_{n \rightarrow \infty} \int_I f_n(t) dt = \int_I \lim_{n \rightarrow \infty} f_n(t) dt \quad (14)$$

- Weierstrass M Test

Theorem:

- If $\forall n \in \mathbb{Z}, |f_n(t)| \leq M_n$ and if $\sum_{n=0}^{\infty} M_n < L$ for some $L \in \mathbb{R}$,
- Then $\sum_{n=0}^{\infty} f_n(t)$ Converges Uniformly on I

9.1.7 Existence Theorem

1. Claim:

(a) If:

- i. $f(y)$ is continuous
- ii. f is Lipschitz w.r.t. $y \in R \equiv \{t, y) : |t - t_0| \leq T \text{ and } |y - y_0| \leq k\}$
- iii. $\sum_{k=1}^{\infty} [y_k(t) - y_{k-1}(t)]$ converges uniformly

(b) Then: \exists a solution to the IVP on the interval $|t - t_0| \leq T_1 = \min(T, \frac{k}{m})$ where $|f(t, y)| \leq M \in R$

2. Proof:

(a) Converting the IVP to an I.E.: $y(t) = y_0 + \int_{t_0}^t f(s, y(s))ds$

(b) Note that: $|y_k(t) - y_{k-1}(t)| \leq \frac{M}{L} \frac{L^n (t-t_0)^n}{n!} \leq \frac{M}{L} \frac{L^n T_1^n}{n!}$

(c) Define: $M_n \equiv \sum_{k=1}^{\infty} \frac{M}{L} \frac{(LT_1)^n}{n!} = \frac{M}{L} (e^{LT_1} - 1)$

(d) Apply the Weierstrass M Test, because $\frac{M}{L} \frac{(LT_1)^n}{n!}$ converges, then $\sum_{k=1}^{\infty} [y_k - y_{k-1}]$ converges

(e) Thus the series $\{y_n : n \geq 1\}$ converges uniformly on the interval.

(f) Therefore \exists a solution to the IVP \square

9.1.8 Uniqueness Theorem

1. Claim:

(a) If $\Phi(t)$ and $\Psi(t)$ are solutions of $y' \equiv f(y, t) \in R$ and if f is Lipschitz w.r.t. $y \in R$

(b) Then $|\Phi(t) - \Psi(t)| \leq e^{L|t-t_0|} |\Phi(t_0) - \Psi(t_0)| = 0$

i. Because they solve the same I.V.P. $|\Phi(t_0) - \Psi(t_0)| = 0$

(c) Equivalently: Then $\Phi(t) = \Psi(t)$

2. Proof:

(a) $E \equiv |\Phi(t) - \Psi(t)|^2$

i. Note that $E \geq 0$

(b) $\frac{d}{dt} E = 2(\Phi(t) - \Psi(t))(\Phi'(t) - \Psi'(t))$

(c) $E' \stackrel{DE}{=} 2(\Phi(t) - \Psi(t))(f(t, \Phi) - f(t, \Psi))$

(d) $E' \stackrel{Lip}{\leq} 2|\Phi(t) - \Psi(t)| L |\Phi(t) - \Psi(t)|$

(e) Thus $E' \leq 2LE \implies E' - 2LE \leq 0 \implies (E(t)e^{-2Lt})' \leq 0$

i. Note that E' is strictly decreasing

(f) Therefore: $e^{-t} E(t) \leq e^{2Lt_0} E(t_0) \implies E(t) \leq e^{2L(t-t_0)} E(t_0)$

(g) Substituting: $|\Phi(t) - \Psi(t)|^2 \leq e^{2L(t-t_0)} |\Phi(t_0) - \Psi(t_0)|^2$

(h) Because of absolute value: $|\Phi(t) - \Psi(t)| \leq e^{2L(t-t_0)} |\Phi(t_0) - \Psi(t_0)|$

(i) Because they solve the same I.V.P. $|\Phi(t_0) - \Psi(t_0)| = 0$

(j) Thus $\Phi(t) = \Psi(t) \square$

9.2 Second Order Linear Eq

9.2.1 The Wronskian:

$$W(f, g)(t) = \begin{vmatrix} f_1(x) & f_2(x) & \cdots & f_n(x) \\ f_1'(x) & f_2'(x) & \cdots & f_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ f_1^{(n-1)}(x) & f_2^{(n-1)}(x) & \cdots & f_n^{(n-1)}(x) \end{vmatrix}, \quad x \in I \quad (15)$$

9.2.2 Existence Theorem

1. Claim: For all D.E. there exists a $y(t)$ that satisfies it locally on some interval
2. Proof:
 - (a) Let $y' = v \rightarrow v' = y''$ 2) Therefore $v' = -py' - qy = -pv - qy$, by plugging into the DE
 - (b) In matrix form:

$$\begin{bmatrix} y \\ v \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -q & -p \end{bmatrix} \begin{bmatrix} y \\ v \end{bmatrix} \quad (16)$$

- (c) Note that this is a linear first order matrix system which there is an existence theorem for

9.2.3 Uniqueness

9.2.4 Second Order Linear Homogenous Diff Eq (S.O.L.H.D.E)

- Form

$$a(x)\frac{d^2y}{dt^2} + b(x)\frac{dy}{dt} + c(x)y = 0 \quad (17)$$

- Theorem: The general solution to S.O.L.H.E

Claim: The general soln of eq1 $\equiv [y'' + p(t)y' + q(t)y = 0]$ is:

$$y_h = c_1y_1 + c_2y_2 \quad (18)$$

- Proof:

– Q1:

Given y_1 and y_2 are solutions, why is $c_1y_1 + c_2y_2$ a solution

1. $Eq1 = D^2(y) + p(t)D(y) + q(t)y = 0$
2. $Eq1 = [D^2 + p(t)D + q(t)]y = 0$
3. Let $L = [D^2 + p(t)D + q(t)] \rightarrow Eq1 \equiv L(y) = 0$
4. Notice the L is a linear operator and thus obeys the superposition principle
5. Thus $y = c_1y_1 + c_2y_2$ is a solution \square

– Q2:

Given 2 indepnt solutions y_1 and y_2 for the DE, \forall IVP and its unique solution y , $\exists (c_1, c_2) \in \mathbb{C}^2$ s.t. $y = c_1y_1 + c_2y_2 \equiv \vec{y} \cdot \vec{c}$

– Q3:

* Abel's Identity

1. If u, v solve the D.E. then $W' + p(t)W = 0 \rightarrow ce^{-\int p(t)dt}$
2. Alternatively: $W' + p(t)W = 0 \implies W' = 0 \implies W = C$

* Finding the general solution

Goal: The general soln is of the form $y = \vec{y} \cdot \vec{c}$

1. Recall the matrix form of the D.E. from the Existence theorem proof.
2. Also Recall that that the equation was only solvable if $W(y_1, y_2)(t_0) \neq 0$
3. Observe that $W' = (uv' + uv'') - (u'V + u'v') = -pW$
4. Lemma: if u, v are linearly dependent, then $W(u, v) = 0$ on I

- Generating Second Solution

1. Claim: if $y_1 \neq 0$ be a solution to the D.E. then,

$$y_2 = Cy_1 \int \frac{e^{-\int p dt}}{y_1^2} \quad (19)$$

and y_2 = solution independent of y_1

2. Proof:

- (a) Consider $(\frac{y_2}{y_1})' = \frac{y_1 y_2' - y_1' y_2}{y_1^2} = \frac{W(y_1, y_2)}{y_1^2}$
- (b) Given that $W' + p(t)W = 0 \implies W(t) = ce^{-\int p(t) dt}$
- (c) Thus: $\int (\frac{y_2}{y_1})' dt = C \int \frac{e^{-\int p dt}}{y_1^2}$
- (d) Solving: $y_2 = Cy_1 \int \frac{e^{-\int p dt}}{y_1^2} \square$

9.2.5 Second Order Linear Inhomogenous Diff Eq (S.O.L.I.D.E)

- Form

$$y^{(n)}(t) + p(t)y'(t) + q(t)y = f(t) \quad (20)$$

- General Solution

1. Claim: The general soln of $y^{(n)}(t) + p(t)y'(t) + q(t)y = f(t)$ is:

$$y = y_h + k(t) \quad (21)$$

- (a) $y_h = c_1 y_1 + c_2 y_2$ is the solution to the homogenous equation i.e. $f(t) = 0$
- (b) Functional Offset ($k(t)$): variation or 'offset' from the homogenous equation

2. Proof:

- (a) Sub-Claim: $y_h + k$ is a solution
 - i. Using Operator notation: $D^2 y + pDy + qy = f \implies [D^2 + pD + q](y) = f$
 - ii. Let $L^2 + pD + q \implies L(y) = f$
 - A. Note that L is linear
 - iii. $L(y_h + k) = L(y_h) + L(y_p)$
 - iv. $L(y_h) = 0, L(y_p) = f \implies L(y_h + k) = f + 0 = f \square$
- (b) Sub-Claim: $\forall y_i$, if y_i is a solution to the S.O.L.I.D.E, then $y_i = y_h + k$
 - i. $[L(y_i) = f \text{ and } L(k) = f] \implies L(y_i - k) = f - f = 0$
 - ii. By Existence of S.O.L.H.D.E, $L(y_i - k) = 0 \implies y_i - k = y_h$
 - iii. Thus $y_i = k + y_h \square$

- Exponential Shift Law

$$P(D)[e^{\alpha u(t)}] = p(D + \alpha)u(t)[e^{\alpha t}] \quad (22)$$

- Exponential-Polynomial Functional offsets

- Form

$$ay'' + by' + cy = e^{\alpha t} g(t); \alpha \in \mathbb{C} \quad (23)$$

- Characteristic Polynomial

$$p(r) = aD^2 + bD + c \quad (24)$$

Note that the DE in Operator notation is: $[aD^2 + bD + c]$

- Finding Particular Solution for S.O.L.E

Theorem:

- * Let k be s.t. $(r - \alpha)^k$ are roots of $p(\alpha)$
- * Then

$$y_p = \frac{t^k e^{\alpha t}}{p^{(k)}(\alpha)} \quad (25)$$

- Method of Undetermined Coefficients

Idea: if $f(t)$ is comprised of strict multiplications (no division) sinusoidal, exponentials, and polynomials then the solution of the S.O.L.E with const coefficients is in terms of the same types you began with.

Cases:

if $f(t) = e^{\alpha t}$ (polynomial of $\deg(k + m)$), then guess $y_p = e^{\alpha t} \sum_{j=0}^k C_j t^j$

- Annihilator Method

1. Given: $(D^2 + aD + b)y = e^{\alpha t} f(t)$, where $f(t)$ is a polynomial
2. Attempt to multiply both sides by a Differential Operator (G) s.t.

$$G(e^{\alpha t} f(t)) = 0 \implies G(D^2 + aD + b)y = 0 \quad (26)$$

3. Notice that this is now a homogenous equation. Solve by finding the roots of the characteristic equation and applying the method of undetermined coefficients

- Lagrange Variation of Parameters

- Equation:

$$y_p = \int \frac{y_1 f(x)}{W(y_1, y_2)} dt + \int \frac{y_2 f(x)}{W(y_1, y_2)} dt \quad (27)$$

- Derivation:
See General Derivation

9.2.6 Foulrier Transform

9.2.7 Strum Comparison Theorem

Theorem:

1. If:
 - (a) $u'' + q_1(t)u = 0$ and $v'' + q_2v = 0$
 - (b) $q_1 > q_2$
2. Then:
 - (a) u vanishes at some point between 2 zeros of v

9.3 General Linear Diff Eq and Variation of Parameters

9.3.1 Form

$$y^{(n)}(x) + \sum_{k=0}^{n-1} a_k(x) y^{(k)}(x) = f(x) \quad (28)$$

9.3.2 Equation:

$$\sum_{k=0}^n [y_k(x) \int \frac{W_k(x)}{W(X)} dx] \quad (29)$$

$W(x) \equiv$ Wronskian determinant of the fundamental system and $W_i(x) \equiv$ the Wronskian determinant of the fundamental system with the $i - th$ column replaced by $(0, 0, \dots, f(x))$

- Derivation:

9.3.3 Theorem for L.D.E

If $u(t) + iv(t)$ is a solution to the D.E. then $u(t) \wedge v(t)$ are solutions

10 First Order Systems of Linear Diff Equations with Const Coefficients

10.1 Form

$$\vec{r}' = A\vec{r} \quad (30)$$

10.1.1 Example 1st Order

$$\vec{r}' = \begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} ax + by \\ cx + dy \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \quad (31)$$

10.2 Solution

1. Guess: $\vec{r} = \vec{a}e^{\lambda t}$, $\vec{a} \in D$

2. Thus $\vec{r}' = \vec{a}e^{\lambda t} \implies \lambda \vec{a} = A\vec{a}$

3. Factoring we get $(A - \lambda I)\vec{a} = \vec{0}$

4. Characteristic Polynomial $\equiv |A - \lambda I|$

(a) For the 2-d version we get

$$|A - \lambda I| = \lambda^2 + tr(A)\lambda + |A| \quad (32)$$

5. Solve for Eigen Value and Corresponding Eigen Vectors

(a) Complete e-value: K-Repeated Eigen value produces K linearly independent eigen vectors

(b) Incomplete e-value: Does not produce enough e-values. Solution: assume solution is:

$$\vec{r} = u(t)\vec{a}e^{\lambda t} + \vec{c} \quad (33)$$

i. Typically $u(t)$ is a polynomial as in the $D-1$ case

10.3 Classification

Let $p \equiv tr(A)$ and $q \equiv |A|$

10.3.1 Theorem: Change of coordinates

1. Claim: $\vec{x}' = A\vec{x} \equiv \vec{y}' = B\vec{y} \iff |A - I| = |B - I|$ and $(\text{tr}[A])^2 - 4|A| \neq 0$
 - (a) $(\text{tr}[A])^2 - 4|A| \neq 0$ excludes repeated roots
2. Alternate version: \exists non-singular matrix U s.t. $\vec{y} = U\vec{x}$ and $B = UAU^{-1}$
 - (a) This is a change of coordinates
3. Light Proof: Suppose $\vec{x}' = A\vec{x} \equiv \vec{y}' = B\vec{y}$
 - (a) By the claim $|B - \lambda I| = |UAU^{-1} - \lambda I|$
 - (b) Note that $\lambda I = \lambda U U^{-1} = U \lambda I U^{-1} \implies |B - \lambda I| = |UAU^{-1} - U(\lambda I)U^{-1}|$
 - (c) Factoring $|U(A - \lambda I)U^{-1}| = |U||A - \lambda I||U^{-1}|$
 - (d) Using the commutative property of multiplication: $|U||U^{-1}||A - \lambda I| = |A - \lambda I|$
 - (e) Thus: $|B - \lambda I| = |A - \lambda I|$