


Comparative analysis of the treatment of asymptomatic and symptomatic complex aortic aneurysms

Jennifer Ci, Thu Vu, Lily Wang | UW Biostatistics
Dr. Sara Zettervall MD, MPH | UW Vascular Surgery
Spring 2023

Our Team



Jennifer Ci




Thu Vu



Lily Wang



Dr. Sara Zettervall, MD, MPH



Dr. Kirsten Dansey, MD, MPH

Scientific Background

> **Complex Aortic Aneurysm**


- **Thoraco-abdominal aortic aneurysm (TAAA):** involves the aorta in both the chest and abdomen at the level of vital branches to the intestines and kidneys
- Open aortic repair associated with 20% mortality

> **Fenestrated-Branched Endovascular Aortic Repair (F-BEVAR)**

- Safer alternative to open repair
- Custom devices for each patient based on their unique anatomy
- **Further research needed for surgical practice and perioperative management**

Primary Objective

- > To compare outcomes of asymptomatic and symptomatic patients with complex aortic aneurysm treated with F-BEVAR



Data Source

> Vascular Quality Initiative (VQI)

- Large, multi-center database
- 14 major vascular registries

> Thoracic and Complex EVAR

- Over 25,000 procedures from 2010 to 2022

> Recorded Variables

- Patient demographics
- Comorbidities
- Operative and anatomic differences
- Short-term and long-term outcomes

Population of Interest

- > **Patients with Complex Aortic Aneurysm treated with F-BEVAR**
- > **Primary Covariate: Asymptomatic or Symptomatic**
 - **Asymptomatic:** Presenting no symptoms, elective (planned procedure)
 - **Symptomatic:** Presenting symptoms, urgent or emergent (procedure given within 24 or 4 hours of presentation, respectively)
- > **Excluded patients with aortic ruptures**

Methods

- > **Cox Proportional-Hazards**

- Primary Outcome: All-Cause Mortality

- > **Logistic Regression using GEE with working independence**

- Binary Secondary Outcomes

- > **Clustered by Center (147)**

- > **Confounders**


- Expert consultation
- Univariate analysis
- Include: demographics, comorbidities, clinical & perioperative factors

Descriptive Statistics

Variable	Asymptomatic (N=3315)	Symptomatic (N=442)	Overall (N=3757)
Age			
Mean (SD)	73.5 (7.89)	70.8 (10.1)	73.2 (8.23)
Sex			
Female	800 (24.1%)	177 (40.0%)	977 (26.0%)
Male	2515 (75.9%)	265 (60.0%)	2780 (74.0%)
Mortality			
Yes	390 (11.8%)	94 (21.3%)	484 (12.9%)
No	2925 (88.2%)	348 (78.7%)	3273 (87.1%)
Follow-Up Days			
Median [IQR]	398 [15, 1046]	309 [11, 660]	391 [13, 1015]

Primary Outcome Results

All-Cause Mortality



Year 1

A: 0.89

S: 0.77

Year 5


A: 0.79

S: 0.66

		Number at risk										
Strata	Asymptomatic	3314	1849	984	802	631	472	339	208	60	8	1
	Symptomatic	442	199	105	90	69	45	31	22	10	1	1
		0	1	2	3	4	5	6	7	8	9	10
		Time (in years)										

Secondary Outcome Results

First Reintervention




Number at risk

Strata	Asymptomatic	2126	1649	771	540	374	254
	Symptomatic	253	176	72	52	31	20
		0	1	2	3	4	5
		Time (in years)					

Secondary Outcome Results

Multivariate Logistic Regression Models



Conclusions

- > **Survival differs between asymptomatic and symptomatic patients**
 - Mortality higher for symptomatic patients
- > **Hospital/ICU Stay & some post-op factors differs**
 - Overall, symptomatic has higher odds
- > **Increase effort to detect patients with TAAA before presenting symptoms**
- > **Provide evidence for insurance companies to cover F-BEVAR procedures**

Limitations & Next Steps

> Limitations

- Retrospective analysis of all hospitals participating in the VQI
- Coding errors, missing data, self-reported data
- Possible other variables not accounted for

> Next Steps

- Submit to and present at the Western Vascular Society Meeting
- Publish results in the Journal of Vascular Surgery
- Future projects focusing on other subgroups (sex and race)

Acknowledgments

Dr. Lloyd Mancl, PhD

Dr. Patrick Heagerty, PhD

Dr. Sara Zettervall, MD, MPH

Dr. Kirsten Dansey, MD, MPH

Thank you! Questions?

