



ENGR 298 - Data Analysis  
Blood Oxygen Data Final Project  
I have followed the honor code 5/11/22

Thomas Wasylenko

## Background:

“Pulse oximetry is routinely used for monitoring patients' oxygen saturation levels with little regard to the variability of this physiological variable. There are few published studies on oxygen saturation variability (OSV), with none describing the variability and its pattern in a healthy adult population. The aim of this study was to characterize the pattern of OSV using several parameters: the regularity (sample entropy analysis), the self-similarity (detrended fluctuation analysis (DFA)), and the complexity (multiscale entropy (MSE) analysis). Secondly, to determine if there were any changes that occur with age.”

## Procedure:

1. Clean the pulse oximeter and place on finger of participants choosing
2. Place the pulse pressure transducer on the adjacent finger
3. Wrap the Respiratory band around the umbilicus of the participant
4. Preferably have participant sitting with the fingers relatively still
5. Test equipment to ensure correct readings
6. Once the equipment has been checked stop the test and start the official recording

# Study

## Parameters:

- Healthy Adult
- Gender
- Smoking Status
- BMI
- Age



Pulse Oximeter Picture  
PhysioNet: Pattern Analysis of Oxygen Saturation Variability

Participant No	Gender	Smoking Status	BMI	Age
1	Female	Non-Smoker	20.5	23
2	Female	Non-Smoker	21.7	21
3	Female	Smoker	23.8	45
4	Female	Ex-Smoker	23.2	66
5	Male	Non-Smoker	25.1	70
6	Male	Non-Smoker	22	20
7	Female	Non-Smoker	19.8	19
8	Female	Non-Smoker	24.7	22
9	Female	Non-Smoker	20.7	20
10	Male	Non-Smoker	23.6	20
11	Female	Non-Smoker	28.2	41
12	Female	Non-Smoker	24.4	20
13	Female	Non-Smoker	24.1	45
14	Male	Non-Smoker	26.8	49
15	Male	Non-Smoker	23.2	22
16	Male	Non-Smoker	20.6	20
17	Femae	Non-Smoker	23.1	21
18	Male	Non-Smoker	26.2	42

Participant No	Gender	Smoking Status	BMI	Age
19	Female	Non-Smoker	24.3	62
20	Male	Non-Smoker	19.9	48
21	Male	Ex-Smoker	24.5	60
22	Female	Ex-Smoker	18.5	56
23	Female	Non-Smoker	24.2	24
24	Male	Non-Smoker	23.9	45
25	Female	Non-Smoker	19.8	38
26	Male	Smoker	24.6	21
27	Male	Non-Smoker	26.7	23
28	Female	Non-Smoker	18.9	21
29	Male	Smoker	26.8	23
30	Female	Ex-Smoker	17.9	21
31	Female	Non-Smoker	21.3	35
32	Male	Non-Smoker	23	20
33	Male	Ex-Smoker	28.4	55
34	Female	Non-Smoker	20.3	20
35	Male	Non-Smoker	26.5	42
36	M	Non-Smoker	21.7	20



# Problem Definition

---

Take the data collected and perform analysis to see how and what physical factors have the greatest importance on a person's blood oxygen values.

---

Question: What physical factor has the greatest affect on blood oxygen values leading to the decrease in a person's levels?

---

Hypothesis:

- Gender will not play significant role
- Smoking status and BMI will be significant

# Process

Format blood oxygen participant data

Perform 1-way Anova tests over all the participants

Generate graphs and charts of the factors compared to each other Ex: (Male vs Female)

Multiple comparison test to see groups of significance

Perform N-way Anova tests to compare each physical factor

# Format One-Way Anova

```
Editor - anova_bloodoxygen.m
1 - filename = "reformat_osv_data.xlsx";
2
3 - gender = xlsread(filename, "A2:A37");
4 - smoking = xlsread(filename, "B2:B37");
5 - bmi = xlsread(filename, "C2:C37");
6 - age = xlsread(filename, "D2:D37");
7 - data = xlsread(filename, "E2:E37");
8
9 - save("pat_data.mat", 'gender', 'smoking', 'bmi', 'age', 'data');|
10
```

Process:

- ◇ Format Data
  - ◇ Male 0, Female 1
  - ◇ None-Smoker 0, Smoker/Ex-Smoker 1
- ◇ Load the data into Matlab
- ◇ Make each factor its own variable
- ◇ Perform Anova test to find significance in factors
- ◇ Perform Analysis of the data handed back

Gender	Smoking	BMI	Age	Average
1	0	20.5	23	98.239
1	0	21.7	21	98.163
1	1	23.8	45	94.412
1	1	23.2	66	96.344
0	0	25.1	70	93.748
0	0	22	20	96.468
1	0	19.8	19	97.955
1	0	24.7	22	98.435
1	0	20.7	20	98.703
0	0	23.6	20	97.873
1	0	28.2	41	98.801
1	0	24.4	20	97.364
1	0	24.1	45	98.009
0	0	26.8	49	98.185
0	0	23.2	22	96.805
0	0	20.6	20	98.606
1	0	23.1	21	99.001
0	0	26.2	42	98.184
1	0	24.3	62	98.704
0	0	19.9	48	98.504
0	1	24.5	60	98.2
1	1	18.5	56	96.524
1	0	24.2	24	96.976
0	0	23.9	45	96.275
1	0	19.8	38	96.87
0	1	24.6	21	96.523
0	0	26.7	23	97.77
1	0	18.9	21	98.33
0	1	26.8	23	98.186
1	1	17.9	21	99.324
1	0	21.3	35	99.508
0	0	23	20	98.645
0	1	28.4	55	97.126
1	0	20.3	20	98.838
0	0	26.5	42	97.585
0	0	21.7	20	98.218

```
Editor - anova_test.m
1 - load pat_data
2
3 - [p_gender,tbl_gender,stats_gender] = anova1(data, gender);
4 - [p_smoking,tbl_smoking,stats_smoking] = anova1(data, smoking);
5 - [p_bmi,tbl_bmi,stats_bmi] = anova1(data, bmi);
6 - [p_age,tbl_age,stats_age] = anova1(data, age);
```

# Format N-Way Anova

Gender	Smoking	BMI	Age	Average
Female	Non-Smoker	Healthy	Young	98.239
Female	Non-Smoker	Healthy	Young	98.163
Female	Smoker	Healthy	Mid	94.412
Female	Ex-Smoker	Healthy	Old	96.344
Male	Non-Smoker	Overweight	Old	93.748
Male	Non-Smoker	Healthy	Young	96.468
Female	Non-Smoker	Healthy	Young	97.955
Female	Non-Smoker	Healthy	Young	98.435
Female	Non-Smoker	Healthy	Young	98.703
Male	Non-Smoker	Healthy	Young	97.873
Female	Non-Smoker	Overweight	Mid	98.801
Female	Non-Smoker	Healthy	Young	97.364
Female	Non-Smoker	Healthy	Mid	98.009
Male	Non-Smoker	Overweight	Mid	98.185
Male	Non-Smoker	Healthy	Young	96.805
Male	Non-Smoker	Healthy	Young	98.606
Female	Non-Smoker	Healthy	Young	99.001
Male	Non-Smoker	Overweight	Mid	98.184
Female	Non-Smoker	Healthy	Mid	98.704
Male	Non-Smoker	Healthy	Mid	98.504
Male	Ex-Smoker	Healthy	Mid	98.2
Female	Ex-Smoker	Underweight	Mid	96.524
Female	Non-Smoker	Healthy	Young	96.976
Male	Non-Smoker	Healthy	Mid	96.275
Female	Non-Smoker	Healthy	Young	96.87
Male	Smoker	Healthy	Young	96.523
Male	Non-Smoker	Overweight	Young	97.77
Female	Non-Smoker	Healthy	Young	98.33
Male	Smoker	Overweight	Young	98.186
Female	Ex-Smoker	Underweight	Young	99.324
Female	Non-Smoker	Healthy	Mid	99.508
Male	Non-Smoker	Healthy	Young	98.645
Male	Ex-Smoker	Overweight	Mid	97.126
Female	Non-Smoker	Healthy	Young	98.838
Male	Non-Smoker	Overweight	Mid	97.585
Male	Non-Smoker	Healthy	Young	98.218

Formatting:

- ◇ Gender
  - ◇ Male, Female
- ◇ Smoking Status
  - ◇ Non-Smoker, Ex-Smoker, Smoker
- ◇ BMI (Per [BMI CDC](#))
  - ◇ <18.5 Underweight
  - ◇ 18.5-25 Healthy
  - ◇ 25-30 Overweight
  - ◇ >30 Obesity
- ◇ Age
  - ◇ 0-30 Young
  - ◇ 31-64 Mid
  - ◇ 65 above Old

```
Editor - anova_nway.m
anova_bloodoxygen.m  anova_test.m  mult_compare.m  test.m  anova_nway.m  +
1  load nway
2
3  %Gender vs Smoking
4  varnamesGS = {'Gender'; 'Smoking'};
5  anovan(data, {chargender charsmoking}, 2, 2, varnamesGS);
6  %Gender vs BMI
7  varnamesGB = {'Gender'; 'BMI'};
8  anovan(data, {chargender charbmi}, 2, 2, varnamesGB);
9  %Gender vs Age
10 varnamesGA = {'Gender'; 'Age'};
11 anovan(data, {chargender charage}, 2, 2, varnamesGA);
12 %Smoking vs BMI
13 varnamesSB = {'Smoking'; 'BMI'};
14 anovan(data, {charsmoking charbmi}, 2, 2, varnamesSB);
15 %Smoking vs Age
16 varnamesSA = {'Smoking'; 'Age'};
17 anovan(data, {charsmoking charage}, 2, 2, varnamesSA);
18 %BMI vs Age
19 varnamesBA = {'BMI'; 'Age'};
20 anovan(data, {charbmi charage}, 2, 2, varnamesBA);
```

Process:


- ◇ Format data
- ◇ Load data as character arrays
- ◇ Perform Anova test comparing all combinations
- ◇ Perform analysis of data



# Multiple Comparison

Process:


- ❖ Collect data from one-way Anova
- ❖ Perform Multiple Comparison for each factor
- ❖ Analyze data to find groups with large significance

A screenshot of a MATLAB Editor window titled 'Editor - mult\_compare.m'. The window has a tab bar at the top with several open files: 'anova\_bloodoxygen.m', 'anova\_test.m', 'mult\_compare.m' (the active file), 'test.m', and 'anova\_nway.m'. To the right of the tab bar is a 'Variables - tbld' window. The main editor area shows four lines of MATLAB code, each preceded by a line number (1-4) and a minus sign. The code uses the 'multcompare' function to perform multiple comparisons on the results of one-way ANOVAs for different variables: gender, smoking, bmi, and age.

```
1 - [c_gender,m_gender,h_gender,gnames_gender] = multcompare(stats_gender);  
2 - [c_smoking,m_smoking,h_smoking,gnames_smoking] = multcompare(stats_smoking);  
3 - [c_bmi,m_bmi,h_bmi,gnames_bmi] = multcompare(stats_bmi);  
4 - [c_age,m_age,h_age,gnames_age] = multcompare(stats_age);
```




# Findings One-Way Anova



Gender Factor Test  
Male 0, Female 1


ANOVA Table					
Source	SS	df	MS	F	Prob>F
Groups	1.868	1	1.868	1.21	0.2792
Error	52.527	34	1.54491		
Total	54.3951	35			



Smoking Factor Test  
Non-Smoker 0, Smoker 1


ANOVA Table					
Source	SS	df	MS	F	Prob>F
Groups	4.027	1	4.02697	2.72	0.1084
Error	50.3681	34	1.48141		
Total	54.3951	35			

# Findings One-Way Anova



BMI Factor Test


ANOVA Table					
Source	SS	df	MS	F	Prob>F
Groups	53.6987	31	1.73222	9.95	0.0185
Error	0.6964	4	0.1741		
Total	54.3951	35			




Age Factor Test

ANOVA Table					
Source	SS	df	MS	F	Prob>F
Groups	36.84	18	2.04667	1.98	0.0826
Error	17.5551	17	1.03265		
Total	54.3951	35			



# Findings Multiple Comparisons




Gender Multi Test  
Male 0, Female 1



BMI Multi Test



Smoking Multi Test  
Non-Smoking 0  
Smoking 1



Age Multi Test



# Findings N-Way Anova

Analysis of Variance					
Source	Sum Sq.	d.f.	Mean Sq.	F	Prob>F
Gender	2.2121	2	1.10604	0.87	0.4306
Smoking	5.5339	2	2.76694	2.17	0.1324
# Gender*Smoking	8.7859	2	4.39296	3.45	0.0454
Error	36.9762	29	1.27504		
Total	54.3951	35			

Gender vs Smoking  
Smoking has greater significance

Analysis of Variance					
Source	Sum Sq.	d.f.	Mean Sq.	F	Prob>F
Gender	2.4348	2	1.21741	0.73	0.4887
BMI	0.0627	2	0.03136	0.02	0.9813
# Gender*BMI	1.4288	1	1.42876	0.86	0.361
Error	49.8045	30	1.66015		
Total	54.3951	35			

Gender vs BMI  
Gender has slightly greater significance

Analysis of Variance					
Source	Sum Sq.	d.f.	Mean Sq.	F	Prob>F
Gender	2.3269	2	1.16347	1.01	0.3765
Age	14.8038	2	7.40188	6.43	0.0049
# Gender*Age	3.1022	2	1.55111	1.35	0.2758
Error	33.39	29	1.15138		
Total	54.3951	35			

Gender vs Age  
Age has greater significance

Analysis of Variance					
Source	Sum Sq.	d.f.	Mean Sq.	F	Prob>F
Smoking	6.5703	2	3.28516	2.35	0.113
BMI	0.8764	2	0.43821	0.31	0.7331
# Smoking*BMI	6.6052	2	3.30262	2.37	0.1118
Error	40.4925	29	1.39629		
Total	54.3951	35			

Smoking vs BMI  
Smoking has greater significance

Analysis of Variance					
Source	Sum Sq.	d.f.	Mean Sq.	F	Prob>F
Smoking	7.6392	2	3.81961	5.54	0.0094
Age	16.7943	2	8.39713	12.17	0.0002
# Smoking*Age	11.8598	3	3.95327	5.73	0.0035
Error	19.3201	28	0.69		
Total	54.3951	35			

Smoking vs Age  
Age has greater significance

Analysis of Variance					
Source	Sum Sq.	d.f.	Mean Sq.	F	Prob>F
BMI	0.0255	2	0.01274	0.01	0.9888
Age	14.8744	2	7.4372	6.56	0.0046
# BMI*Age	7.0695	3	2.3565	2.08	0.1255
Error	31.7242	28	1.13301		
Total	54.3951	35			

BMI vs Age  
Age has greater significance

# Summary

Initial Question: What physical factor has the greatest affect on blood oxygen values leading to the decrease in a person's levels?

Hypothesis:

- ◇ -Gender will not play significant role
- ◇ -Smoking status and BMI will be significant

After Analysis:

- ◇ Age has the greater significance over all other factors
- ◇ Smoking is the next significant factor
- ◇ BMI is least significant

Final Answer in relation to Hypothesis:

My hypothesis was not proven to be correct entirely, only that smoking status plays a significant role.

Challenges:

- ◇ Small Sample Size for proper analysis
  - ◇ Possible Solution: Larger Sample Size with greater variability in factors

[GitHub Link](#)