

# Theoretical modeling of the collective tunneling of a Wigner necklace

Dominik Szombathy,<sup>1,2</sup> Miklós Antal Werner,<sup>1,2</sup> Cătălin Pașcu Moca,<sup>1,3</sup> and Gergely Zaránd<sup>1,2</sup>

<sup>1</sup>Department of Theoretical Physics, Institute of Physics,

Budapest University of Technology and Economics, Budafoki út 8., H-1111 Budapest, Hungary

<sup>2</sup>MTA-BME Quantum Dynamics and Correlations Research Group, Institute of Physics,

Budapest University of Technology and Economics, Budafoki út 8., H-1111 Budapest, Hungary

<sup>3</sup>Department of Physics, University of Oradea, 410087, Oradea, Romania

(Dated: December 8, 2022)

To be written.

## INTRODUCTION

...




FIG. 1: Experimental setup




FIG. 3: Experimental and theoretical calculation of the tunnel splitting in the quartic potential




FIG. 2: Experimental and theoretical calculation of the tunnel splitting in the quartic potential




FIG. 4: Comparison between the theoretical and the experimental polarization

**SUPMAT FIGURES**




FIG. 5: Perpendicular tunnel splitting factors for 1,3,5 and 7 particles





FIG. 6





FIG. 7