

Theoretical modeling of the collective tunneling of a Wigner necklace

Dominik szombathy

November 2, 2022

Abstract

Contents

1	Path Integral Formalism (introduction)	4
2	Wigner Crystals	4
3	Instanton Formalism	4
3.1	Free Particle	4
3.2	Harmonic Oscillator	4
3.3	One particle tunneling	4
3.3.1	One Particle Hamiltonian	4
3.3.2	Action Integral Calculation	4
3.3.3	Tunneling calculation in quartic potentials	4
4	Experimental Considerations	4
4.1	Modeling the Potential and Fitting	4
4.2	Polarization data	4
4.3	Scaling	4
5	Many Body Tunneling Calculations - Instanton	4
5.1	Quartic Hamiltonian model	4
5.1.1	Rescaling and Dimensionless Hamiltonian	4
5.2	Milnikov method	4
5.3	Classical Equilibrium Positions	4
5.4	Harmonic Oscillator Eigenfrequencies and Eigenmodes	4
5.5	Trajectory Calculation	4
5.6	Arc Length Parameterization	4
5.7	Exact Diagonalization (ED)	4

6	Polarization	4
6.1	Classical Polarization Calculation	4
6.2	Polarization using ED	4
7	Open Questions	4
A	Mathematics	4
A.1	Gaussian Integrals	5
A.2	Cauchy's Residue Theorem	5
A.3	Bessel Problem	5
B	Algorithms and Methods	7
B.1	Monte Carlo and Simulated Annealing	7
B.2	Minimalization by Nelder-Mead algorithm	7
C	Links, Data and other stuff	7
D	Research Diary	7

1	Path Integral Formalism (introduction)
2	Wigner Crystals
3	Instanton Formalism
3.1	Free Particle
3.2	Harmonic Oscillator
3.3	One particle tunneling
3.3.1	One Particle Hamiltonian
3.3.2	Action Integral Calculation
3.3.3	Tunneling calculation in quartic potentials
4	Experimental Considerations
4.1	Modeling the Potential and Fitting
4.2	Polarization data
4.3	Scaling
5	Many Body Tunneling Calculations - Instanton
5.1	Quartic Hamiltonian model
5.1.1	Rescaling and Dimensionless Hamiltonian
5.2	Milnikov method
5.3	Classical Equilibrium Positions
5.4	Harmonic Oscillator Eigenfrequencies and Eigenmodes
5.5	Trajectory Calculation
5.6	Arc Length Parameterization
5.7	Exact Diagonalization (ED)
6	Polarization
6.1	Classical Polarization Calculation
6.2	Polarization using ED
7	Open Questions
A	Mathematics

A.1 Gaussian Integrals

A.2 Cauchy's Residue Theorem

A.3 Besel Problem

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \quad (\text{A.3.1})$$

Fourier Way Let's consider first an arbitrary function $f(x)$. One could write the fourier series of this function as

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left\{ a_n \cos\left(\frac{n\pi l}{L}\right) + b_n \sin\left(\frac{n\pi l}{L}\right) \right\}, \quad (\text{A.3.2})$$

where $l \in [l_i, l_f]$ and $L = l_f - l_i$. Let $f(x) = x^2$ and $l_i = -l_f = -\pi$ so that $L = 2\pi$. The coefficients can be calculated as

$$a_0 = \frac{2}{L} \int_{-\pi}^{\pi} f(x) dx = \frac{2}{\pi} \int_0^{\pi} dx x^2 = \frac{2}{3} \pi^2 \quad (\text{A.3.3})$$

$$\begin{aligned} a_n &= \frac{2}{L} \int_{-\pi}^{\pi} dx f(x) \cos\left(\frac{n\pi 2x}{L}\right) \\ &= \frac{2}{\pi} \left(\frac{x^2 \sin(nx)}{n} \right) \Big|_0^{\pi} - \frac{2}{\pi} \int_0^{\pi} dx \frac{2x \sin(nx)}{n} \\ &= \frac{4(-1)^n}{n^2}. \end{aligned} \quad (\text{A.3.4})$$

Considering the problem at hand, consider the function as x^2 , so $b_n = 0$ because x^2 is even. The Fourier series then takes the form

$$\begin{aligned} x^2 &= \frac{2\pi^2}{3 \cdot 2} + \sum_{n=1}^{\infty} \frac{4(-1)^n}{n^2} \cos\left(\frac{2n\pi x}{L}\right) \\ &= \frac{\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4(-1)^n}{n^2} (-1)^n. \end{aligned} \quad (\text{A.3.5})$$

Choose x^2 to be equal to 0,

$$\pi^2 = \frac{\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4}{n^2}. \quad (\text{A.3.6})$$

Then subtract the a_0 term and divide by 4

$$\frac{\pi^2}{6} = \sum_{n=1}^{\infty} \frac{1}{n^2} \quad (\text{A.3.7})$$

Complex Integral Way Consider the following integral:

$$I = \int_C \frac{1}{z^2} f(z) dz. \quad (\text{A.3.8})$$

Let $f(z)$ be the Fermi function:

$$f(z) = \frac{1}{1 + e^{i\pi z}}, \quad (\text{A.3.9})$$

then we have several first order poles at $z = \pm(2n + 1) = p_n$ and a second order pole at $z = 0$.

The residue from the 1st order poles:

$$\text{Res}(f(z), p_n) = \lim_{z \rightarrow p_n} (z - p_n) f(z) = \frac{i}{\pi p_n^2}, \quad (\text{A.3.10})$$

and from the 2nd order pole:

$$\text{Res}(f(z), 0) = \lim_{z \rightarrow 0} \frac{d}{dz} (z^2 f(z)) = \frac{-\pi i}{4}. \quad (\text{A.3.11})$$

Writing now Cauchy's residue theorem A.2

$$\int_C g(z) dz = 2\pi i \sum \text{Res}(g(z)) \quad (\text{A.3.12})$$

$$\int_C f(z) dz = 2\pi i \left(\frac{-\pi i}{4} \right) + 2\pi i \sum_{n=0}^{\infty} \frac{2i}{\pi(2n+1)^2} = 0 \quad (\text{A.3.13})$$

$$\frac{\pi i}{4} = \sum_{n=0}^{\infty} \frac{2i}{\pi(2n+1)^2} \quad (\text{A.3.14})$$

$$\frac{\pi^2}{8} = \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \quad (\text{A.3.15})$$

$$= \frac{3}{4} \sum_{n=1}^{\infty} \frac{1}{n^2} \quad (\text{A.3.16})$$

$$\frac{\pi^2}{6} = \sum_{n=1}^{\infty} \frac{1}{n^2} \quad (\text{A.3.17})$$

B Algorithms and Methods

B.1 Monte Carlo and Simulated Annealing

B.2 Minimalization by Nelder-Mead algorithm

C Links, Data and other stuff

D Research Diary

[1], [2]

References

- [1] Anna Andereg, 2022, Example 1.
- [2] Brenda Bradshaw, 2021, Example 2.