

Valid Inequalities for the Cumulative Constraint and the Cumulative Job Shop Scheduling Problem

Tallys Yunes ¹ Dimitris Magos ² Ioannis Mourtos ³

¹ Dept. of Management Science, University of Miami
tallys@miami.edu

² Dept. of Informatics, Technological Educational Institute of Athens
dmagos@teiath.gr

³ Dept. of Mgmt. Science and Technology, Athens Univ. of Econ. & Business
mourtos@aueb.gr

Tepper School of Business, March 25, 2011

Problem Description

Given n jobs (p_j = processing time, c_j =resource consumption) and a single machine with capacity C , the constraint

$$\text{cumulative}((s_1, \dots, s_n), (p_1, \dots, p_n), (c_1, \dots, c_n), C)$$

states that the job start times $s_j \in [r_j, d_j - p_j]$ must be such that the machine capacity is never exceeded


Problem Description

Given n jobs (p_j = processing time, c_j =resource consumption) and a single machine with capacity C , the constraint

$$\text{cumulative}((s_1, \dots, s_n), (p_1, \dots, p_n), (c_1, \dots, c_n), C)$$

states that the job start times $s_j \in [r_j, d_j - p_j]$ must be such that the machine capacity is never exceeded

j	p_j	c_j	r_j	d_j
1	5	1	0	5
2	3	3	0	5
3	4	2	1	7



Problem Description (cont.)

Many applications:

- Production planning and scheduling
- Resource-constrained project scheduling
- Berth allocation at container ports



Problem Description (cont.)

Many applications:

- Production planning and scheduling
- Resource-constrained project scheduling
- Berth allocation at container ports
- Baking (pizza, cakes, cookies,...)



Problem Description (cont.)

Many applications:

- Production planning and scheduling
- Resource-constrained project scheduling
- Berth allocation at container ports
- Baking (pizza, cakes, cookies,...)



As the number of jobs and the time horizon (especially) get larger, optimizing with even a **single** cumulative can be challenging for **both** MIP and CP

A library of single-machine **Cumulative Scheduling Problems**:

CuSPLIB: <http://moya.bus.miami.edu/~tallys/cusplib>

Outline

- Problem Description and Related Work
- Revisiting Identical Jobs
- Arbitrary Resource Consumption
- Cumulative Job-Shop Scheduling
- Preliminary Experiments
- Conclusion

Some Related Work

Mostly disjunctive, rather than cumulative, with a few exceptions:

- [Queyranne and Schulz '95](#): parallel machines with non-stationary speeds; $p_j = 1$; generalization of our problem
- [Hooker and Yan '01](#): facet-defining inequalities for identical jobs; valid inequalities for the general case
- [Hooker '07](#): valid inequalities for the general case
- [Hardin, Nemhauser, and Savelsbergh '08](#): $r_j = 0$, $d_j = \infty$, arbitrary c_j , p_j ; x_{jt} variables; lifted cover-clique inequalities (some computation: 25 instances with 15 jobs each)

Basic Definitions

Jobs are indexed by $N = \{1, \dots, n\}$

r_j^{lb}	: earliest release date
r_j^{ub}	: latest release date
p_j	: processing time
c_j	: resource consumption
C	: capacity of the resource

Basic Definitions

Jobs are indexed by $N = \{1, \dots, n\}$

r_j^{lb}	: earliest release date
r_j^{ub}	: latest release date
p_j	: processing time
c_j	: resource consumption
C	: capacity of the resource

Let s_j denote the start time of job $j \in N$, such that

$$\sum_{j \in N_t} c_j \leq C, \quad \forall t \quad (1)$$

$$r_j^{lb} \leq s_j \leq r_j^{ub} (= d_j - p_j), \quad \forall j \in N \quad (2)$$

where $N_t = \{j \in N : s_j \leq t < s_j + p_j\}$

Identical Jobs

We initially assume that all $j \in N$ are **identical**, i.e.

$$r_j^{lb} = r_0, r_j^{ub} = r_1, p_j = p_0, c_j = c_0$$

Identical Jobs

We initially assume that all $j \in N$ are **identical**, i.e.

$$r_j^{lb} = r_0, r_j^{ub} = r_1, p_j = p_0, c_j = c_0$$

Given $K \subseteq N$, let $s(K) = \sum_{j \in K} s_j$

Identical Jobs

We initially assume that all $j \in N$ are **identical**, i.e.

$$r_j^{lb} = r_0, r_j^{ub} = r_1, p_j = p_0, c_j = c_0$$

Given $K \subseteq N$, let $s(K) = \sum_{j \in K} s_j$

Let $\lambda = \left\lfloor \frac{C}{c_0} \right\rfloor$ = maximum number of jobs running at a time

Identical Jobs

We initially assume that all $j \in N$ are **identical**, i.e.

$$r_j^{lb} = r_0, r_j^{ub} = r_1, p_j = p_0, c_j = c_0$$

Given $K \subseteq N$, let $s(K) = \sum_{j \in K} s_j$

Let $\lambda = \left\lfloor \frac{C}{c_0} \right\rfloor =$ maximum number of jobs running at a time

Let $\rho(K) = \left\lceil \frac{|K|}{\lambda} \right\rceil - 1 =$ periods required to run $|K|$ jobs, minus 1

Identical Jobs

We initially assume that all $j \in N$ are **identical**, i.e.

$$r_j^{lb} = r_0, r_j^{ub} = r_1, p_j = p_0, c_j = c_0$$

Given $K \subseteq N$, let $s(K) = \sum_{j \in K} s_j$

Let $\lambda = \left\lfloor \frac{C}{c_0} \right\rfloor =$ maximum number of jobs running at a time

Let $\rho(K) = \left\lceil \frac{|K|}{\lambda} \right\rceil - 1 =$ periods required to run $|K|$ jobs, minus 1

$$s(K) \geq f(K) = |K|r_0 + p_0\rho(K) \left(|K| - \frac{\lambda}{2}(\rho(K) + 1) \right)$$

Identical Jobs

We initially assume that all $j \in N$ are **identical**, i.e.

$$r_j^{lb} = r_0, r_j^{ub} = r_1, p_j = p_0, c_j = c_0$$

Given $K \subseteq N$, let $s(K) = \sum_{j \in K} s_j$


Let $\lambda = \left\lfloor \frac{C}{c_0} \right\rfloor =$ maximum number of jobs running at a time

Let $\rho(K) = \left\lceil \frac{|K|}{\lambda} \right\rceil - 1 =$ periods required to run $|K|$ jobs, minus 1

$$s(K) \geq f(K) = |K|r_0 + p_0\rho(K) \left(|K| - \frac{\lambda}{2}(\rho(K) + 1)\right)$$


$$s(K) \leq g(K) = |K|r_1 - p_0\rho(K) \left(|K| - \frac{\lambda}{2}(\rho(K) + 1)\right)$$

Graphical Intuition Behind $f(K)$




$$f(K) =$$

Graphical Intuition Behind $f(K)$



$$f(K) = \sum_{i=0}^{\rho(K)-1} (r_0 + ip_0)\lambda$$

Graphical Intuition Behind $f(K)$



$$f(K) = \sum_{i=0}^{\rho(K)-1} (r_0 + ip_0)\lambda + (r_0 + \rho(K)p_0)(|K| - \lambda\rho(K))$$

Cumulative Polyhedra with Identical Jobs

f is supermodular and g is submodular on 2^N

Cumulative Polyhedra with Identical Jobs

f is supermodular and g is submodular on 2^N

The lower problem polyhedron of start times (r_0 finite, $r_1 = \infty$) is the extended contrapolymatroid associated with f :

$$EP(f) = \{s \in \mathbb{R}^n : s(K) \geq f(K), \forall K \subseteq N\}$$

$$B(f) = \{s \in EP(f) : s(N) = f(N)\}$$

Cumulative Polyhedra with Identical Jobs

f is supermodular and g is submodular on 2^N

The **lower** problem polyhedron of start times (r_0 finite, $r_1 = \infty$) is the **extended contrapolymatroid** associated with f :

$$\begin{aligned} EP(f) &= \{s \in \mathbb{R}^n : s(K) \geq f(K), \forall K \subseteq N\} \\ B(f) &= \{s \in EP(f) : s(N) = f(N)\} \end{aligned}$$

The **upper** problem polyhedron start times ($r_0 = -\infty$, r_1 finite) is the **extended polymatroid** associated with g :

$$\begin{aligned} EP(g) &= \{s \in \mathbb{R}^n : s(K) \leq g(K), \forall K \subseteq N\} \\ B(g) &= \{s \in EP(g) : s(N) = g(N)\} \end{aligned}$$

Cumulative Polyhedra with Identical Jobs (cont.)

Theorem

For $j \in N$, $s_j \geq r_0$ define extreme rays of $EP(f)$ and, for $|K| > \lfloor C/c_0 \rfloor$, $s(K) \geq f(K)$ define facets of $EP(f)$

Cumulative Polyhedra with Identical Jobs (cont.)

Theorem

For $j \in N$, $s_j \geq r_0$ define extreme rays of $EP(f)$ and, for $|K| > \lfloor C/c_0 \rfloor$, $s(K) \geq f(K)$ define facets of $EP(f)$

Theorem

$B(f)$ is completely described by $s_j = r_0$ ($j \in N$) and the inequalities $s(K) \geq f(K)$ when $|K| > \lfloor C/c_0 \rfloor$ and either $|K| = |N| - 1$ or

$$\frac{|N|}{\left\lfloor \frac{C}{c_0} \right\rfloor} > \left\lfloor \frac{|K|}{\left\lfloor \frac{C}{c_0} \right\rfloor} \right\rfloor + 1$$

Cumulative Polyhedra with Identical Jobs (cont.)

Theorem

For $j \in N$, $s_j \geq r_0$ define extreme rays of $EP(f)$ and, for $|K| > \lfloor C/c_0 \rfloor$, $s(K) \geq f(K)$ define facets of $EP(f)$

Theorem

$B(f)$ is completely described by $s_j = r_0$ ($j \in N$) and the inequalities $s(K) \geq f(K)$ when $|K| > \lfloor C/c_0 \rfloor$ and either $|K| = |N| - 1$ or

$$\frac{|N|}{\left\lfloor \frac{C}{c_0} \right\rfloor} > \left\lfloor \frac{|K|}{\left\lfloor \frac{C}{c_0} \right\rfloor} \right\rfloor + 1$$

Theorem

*When $r_1 \geq r_0 + \rho(N)p_0$, the convex hull of all feasible schedules $P = EP(f) \cap EP(g)$ is a **generalized polymatroid***

Cumulative Polyhedra with Identical Jobs (cont.)

- Can optimize a linear function of s over $EP(f)$ and $EP(g)$ in polynomial time

Cumulative Polyhedra with Identical Jobs (cont.)

- Can optimize a linear function of s over $EP(f)$ and $EP(g)$ in polynomial time
 - Greedy algorithm:
$$s_{\pi(j)} = f(\{\pi(1), \dots, \pi(j)\}) - f(\{\pi(1), \dots, \pi(j-1)\})$$

Cumulative Polyhedra with Identical Jobs (cont.)

- Can optimize a linear function of s over $EP(f)$ and $EP(g)$ in polynomial time
 - Greedy algorithm:
$$s_{\pi(j)} = f(\{\pi(1), \dots, \pi(j)\}) - f(\{\pi(1), \dots, \pi(j-1)\})$$
 - $O(n \log n)$ separation: sort s_j 's in non-decreasing order

Cumulative Polyhedra with Identical Jobs (cont.)

- Can optimize a linear function of s over $EP(f)$ and $EP(g)$ in polynomial time
 - Greedy algorithm:
$$s_{\pi(j)} = f(\{\pi(1), \dots, \pi(j)\}) - f(\{\pi(1), \dots, \pi(j-1)\})$$
 - $O(n \log n)$ separation: sort s_j 's in non-decreasing order
- **However**, inequalities $s(K) \geq f(K)$ and $s(K) \leq g(K)$ can still be useful for problems with **multiple** cumulative constraints!

Cumulative Polyhedra with Identical Jobs (cont.)

- Can optimize a linear function of s over $EP(f)$ and $EP(g)$ in polynomial time
 - Greedy algorithm:
$$s_{\pi(j)} = f(\{\pi(1), \dots, \pi(j)\}) - f(\{\pi(1), \dots, \pi(j-1)\})$$
 - $O(n \log n)$ separation: sort s_j 's in non-decreasing order
- **However**, inequalities $s(K) \geq f(K)$ and $s(K) \leq g(K)$ can still be useful for problems with **multiple** cumulative constraints!
- Hold that thought...

Cumulative Scheduling with Arbitrary c_j

Cumulative Scheduling with Arbitrary c_j

For all $j \in N$, $r_j^{lb} = r_0$, $r_j^{ub} = \infty$, $p_j = p_0$, **but** c_j arbitrary

Cumulative Scheduling with Arbitrary c_j

For all $j \in N$, $r_j^{lb} = r_0$, $r_j^{ub} = \infty$, $p_j = p_0$, **but** c_j arbitrary

Definition

Given $K \subseteq N$, the ordered tuple $Q = (J_1, \dots, J_q)$, with $J_i \subset K$, is a **feasible partition** of K if

- (i) $\bigcup_{i=1}^q J_i = K$;
- (ii) $J_{i_1} \cap J_{i_2} = \emptyset$, for any $i_1 \neq i_2 \in \{1, \dots, q\}$;
- (iii) $\sum_{j \in J_i} c_j \leq C$, for all $i \in \{1, \dots, q\}$.

Cumulative Scheduling with Arbitrary c_j

For all $j \in N$, $r_j^{lb} = r_0$, $r_j^{ub} = \infty$, $p_j = p_0$, **but** c_j arbitrary

Definition

Given $K \subseteq N$, the ordered tuple $Q = (J_1, \dots, J_q)$, with $J_i \subset K$, is a **feasible partition** of K if

- (i) $\bigcup_{i=1}^q J_i = K$;
- (ii) $J_{i_1} \cap J_{i_2} = \emptyset$, for any $i_1 \neq i_2 \in \{1, \dots, q\}$;
- (iii) $\sum_{j \in J_i} c_j \leq C$, for all $i \in \{1, \dots, q\}$.

If Q satisfies $|J_1| \geq |J_2| \geq \dots \geq |J_q|$, it is called a **decreasing feasible partition**

Cumulative Scheduling with Arbitrary c_j (cont.)

It's feasible to assign $s_j = r_0 + (i - 1)p_0$ for all $J_i \in Q$, which yields

$$\sum_{j \in K} s_j = \sum_{i=1}^q (r_0 + (i - 1)p_0) |J_i| = h(K, Q)$$

Cumulative Scheduling with Arbitrary c_j (cont.)

It's feasible to assign $s_j = r_0 + (i - 1)p_0$ for all $J_i \in Q$, which yields

$$\sum_{j \in K} s_j = \sum_{i=1}^q (r_0 + (i - 1)p_0) |J_i| = h(K, Q)$$

Lemma

Given $K \subseteq N$, the following inequality is valid for the cumulative scheduling polyhedron

$$\sum_{j \in K} s_j \geq \min_{Q \in \mathfrak{P}(K)} h(K, Q)$$

where $\mathfrak{P}(K)$ is the set of all decreasing feasible partitions of K

Facets for Cumulative Scheduling with Arbitrary c_j

$$\sum_{j \in K} s_j \geq \min_{Q \in \mathfrak{P}(K)} h(K, Q) = \min_{Q \in \mathfrak{P}(K)} \sum_{i=1}^{|Q|} (r_0 + (i-1)p_0) |J_i|$$

Theorem

Given $K \subseteq N$, let $Q^* = (J_1, \dots, J_q) = \operatorname{argmin}_{Q \in \mathfrak{P}(K)} h(K, Q)$.
The resulting inequality defines a facet of the cumulative scheduling polyhedron if

- (i) $q \geq 2$
- (ii) $\sum_{j \in J_i} c_j - \min_{j \in J_i} c_j + \min_{j \in J_{i+1}} c_j \leq C, \quad \forall i = 1, \dots, q-1$
- (iii) Either $|J_q| = 1$ or $\sum_{j \in J_q} c_j - \min_{j \in J_q} c_j + c_{j^*} \leq C$,
for some $j^* \in \bigcup_{i=1}^{q-1} J_i$

What Kind of Partition Minimizes $h(K, Q)$?

$$\sum_{j \in K} s_j \geq \min_{Q \in \mathfrak{P}(K)} h(K, Q) = \min_{Q \in \mathfrak{P}(K)} \sum_{i=1}^{|Q|} (r_0 + (i-1)p_0) |J_i|$$

What Kind of Partition Minimizes $h(K, Q)$?

$$\sum_{j \in K} s_j \geq \min_{Q \in \mathfrak{P}(K)} h(K, Q) = \min_{Q \in \mathfrak{P}(K)} \sum_{i=1}^{|Q|} (r_0 + (i-1)p_0) |J_i|$$

Definition

Let $Q = (J_1, \dots, J_q)$ and $Q' = (J'_1, \dots, J'_q)$ be two elements of $\mathfrak{P}(q, K)$. Q' is said to **majorize** Q (denoted $Q' \succ Q$) if $\sum_{i=1}^v |J'_i| \geq \sum_{i=1}^v |J_i|$, for all $v \in \{1, \dots, q-1\}$, and $\sum_{i=1}^q |J'_i| = \sum_{i=1}^q |J_i|$. Moreover, Q' is called a **majorizer** of $\mathfrak{P}(q, K)$ if $Q' \succ Q$ for all $Q \in \mathfrak{P}(q, K)$. The set of all majorizers of $\mathfrak{P}(q, K)$ is denoted by $\mathfrak{P}^\succ(q, K)$

What Kind of Partition Minimizes $h(K, Q)$?

$$\sum_{j \in K} s_j \geq \min_{Q \in \mathfrak{P}(K)} h(K, Q) = \min_{Q \in \mathfrak{P}(K)} \sum_{i=1}^{|Q|} (r_0 + (i-1)p_0) |J_i|$$

Definition

Let $Q = (J_1, \dots, J_q)$ and $Q' = (J'_1, \dots, J'_q)$ be two elements of $\mathfrak{P}(q, K)$. Q' is said to **majorize** Q (denoted $Q' \succ Q$) if $\sum_{i=1}^v |J'_i| \geq \sum_{i=1}^v |J_i|$, for all $v \in \{1, \dots, q-1\}$, and $\sum_{i=1}^q |J'_i| = \sum_{i=1}^q |J_i|$. Moreover, Q' is called a **majorizer** of $\mathfrak{P}(q, K)$ if $Q' \succ Q$ for all $Q \in \mathfrak{P}(q, K)$. The set of all majorizers of $\mathfrak{P}(q, K)$ is denoted by $\mathfrak{P}^\succ(q, K)$

Proposition

Given $K \subseteq N$ and a fixed partition size q , the value of $h(K, Q)$ is minimized by a majorizer of $\mathfrak{P}(q, K)$

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

$Q_2 \succ Q_1$, $h(K, Q_2) = 14$

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

$Q_2 \succ Q_1$, $h(K, Q_2) = 14$

But $Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields $h(K, Q_3) = 13$

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

$Q_2 \succ Q_1$, $h(K, Q_2) = 14$

But $Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields $h(K, Q_3) = 13$

A majorizer may not exist:

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

$Q_2 \succ Q_1$, $h(K, Q_2) = 14$

But $Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields $h(K, Q_3) = 13$

A majorizer may not exist:

$K = \{1, \dots, 9\}$, $C = 8$, $c = (1, 1, 1, 1, 3, 3, 3, 3, 3)$, $q^* = 3$

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

$Q_2 \succ Q_1$, $h(K, Q_2) = 14$

But $Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields $h(K, Q_3) = 13$

A majorizer may not exist:

$K = \{1, \dots, 9\}$, $C = 8$, $c = (1, 1, 1, 1, 3, 3, 3, 3, 3)$, $q^* = 3$

$Q_1 = (\{1, 2, 3, 4, 5\}, \{6, 7\}, \{8, 9\})$

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

$Q_2 \succ Q_1$, $h(K, Q_2) = 14$

But $Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields $h(K, Q_3) = 13$

A majorizer may not exist:

$K = \{1, \dots, 9\}$, $C = 8$, $c = (1, 1, 1, 1, 3, 3, 3, 3, 3)$, $q^* = 3$

$Q_1 = (\{1, 2, 3, 4, 5\}, \{6, 7\}, \{8, 9\})$

$Q_2 = (\{1, 2, 5, 6\}, \{3, 4, 7, 8\}, \{9\})$

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

$Q_2 \succ Q_1$, $h(K, Q_2) = 14$

But $Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields $h(K, Q_3) = 13$

A majorizer may not exist:

$K = \{1, \dots, 9\}$, $C = 8$, $c = (1, 1, 1, 1, 3, 3, 3, 3, 3)$, $q^* = 3$

$Q_1 = (\{1, 2, 3, 4, 5\}, \{6, 7\}, \{8, 9\})$

$Q_2 = (\{1, 2, 5, 6\}, \{3, 4, 7, 8\}, \{9\})$

Neither $Q_1 \succ Q_2$, nor $Q_2 \succ Q_1$

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

$Q_2 \succ Q_1$, $h(K, Q_2) = 14$

But $Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields $h(K, Q_3) = 13$

A majorizer may not exist:

$K = \{1, \dots, 9\}$, $C = 8$, $c = (1, 1, 1, 1, 3, 3, 3, 3, 3)$, $q^* = 3$

$Q_1 = (\{1, 2, 3, 4, 5\}, \{6, 7\}, \{8, 9\})$

$Q_2 = (\{1, 2, 5, 6\}, \{3, 4, 7, 8\}, \{9\})$

Neither $Q_1 \succ Q_2$, nor $Q_2 \succ Q_1$

No majorizer of size 3 exists.

What Kind of Partition Minimizes $h(K, Q)$? (cont.)

Intuitive guess: A majorizer with minimum size q^* ?

Smaller $q \not\Rightarrow$ smaller $h(K, Q)$:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_1 = (\{1, 2, 3, 4, 9\}, \{5, 6, 7, 8, 10\})$, $Q_2 = (\{1, 2, 3, 4, 5, 9\}, \{6, 7, 8, 10\})$

$Q_2 \succ Q_1$, $h(K, Q_2) = 14$

But $Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields $h(K, Q_3) = 13$

A majorizer may not exist:

$K = \{1, \dots, 9\}$, $C = 8$, $c = (1, 1, 1, 1, 3, 3, 3, 3, 3)$, $q^* = 3$

$Q_1 = (\{1, 2, 3, 4, 5\}, \{6, 7\}, \{8, 9\})$

$Q_2 = (\{1, 2, 5, 6\}, \{3, 4, 7, 8\}, \{9\})$

Neither $Q_1 \succ Q_2$, nor $Q_2 \succ Q_1$

No majorizer of size 3 exists.

We suspect it is NP-Hard (GAP), but no proof yet

A Lower Bound on $h(K, Q)$

Let $K = \{1, \dots, |K|\} \subseteq N$

$$\min \sum_{j=1}^{|K|} \sum_{t=1}^{|K|} \frac{(r_0 + (t-1)p_0)}{c_j} y_{jt}$$

$$\sum_{t=1}^{|K|} y_{jt} = c_j, \quad \forall j \in K$$

$$\sum_{j=1}^{|K|} y_{jt} \leq C, \quad \forall t \in K$$

$$y_{jt} \in \{0, c_j\}, \quad \forall j, t \in K$$

where y_{jt} = resource consumption of j at time t

A Lower Bound on $h(K, Q)$

Let $K = \{1, \dots, |K|\} \subseteq N$

$$\begin{aligned} \min \quad & \sum_{j=1}^{|K|} \sum_{t=1}^{|K|} \frac{(r_0 + (t-1)p_0)}{c_j} y_{jt} \\ & \sum_{t=1}^{|K|} y_{jt} = c_j, \quad \forall j \in K \\ & \sum_{j=1}^{|K|} y_{jt} \leq C, \quad \forall t \in K \\ & y_{jt} \in \{0, c_j\}, \quad \forall j, t \in K \end{aligned}$$

where y_{jt} = resource consumption of j at time t

The LP relaxation of this model is a [transportation problem](#)

A Lower Bound on $h(K, Q)$ (cont.)

Recall previous example:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields minimum $h(K, Q_3) = 13$

A Lower Bound on $h(K, Q)$ (cont.)

Recall previous example:

$K = \{1, \dots, 10\}$, $r_0 = 1$, $p_0 = 1$, $C = 10$, $c = (1, 1, 1, 1, 1, 1, 1, 1, 5, 6)$

$Q_3 = (\{1, \dots, 8\}, \{9\}, \{10\})$ yields minimum $h(K, Q_3) = 13$

The **transportation problem relaxation** gives:

$$y_{11} = y_{21} = \dots = y_{81} = 1$$

$$y_{91} = 2, y_{92} = 3$$

$$y_{10,2} = 6$$

For a value of 11.6, which can be rounded up to 12

Generalization of Job-Shop Scheduling

Generalization of Job-Shop Scheduling

Traditional job-shop:

- m **disjunctive** machines, n jobs
- s_{ij} = start time of job j on machine i
- Each job visits (all) machines in a **specific** order

Generalization of Job-Shop Scheduling

Traditional job-shop:

- m **disjunctive** machines, n jobs
- s_{ij} = start time of job j on machine i
- Each job visits (all) machines in a **specific** order

Known Fact: $J3|p_{ij} = 1|C_{\max}$ or $\sum C_j$ is NP-Hard

Generalization of Job-Shop Scheduling

Traditional job-shop:

- m **disjunctive** machines, n jobs
- s_{ij} = start time of job j on machine i
- Each job visits (all) machines in a **specific** order

Known Fact: $J3|p_{ij} = 1|C_{\max}$ or $\sum C_j$ is NP-Hard

Cumulative job-shop:

- m **cumulative** machines, n jobs, same s_{ij} variables
- Each job visits (all) machines in a specific order
- Jobs have identical c_{ij} on each machine, but may differ across machines

Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

Sample instance:

$$m = 5: C_i = (3, 5, 11, 4, 2)$$

$$n = 10: c_i = (1, 2, 3, 2, 1)$$

Machine sequences:

Job 1: 1, 2, 3, 4, 5

Job 2: 5, 1, 3, 2, 4

Job 3: 2, 5, 1, 4, 3

Job 4: 2, 4, 5, 3, 1

Job 5: 3, 5, 4, 1, 2


Job 6: 4, 3, 2, 5, 1

Job 7: 5, 1, 2, 4, 3

Job 8: 1, 3, 2, 5, 4


Job 9: 2, 3, 1, 4, 5

Job 10: 3, 4, 5, 1, 2



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$


How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- Machine sequencing constraints impose initial release dates on jobs: r_{ij}



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- Machine sequencing constraints impose initial release dates on jobs: r_{ij}

Job 1: 1, 2, 3, 4, 5

Job 2: 5, 1, 3, 2, 4

Job 3: 2, 5, 1, 4, 3

Job 4: 2, 4, 5, 3, 1

Job 5: 3, 5, 4, 1, 2


Job 6: 4, 3, 2, 5, 1

Job 7: 5, 1, 2, 4, 3

Job 8: 1, 3, 2, 5, 4

Job 9: 2, 3, 1, 4, 5

Job 10: 3, 4, 5, 1, 2



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- Machine sequencing constraints impose initial release dates on jobs: r_{ij}

Job 1: 1, 2, 3, 4, 5

Job 2: 5, 1, 3, 2, 4

Job 3: 2, 5, 1, 4, 3

Job 4: 2, 4, 5, 3, 1

Job 5: 3, 5, 4, 1, 2


Job 6: 4, 3, 2, 5, 1

Job 7: 5, 1, 2, 4, 3

Job 8: 1, 3, 2, 5, 4

Job 9: 2, 3, 1, 4, 5

Job 10: 3, 4, 5, 1, 2



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- Machine sequencing constraints impose initial release dates on jobs: r_{ij}

Job 1: 1, 2, 3, 4, 5

Job 2: 5, 1, 3, 2, 4

Job 3: 2, 5, 1, 4, 3

Job 4: 2, 4, 5, 3, 1

Job 5: 3, 5, 4, 1, 2

Job 6: 4, 3, 2, 5, 1


Job 7: 5, 1, 2, 4, 3

Job 8: 1, 3, 2, 5, 4

Job 9: 2, 3, 1, 4, 5

Job 10: 3, 4, 5, 1, 2

- At machine i , jobs with $r_{ij} = r$ form class r



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- Machine sequencing constraints impose initial release dates on jobs: r_{ij}

Job 1: 1, 2, 3, 4, 5

Job 2: 5, 1, 3, 2, 4

Job 3: 2, 5, 1, 4, 3

Job 4: 2, 4, 5, 3, 1

Job 5: 3, 5, 4, 1, 2

Job 6: 4, 3, 2, 5, 1


Job 7: 5, 1, 2, 4, 3

Job 8: 1, 3, 2, 5, 4

Job 9: 2, 3, 1, 4, 5

Job 10: 3, 4, 5, 1, 2


- At machine i , jobs with $r_{ij} = r$ form **class r**
- Separate $s(K) \geq f(K)$ for each class



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- If **infeasible**:




Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- If infeasible:

Either 3, 4, or 10 must have $r_{5j} < 2$




Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- If **infeasible**:

Either 3, 4, or 10 must have $r_{5j} < 2$

Say it's job 3



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$


How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- If **infeasible**:

Either 3, 4, or 10 must have $r_{5j} < 2$

Say it's job 3

Branch: $s_{53} \leq 1$ and $s_{53} \geq 2$



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:


- If **infeasible**:

Either 3, 4, or 10 must have $r_{5j} < 2$

Say it's job 3

Branch: $s_{53} \leq 1$ and $s_{53} \geq 2$

Update: $r_{53}^{ub} = 1$ and $r_{53}^{lb} = 2$



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- If **infeasible**:


Either 3, 4, or 10 must have $r_{5j} < 2$

Say it's job 3

Branch: $s_{53} \leq 1$ and $s_{53} \geq 2$

Update: $r_{53}^{ub} = 1$ and $r_{53}^{lb} = 2$

- Branching alters job classes: allows cuts to remove violations



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- If **infeasible**:


Either 3, 4, or 10 must have $r_{5j} < 2$

Say it's job 3

Branch: $s_{53} \leq 1$ and $s_{53} \geq 2$

Update: $r_{53}^{ub} = 1$ and $r_{53}^{lb} = 2$

- Branching alters job classes: allows cuts to remove violations
- Branching imposes upper bounds on s_{ij} :
separate $s(K) \leq g(K)$ as well



Cumulative Job-Shop: $p_{ij} = 1, \min \sum s_{ij}$

How to use $s(K) \geq f(K)$ and $s(K) \leq g(K)$ cuts:

- If **infeasible**:


Either 3, 4, or 10 must have $r_{5j} < 2$

Say it's job 3

Branch: $s_{53} \leq 1$ and $s_{53} \geq 2$

Update: $r_{53}^{ub} = 1$ and $r_{53}^{lb} = 2$

- Branching alters job classes: allows cuts to remove violations
- Branching imposes upper bounds on s_{ij} :
separate $s(K) \leq g(K)$ as well
- How to pick violation and job to branch on?
Still experimenting with this...



Before we continue...

Before we continue...



Before we continue...

Potential Branching Heuristic:

Before we continue...

Potential Branching Heuristic:

Have generic problem \mathcal{P}

Before we continue...

Potential Branching Heuristic:

Have generic problem \mathcal{P}

Have family \mathcal{F} of cuts
that work when \mathcal{P} is restricted
to satisfy certain conditions \mathcal{R}

Before we continue...

Potential Branching Heuristic:

Have generic problem \mathcal{P}

Have family \mathcal{F} of cuts
that work when \mathcal{P} is restricted
to satisfy certain conditions \mathcal{R}

Bias branching on \mathcal{P} to bring it
closer to satisfying \mathcal{R} during search

Before we continue...

Potential Branching Heuristic:

Have generic problem \mathcal{P}

Have family \mathcal{F} of cuts
that work when \mathcal{P} is restricted
to satisfy certain conditions \mathcal{R}

Bias branching on \mathcal{P} to bring it
closer to satisfying \mathcal{R} during search

Separate \mathcal{F} cuts when violated

Before we continue...

Potential Branching Heuristic:

Have generic problem \mathcal{P}

Have family \mathcal{F} of cuts
that work when \mathcal{P} is restricted
to satisfy certain conditions \mathcal{R}

Bias branching on \mathcal{P} to bring it
closer to satisfying \mathcal{R} during search

Separate \mathcal{F} cuts when violated

Preliminary Computational Experiments

Initial set of instances generated by hand:

- Number of machines $m = 5$ and $C_i = (3, 5, 11, 4, 2)$
- Number of jobs $n \in \{10, 15, 20, 25, 30, 35, 40, 50, 60\}$
- Resource consumption $c_i = (1, 2, 3, 2, 1)$
- $p_{ij} = 1$ and time horizon $T = 30$

Preliminary Computational Experiments

Initial set of instances generated by hand:

- Number of machines $m = 5$ and $C_i = (3, 5, 11, 4, 2)$
- Number of jobs $n \in \{10, 15, 20, 25, 30, 35, 40, 50, 60\}$
- Resource consumption $c_i = (1, 2, 3, 2, 1)$
- $p_{ij} = 1$ and time horizon $T = 30$

Solution methods:

- Default CPLEX 12.2 with typical x_{ijt} formulation

Preliminary Computational Experiments

Initial set of instances generated by hand:

- Number of machines $m = 5$ and $C_i = (3, 5, 11, 4, 2)$
- Number of jobs $n \in \{10, 15, 20, 25, 30, 35, 40, 50, 60\}$
- Resource consumption $c_i = (1, 2, 3, 2, 1)$
- $p_{ij} = 1$ and time horizon $T = 30$

Solution methods:

- Default CPLEX 12.2 with typical x_{ijt} formulation
- B&C with s_{ij} variables, classed f and g cuts, and 2 branching strategies:

Preliminary Computational Experiments

Initial set of instances generated by hand:

- Number of machines $m = 5$ and $C_i = (3, 5, 11, 4, 2)$
- Number of jobs $n \in \{10, 15, 20, 25, 30, 35, 40, 50, 60\}$
- Resource consumption $c_i = (1, 2, 3, 2, 1)$
- $p_{ij} = 1$ and time horizon $T = 30$

Solution methods:

- Default CPLEX 12.2 with typical x_{ijt} formulation
- B&C with s_{ij} variables, classed f and g cuts, and 2 branching strategies:
 - EFC = Earliest violation, first candidate job
 - ESDFF = Earliest viol., smallest s_{ij} domain, fractional s_{ij} **first**

Preliminary Computational Experiments

Initial set of instances generated by hand:

- Number of machines $m = 5$ and $C_i = (3, 5, 11, 4, 2)$
- Number of jobs $n \in \{10, 15, 20, 25, 30, 35, 40, 50, 60\}$
- Resource consumption $c_i = (1, 2, 3, 2, 1)$
- $p_{ij} = 1$ and time horizon $T = 30$

Solution methods:

- Default CPLEX 12.2 with typical x_{ijt} formulation
- B&C with s_{ij} variables, classed f and g cuts, and 2 branching strategies:
 - EFC = Earliest violation, first candidate job
 - ESDFF = Earliest viol., smallest s_{ij} domain, fractional s_{ij} **first**
- **Latest** violation seems to perform worse than earliest

n	Optimal Solution	CPLEX Default			CPLEX 1-Thread		
		Best	Nodes	Time	Best	Nodes	Time
10	54		1	0.1		1	0.1
15	98		4K	57		733	7
20	154		12K	122		3K	26
25	225		51K	4K		17K	817
30	306		83K	11K		233K	79K
35	402	414	131K	$2K^m$	405	161K	$43K^m$
40	508	525	91K	$4K^m$	516	108K	$8K^m$
50	760*	791	114K	$3K^m$	784	89K	$15K^m$
60	1,062*	<i>no sol.</i>	90K	$13K^m$	<i>no sol.</i>	67K	$44K^m$

n	Optimal Solution	CPLEX Default			CPLEX 1-Thread		
		Best	Nodes	Time	Best	Nodes	Time
10	54		1	0.1		1	0.1
15	98		4K	57		733	7
20	154		12K	122		3K	26
25	225		51K	4K		17K	817
30	306		83K	11K		233K	79K
35	402	414	131K	2K ^m	405	161K	43K ^m
40	508	525	91K	4K ^m	516	108K	8K ^m
50	760*	791	114K	3K ^m	784	89K	15K ^m
60	1,062*	<i>no sol.</i>	90K	13K ^m	<i>no sol.</i>	67K	44K ^m

n	EFC			ESDFF		
	Best	Nodes	Time	Best	Nodes	Time
10		15	0.1		15	0.1
15		393K	2K		10K	31
20		221	1		8K	30
25		119K	25K		1K	14
30		2K	32		11K	392
35		38K	26K		21K	6K
40		14K	724		7K	395
50	762	159K	24h ^t	762	77K	24h ^t
60	1,097	66K	24h ^t	1108	68K	24h ^t

Preliminary Computational Experiments (cont.)

Modified job-shop instances from OR-Library

- 81 instances taken from 6 articles from 1988 to 1992
- Number of jobs $n \in \{10, 15, 20, 30, 50\}$
- Number of machines $m \in \{5, 10, 15, 20\}$
- **Modifications:**
 - $p_{ij} = 1$
 - $C = 6$ for all machines
 - c_i can be 2 or 3 (randomly chosen)

Preliminary Computational Experiments (cont.)

Modified job-shop instances from OR-Library

- 81 instances taken from 6 articles from 1988 to 1992
- Number of jobs $n \in \{10, 15, 20, 30, 50\}$
- Number of machines $m \in \{5, 10, 15, 20\}$
- **Modifications:**
 - $p_{ij} = 1$
 - $C = 6$ for all machines
 - c_i can be 2 or 3 (randomly chosen)

Solution methods:

- Default CPLEX 12.2 with typical x_{ijt} formulation

Preliminary Computational Experiments (cont.)

Modified job-shop instances from OR-Library

- 81 instances taken from 6 articles from 1988 to 1992
- Number of jobs $n \in \{10, 15, 20, 30, 50\}$
- Number of machines $m \in \{5, 10, 15, 20\}$
- **Modifications:**
 - $p_{ij} = 1$
 - $C = 6$ for all machines
 - c_i can be 2 or 3 (randomly chosen)

Solution methods:

- Default CPLEX 12.2 with typical x_{ijt} formulation
- B&C with s_{ij} variables, classed f and g cuts, and 4 branching strategies:

Preliminary Computational Experiments (cont.)

Modified job-shop instances from OR-Library

- 81 instances taken from 6 articles from 1988 to 1992
- Number of jobs $n \in \{10, 15, 20, 30, 50\}$
- Number of machines $m \in \{5, 10, 15, 20\}$
- **Modifications:**
 - $p_{ij} = 1$
 - $C = 6$ for all machines
 - c_i can be 2 or 3 (randomly chosen)

Solution methods:

- Default CPLEX 12.2 with typical x_{ijt} formulation
- B&C with s_{ij} variables, classed f and g cuts, and 4 branching strategies:
 - EFC = Earliest violation, first candidate job
 - ESDFF = Earliest viol., smallest s_{ij} domain, fractional s_{ij} **first**
 - ESDFL = Earliest viol., smallest s_{ij} domain, fractional s_{ij} **last**
 - SODE = Smallest overall s_{ij} domain, earliest viol. first if tied

Instances with $n = 10$: 5 with $m = 5$, 18 with $m = 10$

Instances with $n = 10$: 5 with $m = 5$, 18 with $m = 10$

All methods found all 23 optimal solutions

Instances with $n = 10$: 5 with $m = 5$, 18 with $m = 10$

All methods found all 23 optimal solutions

# inst.	CPLEX		B&C best		B&C avg.	
	Nodes	Time	Nodes	Time	Nodes	Time

Instances with $n = 10$: 5 with $m = 5$, 18 with $m = 10$

All methods found all 23 optimal solutions

	# inst.	CPLEX		B&C best		B&C avg.	
		Nodes	Time	Nodes	Time	Nodes	Time
$m = 5$	4	1	0.10	37	0.08	same	same
	1	1	0.09	1,079	1.7	same	same

Instances with $n = 10$: 5 with $m = 5$, 18 with $m = 10$

All methods found all 23 optimal solutions

	# inst.	CPLEX		B&C best		B&C avg.	
		Nodes	Time	Nodes	Time	Nodes	Time
$m = 5$	4	1	0.10	37	0.08	same	same
	1	1	0.09	1,079	1.7	same	same
$m = 10$	16	32	0.52	82	0.25	97	0.28
	2	423	6.70	57,163	140	81,951	212

Instances with $n = 15$: 5 for each $m = 5, 10$, and 15

Instances with $n = 15$: 5 for each $m = 5, 10$, and 15

CPLEX found all 15 optimal solutions

B&C methods found all 15, but didn't prove one of them

Instances with $n = 15$: 5 for each $m = 5, 10$, and 15

CPLEX found all 15 optimal solutions

B&C methods found all 15, but didn't prove one of them

# inst.	CPLEX		B&C best		B&C avg.	
	Nodes	Time	Nodes	Time	Nodes	Time

Instances with $n = 15$: 5 for each $m = 5, 10$, and 15

CPLEX found all 15 optimal solutions

B&C methods found all 15, but didn't prove one of them

	# inst.	CPLEX		B&C best		B&C avg.	
		Nodes	Time	Nodes	Time	Nodes	Time
$m = 5$	4	243	1.05	189	0.56	270	0.76
	1	41	0.77	146,162	388	166,955	446

Instances with $n = 15$: 5 for each $m = 5, 10$, and 15

CPLEX found all 15 optimal solutions

B&C methods found all 15, but didn't prove one of them

	# inst.	CPLEX		B&C best		B&C avg.	
		Nodes	Time	Nodes	Time	Nodes	Time
$m = 5$	4	243	1.05	189	0.56	270	0.76
	1	41	0.77	146,162	388	166,955	446
$m = 10$	4	310	2.89	639	2.52	676	2.64
	1	600	2.95	508,659 ^m		509,876 ^m	

Instances with $n = 15$: 5 for each $m = 5, 10$, and 15

CPLEX found all 15 optimal solutions

B&C methods found all 15, but didn't prove one of them

	# inst.	CPLEX		B&C best		B&C avg.	
		Nodes	Time	Nodes	Time	Nodes	Time
$m = 5$	4	243	1.05	189	0.56	270	0.76
	1	41	0.77	146,162	388	166,955	446
$m = 10$	4	310	2.89	639	2.52	676	2.64
	1	600	2.95	508,659 ^m		509,876 ^m	
$m = 15$	4	194	2.51	208	1.21	211	1.22
	1	150	1.94	73,346	366	75,161	375

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

# Instances	$m = 5$	$m = 10$	$m = 15$	$m = 20$	Total
Tied					
CPLEX better					
ESDFF better					
Total					28

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

# Instances	$m = 5$	$m = 10$	$m = 15$	$m = 20$	Total
Tied			1 ^{<i>a</i>}		1
CPLEX better					
ESDFF better					
Total					28

^{*a*}: neither method found optimum

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

# Instances	$m = 5$	$m = 10$	$m = 15$	$m = 20$	Total
Tied			1 ^a		1
CPLEX better	3 ^b				
ESDFF better					
Total					28

^a: neither method found optimum

^b: one 20x faster; one optimum ESDFF didn't find

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

# Instances	$m = 5$	$m = 10$	$m = 15$	$m = 20$	Total
Tied			1 ^a		1
CPLEX better	3 ^b	10 ^c			
ESDFF better					
Total		10			28

^a: neither method found optimum

^b: one 20x faster; one optimum ESDFF didn't find

^c: one non-optimal: loses to EFC: value of 269 vs. 266

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

# Instances	$m = 5$	$m = 10$	$m = 15$	$m = 20$	Total
Tied			1 ^a		1
CPLEX better	3 ^b	10 ^c	4 ^d		
ESDFF better					
Total		10			28

^a: neither method found optimum

^b: one 20x faster; one optimum ESDFF didn't find

^c: one non-optimal: loses to EFC: value of 269 vs. 266

^d: one 3x faster; one optimum found by neither

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

# Instances	$m = 5$	$m = 10$	$m = 15$	$m = 20$	Total
Tied			1 ^a		1
CPLEX better	3 ^b	10 ^c	4 ^d	1	18
ESDFF better					
Total		10			28

^a: neither method found optimum

^b: one 20x faster; one optimum ESDFF didn't find

^c: one non-optimal: loses to EFC: value of 269 vs. 266

^d: one 3x faster; one optimum found by neither

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

# Instances	$m = 5$	$m = 10$	$m = 15$	$m = 20$	Total
Tied			1 ^a		1
CPLEX better	3 ^b	10 ^c	4 ^d	1	18
ESDFF better	3 ^e				
Total	6	10			28

^a: neither method found optimum

^b: one 20x faster; one optimum ESDFF didn't find

^c: one non-optimal: loses to EFC: value of 269 vs. 266

^d: one 3x faster; one optimum found by neither

^e: two 10x faster

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

# Instances	$m = 5$	$m = 10$	$m = 15$	$m = 20$	Total
Tied			1 ^a		1
CPLEX better	3 ^b	10 ^c	4 ^d	1	18
ESDFF better	3 ^e		3 ^f		
Total	6	10	8		28

^a: neither method found optimum

^b: one 20x faster; one optimum ESDFF didn't find

^c: one non-optimal: loses to EFC: value of 269 vs. 266

^d: one 3x faster; one optimum found by neither

^e: two 10x faster

^f: two optima found by neither

Instances with $n = 20$: $m \in \{5, 10, 15, 20\}$, 28 total

No method ran out of time; always found opt. or feasible solutions

# Instances	$m = 5$	$m = 10$	$m = 15$	$m = 20$	Total
Tied			1 ^a		1
CPLEX better	3 ^b	10 ^c	4 ^d	1	18
ESDFF better	3 ^e		3 ^f	3 ^g	9
Total	6	10	8	4	28

^a: neither method found optimum

^b: one 20x faster; one optimum ESDFF didn't find

^c: one non-optimal: loses to EFC: value of 269 vs. 266

^d: one 3x faster; one optimum found by neither

^e: two 10x faster

^f: two optima found by neither

^g: all 2x faster

Instances with $n = 30$, $m = 10$

Originally: S. Lawrence, *GSIA Tech. Report*, 1984

Instances with $n = 30$, $m = 10$

Originally: S. Lawrence, *GSIA Tech. Report*, 1984

Inst.	CPLEX			EFC		ESDFF		ESDFL		SODE	
	Best	Nodes	Time	Best	Nodes	Best	Nodes	Best	Nodes	Best	Nodes
la31	411	132K	24h	426	192K	428	203K	428	189K	-	191K
la32	402*	130K	38K	418	196K	415	194K	425	179K	409	190K
la33	410	132K ^m	45K	411	192K	407	186K	407	351K	412	189K
la34	409	138K ^m	76K	401	193K	399	534K	404	190K	400	195K
la35	392*	124K	52K	419	191K	419	188K	420	192K	420	192K

All four B&C algorithms ran out of memory (4GB)

Instances with $n = 50$, $m = 10$

Originally: Storer, Wu, and Vaccari, *Management Science*, 1992

Instances with $n = 50$, $m = 10$

Originally: Storer, Wu, and Vaccari, *Management Science*, 1992

Inst.	CPLEX			ESDFF			
	Best	Nodes	Time	Best	Nodes	Time	TTF
swv11	-	58K ^m	7,015	1115 (10.3%)	93K	24h	43
swv12	-	59K ^m	8,565	1206 (16.25%)	91K	24h	41
swv13	-	54K ^m	16,530	1108 (9.84%)	98K	24h	15
swv14	-	55K ^m	12,510	1119 (11.53%)	74K	24h	265
swv15	-	62K ^m	2,904	1118 (10.29%)	91K	24h	45
swv16	-	61K ^m	2,734	-	140K ^m	-	-
swv17	-	54K ^m	16,961	967 (8.79%)	116K ^m	-	51
swv18	-	56K ^m	12,082	960 (6.25%)	134K ^m	-	16,759
swv19	-	51K ^m	55,676	940 (5.74%)	98K ^m	-	288
swv20	-	51K ^m	38,033	934 (6.10%)	136K ^m	-	1,402

TTF = time till first feasible solution (in seconds)

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes
- Inequalities remain valid (though not strong) for general case

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes
- Inequalities remain valid (though not strong) for general case
 - use $r_0 = \min r_j$, $p_0 = \min p_j$, $c_0 = \min c_j$ in $f(K)$

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes
- Inequalities remain valid (though not strong) for general case
 - use $r_0 = \min r_j$, $p_0 = \min p_j$, $c_0 = \min c_j$ in $f(K)$
- Facet-defining inequalities for arbitrary c_j (untested)

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes
- Inequalities remain valid (though not strong) for general case
 - use $r_0 = \min r_j$, $p_0 = \min p_j$, $c_0 = \min c_j$ in $f(K)$
- Facet-defining inequalities for arbitrary c_j (untested)
- Preliminary experiments show some promise

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes
- Inequalities remain valid (though not strong) for general case
 - use $r_0 = \min r_j$, $p_0 = \min p_j$, $c_0 = \min c_j$ in $f(K)$
- Facet-defining inequalities for arbitrary c_j (untested)
- Preliminary experiments show some promise
 - especially as n increases

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes
- Inequalities remain valid (though not strong) for general case
 - use $r_0 = \min r_j$, $p_0 = \min p_j$, $c_0 = \min c_j$ in $f(K)$
- Facet-defining inequalities for arbitrary c_j (untested)
- Preliminary experiments show some promise
 - especially as n increases
- **To do:** find heuristic separation for arbitrary- c_j facets

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes
- Inequalities remain valid (though not strong) for general case
 - use $r_0 = \min r_j$, $p_0 = \min p_j$, $c_0 = \min c_j$ in $f(K)$
- Facet-defining inequalities for arbitrary c_j (untested)
- Preliminary experiments show some promise
 - especially as n increases
- To do: find heuristic separation for arbitrary- c_j facets
- To do: continue with cumulative job-shop experiments

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes
- Inequalities remain valid (though not strong) for general case
 - use $r_0 = \min r_j$, $p_0 = \min p_j$, $c_0 = \min c_j$ in $f(K)$
- Facet-defining inequalities for arbitrary c_j (untested)
- Preliminary experiments show some promise
 - especially as n increases
- To do: find heuristic separation for arbitrary- c_j facets
- To do: continue with cumulative job-shop experiments
- To do: find other applications with multiple cumulative constraints

Conclusions and Future Work

- Complete characterization of simple cumulative polytopes
- Inequalities remain valid (though not strong) for general case
 - use $r_0 = \min r_j$, $p_0 = \min p_j$, $c_0 = \min c_j$ in $f(K)$
- Facet-defining inequalities for arbitrary c_j (untested)
- Preliminary experiments show some promise
 - especially as n increases
- **To do:** find heuristic separation for arbitrary- c_j facets
- **To do:** continue with cumulative job-shop experiments
- **To do:** find other applications with multiple cumulative constraints
- **To do:** allow other parameters to differ: r_j , p_j

The End

Thank You!

Any Questions?