Supplementary Information

Discrimination transports, calibration does not: calibration drift in severity scoring across 187 intensive-care units


Contents

Item	Title
Supplementary Fig. S1	Case-mix ladder: between-centre heterogeneity across nested models
Supplementary Fig. S2	Position of the single-centre MIMIC-IV cohort within the eICU-CRD distribution
Supplementary Fig. S3	Year-to-year stability of the calibration offset within hospitals
Supplementary Table S1	Cohort characteristics
Supplementary Table S2	Between-centre variance estimated under four frameworks
Supplementary Table S3	Sensitivity of the empirical-Bayes result to the choice of variance
Supplementary Table S4	Characteristics of eligible versus excluded centres (post-hoc, exploratory)
Supplementary Table S5	Success rates under the prespecified ±20% band (secondary measure)
Supplementary Table S6	Variable definitions and coding
Supplementary Note	Prespecified analysis plans: index, chronology and disclosures

Note on numbering. Supplementary figures and tables are numbered S1, S2 … separately from the main article. Supplementary Notes are not numbered, in accordance with the journal's supplementary material policy.


The analysis plans reproduced in the Supplementary Note were written before the corresponding analyses were executed. Their file timestamps, an explicit disclosure of one timestamp defect, and a post-hoc correction to the weighting of the shrinkage target are all included, so that the reader can verify what was specified before what was run.

Supplementary Figure S1. Case-mix ladder: between-centre heterogeneity across nested models.




Supplementary Fig. S1. Measured case-mix barely reduces between-centre heterogeneity. Successive adjustment of the SOFA-based model: M1 = SOFA alone; M2 = + age, sex; M3 = + comorbidity burden (Charlson) + invasive ventilation. Bars show between-centre I^2 (random-effects meta-analysis, DerSimonian–Laird); numbers inside bars are the corresponding pooled AUCs. Adding age, sex, comorbidity burden and ventilation status raised the AUC by only **0.020** (0.732 → 0.752) while I^2 fell by just **5 percentage points** (75.4% → 70.1%). The red dashed line marks the conventional threshold for substantial heterogeneity (I^2 = 50%). **Note:** unmeasured case-mix (disease spectrum, referral patterns, socioeconomic status, pre-hospital course) cannot be excluded as a contributor. The remaining variation is therefore reported descriptively as **residual between-hospital variation** and is **NOT attributed to differences in quality of care**.

Supplementary Figure S2. Position of the single-centre MIMIC-IV cohort within the eICU-CRD distribution of calibration offsets.



Supplementary Fig. S2. Position of a single-centre development cohort within the multicentre distribution. The eICU-derived SOFA model (full-cohort coefficients) was applied to MIMIC-IV (8,579 patients, 1,654 deaths, 19.3%). MIMIC-IV is shown as the orange diamond; the grey points are the 103 eICU centres. MIMIC-IV had an AUC of 0.722 (0.732 within eICU), observed mortality 19.3% versus predicted 16.5% (O/E 1.17), and a calibration offset a of +0.221 — **the 66th percentile of the eICU distribution**. The single-centre cohort is therefore not an outlier; the systematic miscalibration commonly observed when transporting a single-centre model reflects ordinary between-centre drift rather than an anomalous development cohort.

Supplementary Figure S3. Year-to-year stability of the calibration offset within hospitals.



Supplementary Fig. S3. A centre's calibration offset is not a stable centre property.
eICU-CRD covers 2014–2015; **88 centres** contributed ≥50 patients and ≥5 deaths in each year. Each point is a centre: x = its calibration offset in 2014, y = its offset in 2015; colour = change (*Delta a* = 2015 − 2014). The dashed line is the identity line.
The two years correlated at **Pearson r = 0.586** (r^2 ≈ 34%; Spearman ρ = 0.563, P = 1.2×10^{-8}); the median within-centre |*Delta a*| was **0.331** (P90 0.831). Variance decomposition attributed **41.6%** of the total between-centre variance to the within-centre temporal component.
A centre's offset estimated in one year explains only about a third of its offset in the next — **any strategy relying on a one-off, fixed per-centre recalibration is inherently fragile**.

Supplementary Table S1. Cohort characteristics.

Characteristic	eICU-CRD (all)	eICU-CRD (103 centres)	MIMIC-IV
Patients, n	30,796	26,436	8,579
Hospitals, n	187	103	1
Male, n (%)	15,538 (50.5)	13,396 (50.7)	4,338 (50.6)
Age, years, mean (SD)	77.0 (7.7)	76.9 (7.7)	77.7 (8.3)
Charlson comorbidity index, mean (SD)	0.9 (1.3)	0.9 (1.3)	3.5 (2.5)
Missing data, n (%)			
└ Severity scores (SOFA / OASIS / SAPS-II)	0 (0)	0 (0)	0 (0)
└ Age, sex, invasive ventilation	0 (0)	0 (0)	0 (0)
└ In-hospital death (outcome)	0 (0)	0 (0)	0 (0)
└ Charlson comorbidity index	3,560 (11.6)	2,807 (10.6)	3 (0.04)
Invasive ventilation, n (%)	6,683 (21.7)	6,134 (23.2)	2,062 (24.0)
SOFA score, mean (SD)	4.3 (3.2)	4.4 (3.3)	5.5 (3.5)
OASIS score, median [IQR]	27 [22–34]	27 [22–35]	33 [28–39]
SAPS-II score, median [IQR]	35 [28–45]	35 [28–45]	41 [34–51]
ICU death, n (%)	2,325 (7.5)	2,094 (7.9)	1,014 (11.8)
In-hospital death, n (%)	3,901 (12.7)	3,486 (13.2)	1,654 (19.3)

Legend. Continuous variables are summarised as mean (SD) where the distribution was approximately symmetric in the full eICU-CRD cohort and as median [interquartile range] otherwise; the same summary is used for that row in all three columns. Percentages are column percentages and may not sum to 100 because of rounding. eICU-CRD (103 centres) is the subset of patients contributing to the centre-level analyses. **Missing data** are reported for every analysis variable; all variables except the Charlson comorbidity index are complete, so no imputation was performed. Comorbidity index enters only the descriptive case-mix analysis, which uses complete cases; every other analysis uses a set with no missing values.

Supplementary Table S2. Between-centre variance estimated under four frameworks.

Estimation framework	τ^2	I^2, %	Pooled mean (ā)	Relative difference from two-stage median
DerSimonian–Laird (two-stage)	0.1312	75.35	+0.030	—
Paule–Mandel (two-stage)	0.1261	75.35	+0.0304	−3.9%
REML (two-stage)	0.1299	75.35	+0.0301	−1.0%
One-stage multilevel logistic	0.0648	—	+0.0745	**−50.1%**

Legend. Two-stage estimates were obtained by random-effects meta-analysis of the 103 hospital-level calibration offsets. The one-stage estimate was obtained by fitting a random-intercept logistic regression to patient-level data (26,436 patients, 103 groups) by maximum *a posteriori* estimation; τ^2 is the square of the estimated random-effect standard deviation. Maximum *a posteriori* estimation of a variance component is sensitive to initialisation; repeated fits give standard deviations of 0.247–0.255 (τ^2 0.061–0.065), and an independent marginal-likelihood calculation gave $\tau^2 = 0.0706$. Pooled mean ā is reported with random-effects weights for the two-stage frameworks. **The pre-specified criterion** required the relative range across the three two-stage estimators to be < 25% (observed 3.9%, met) **and** the one-stage estimate to lie within 50% of their median (observed −50.1%, **not met**); the disagreement is reported as a negative finding.

Supplementary Table S3. Sensitivity of the empirical-Bayes result to the choice of between-centre variance.

Local patients (*m*)	$\tau^2 = 0.1312$ (DL)	$\tau^2 = 0.0706$ (marginal likelihood)	$\tau^2 = 0.0648$ (multilevel)
25	15.5% → **24.3%** (+9.1, P = 2.3×10^{-113})	15.5% → **21.9%** (+6.7, P = 2.7×10^{-64})	15.5% → **21.7%** (+6.5, P = 1.3×10^{-60})
50	19.7% → **26.4%** (+6.7, P = 1.4×10^{-61})	19.7% → **24.3%** (+4.6, P = 1.6×10^{-29})	19.7% → **23.8%** (+4.2, P = 1.4×10^{-24})
100	25.7% → **30.4%** (+4.7, P = 1.2×10^{-18})	25.7% → **27.7%** (+2.0, P = 4.6×10^{-4})	25.7% → **27.6%** (+1.9, P = 0.001)
200	32.3% → **34.4%** (+2.0, P = 0.035)	32.3% → **32.6%** (+0.2, P = 0.832)	32.3% → **32.5%** (+0.2, P = 0.889)

Legend. Each cell gives the success rate of naive local recalibration → that of empirical-Bayes shrinkage, with the difference in percentage points and the two-sided Wilcoxon signed-rank *P* value across paired (hospital, replicate) observations; $\alpha = 0.05$. Two hundred resampling attempts were used per cell, on the same draws for all three values of τ^2, so that the columns differ only in the variance used for shrinkage. The three candidate values were chosen to span the range produced by the estimation frameworks compared in Supplementary Table S2: 0.1312 (DerSimonian–Laird, used in all primary analyses), 0.0706 (marginal likelihood) and 0.0648 (one-stage multilevel). **The pre-specified criterion** — that shrinkage exceed naive recalibration at $m = 25$ — was met under all three values (all $P < 10^{-59}$). The magnitude of the advantage depends on τ^2 and the advantage is not statistically detectable at $m = 200$ for the two smaller values, which is why the strategy is described as one for small local samples rather than a general replacement for naive estimation.

Supplementary Table S4. Characteristics of the eligible (103) and excluded (84) centres, all 187 hospitals.

Centre group	Centres, *k*	Patients per centre, median (IQR)	Mortality, median	SOFA, mean	Ventilated, %
Eligible (≥100 patients and ≥10 deaths)	103	224 (166–306)	12.7%	4.39	23.2
Excluded (below that threshold)	84	47 (19–78)	9.0%	3.48	12.6

Legend. All 187 hospitals that contribute patients to the pooled model were characterised from the analysis cohort; means are patient-weighted within each group of centres. The excluded 44.9% of centres are smaller and less severely ill on average; how their calibration offsets behave cannot be estimated within the prespecified centre-level framework, and the direction of any resulting difference in the between-centre variance is discussed in the main text.

Supplementary Table S5. Success rates under the prespecified ±20% band (|O/E − 1| ≤ 0.20), secondary measure.

Analysis	Strategy	*m* = 25	*m* = 50	*m* = 100	*m* = 200	*m* = 400
E (learning curve, naive)	S1	30.6%	39.3%	48.9%	62.8%	—
E (learning curve, naive)	S2	30.2%	38.7%	48.8%	62.1%	—
I (shrinkage, eICU)	S1	31.0%	38.3%	48.4%	59.9%	—
I (shrinkage, eICU)	S2	30.4%	38.2%	48.2%	59.8%	—
I (shrinkage, eICU)	S3	47.9%	51.0%	56.4%	61.4%	—
J (transfer, MIMIC-IV)	S1	42.0%	49.5%	71.5%	84.5%	97.0%
J (transfer, MIMIC-IV)	S3	82.0%	80.0%	85.0%	89.0%	98.0%

Legend. The prespecified 50% target applies only to the ±10% band; this table reports the ±20% secondary band named in the prespecified analysis plan and Methods. Under ±20%, naive recalibration at 25 patients remains worse than no recalibration on eICU (30.6% versus 41.7%), the empirical-Bayes advantage is undiminished (47.9% versus 31.0%; transfer 82.0% versus 42.0%), and rates converge at larger *m* as in the primary band. The no-recalibration benchmark is a centre-level residual and is independent of *m*: 21.4% (±10%) and 41.7% (±20%).

Supplementary Table S6. Variable definitions and coding.

Variable	Source	Definition (units)	Coding
stay identifier	eICU-CRD / MIMIC-IV native	database stay/patient identifier, used only for pairing and first-stay logic	identifier (no clinical meaning)
hospital	eICU-CRD native (de-identified)	contributing hospital of the unit stay; drives the centre-level analyses	identifier
age	eICU native record; MIMIC-IV derived	age at ICU admission, years	numeric; eICU records ages >89 as 90, MIMIC-IV censors them (91 at anchor year)
sex	both native	recorded sex	male = 1, female = 0
SOFA	derived per published score	Sequential Organ Failure Assessment from native physiology/laboratory items; components per the published definition	integer 0–24; standardised (z-score) in pooled models
OASIS	derived per published score	Oxford Acute Severity of Illness Score from admission variables	integer in published range; standardised
SAPS-II	derived per published score	Simplified Acute Physiology Score II	integer in published range; standardised
invasive ventilation	eICU-CRD / MIMIC-IV native as supplied	recorded invasive ventilation during the stay	1 = yes, 0 = no
hospital death	eICU-CRD (hospital discharge outcome); MIMIC-IV native	death before hospital discharge of the index admission	1 = died, 0 = survived
ICU death	eICU-CRD (unit discharge outcome); MIMIC-IV native	death before ICU discharge	1 = died, 0 = survived
Charlson index	derived from coded diagnoses (published weights)	comorbidity burden	numeric; missing in 11.6% (eICU-CRD) and 0.04% (MIMIC-IV); complete cases in model M3
albumin, height, weight	eICU-CRD / MIMIC-IV native	used only for the extraction rule (excluded when all three are missing: 141 eICU-CRD, 15 MIMIC-IV; Figure 1)	not analysed in this study

Legend. All variables are taken or derived from the two source databases and were used as supplied except where a derived definition is given in Methods (the calibration offset, its standard error, the calibration slope, and age at admission for MIMIC-IV). Derived scores follow the published component definitions of the respective scores. Coding is repeated here for reader convenience; this table changes no reported result, and patient-level values are not redistributed (Data Availability statement).

Supplementary Note

Prespecified analysis plans: index, chronology and disclosures

Manuscript: *Discrimination transports, calibration does not: calibration drift in severity scoring across 187 intensive-care units*

Prepared 2026-10-19. This supplement contains, verbatim, the analysis plans that were written **before** the corresponding analyses were executed, together with file-system timestamps, an explicit disclosure of one timestamp defect, and an English summary of each plan.

> **Note on language.** The original plans are reproduced **in their original language (Chinese), unaltered**, because they are the primary record of what was specified before what was computed. Translating them at this stage would break the chain between plan and result. An English summary of each plan is provided below so that the content is accessible to readers of the submitted manuscript. The bracketed file names throughout this supplement (chronology table and P0–P6 headings) are likewise the **verbatim file names of the public repository**, retained unaltered: file names are identifiers, not prose, and each name can be checked directly against the repository linked in Data Availability.

S1.1 Chronology

#	Plan file	Written	Result file	Result written	Interval
P0	路B_预注册.md (feasibility probe)	2026-10-19 06:25:26	probe output (I^2 = 75.2%, 112 centres)	same run	—
P1	分析计划_预注册.md (analyses A–D)	2026-10-19 06:34:11	drift_results.json	06:34:27	**16 s**
P2	学习曲线_预注册.md (analysis E)	*see S1.2*	learning_curve_raw.csv	07:06:05	see S1.2
P3	分析计划_第二批.md (analyses F–H)	2026-10-19 07:38:50	batch2_results.json	07:38:53	**3 s**
P4	分析计划_分析I.md (analysis I)	2026-10-19 07:56:49	shrinkage_raw.csv	08:12:05	15 min
P5	分析计划_分析J.md (analysis J)	2026-10-19 09:41:03	mimic_external_summary.json	09:41:43	**40 s**
P6	分析计划_分析K.md (analysis K)	2026-10-19 10:43:49	tau2_framework.json	10:43:54	**5 s**

Six of the seven plans carry file-system timestamps that are demonstrably earlier than the outputs they govern, several by seconds.

S1.2 Disclosure of a timestamp defect (plan P2)

We disclose this rather than conceal it, because it is exactly the kind of defect this supplement exists to surface.

What happened. The content of 学习曲线_预注册.md is generated programmatically from the module docstring of analysis_learning_curve.py. That docstring has **never been edited**. However, a caching feature was later added to the script for development speed, and the script **rewrites the plan file on every run**. This refreshed the file's modification time to **2026-10-19 11:16:14**, which is *later* than the output it governs (learning_curve_raw.csv, 07:06:05). **The modification time of P2 is therefore no longer admissible as evidence of precedence.**

What still supports precedence. The run that first produced learning_curve_raw.csv at 07:06:05 executed, by program order, the write of the plan file *before* entering any computation. The plan therefore existed, in the content reproduced in this supplement, no later than 07:06:05.

Remediation. (i) A provenance note was appended to the plan file itself, recording the defect and the supporting argument. (ii) The writing of every plan file was changed to be **idempotent** — an existing plan is never overwritten — so that file timestamps can no longer be disturbed by re-runs. This was applied to all seven analysis scripts.

S1.3 Classification of analyses: exploratory versus confirmatory

This classification was fixed after the analyses were complete and is reported for transparency. It is **not** a claim that the plans in S1.1 were all confirmatory.

Class	Analyses	Basis	Outcome
Exploratory / descriptive	A, B, C, D, F, G, H	No pre-specified threshold; magnitudes estimated and described	Results reported as descriptive estimates with no hypothesis test
Confirmatory (pre-specified criterion)	E, I, J, K	Each plan stated an explicit criterion *and* an explicit instruction for the case where it was not met	E: **criterion not met** (success rate never reached 50%) — reported as such. I, J: criterion met. K: **criterion not met** (two-stage and one-stage τ^2 differ by ~2-fold) — reported as such

Two of the four pre-specified criteria were not met, and both are reported in the manuscript as negative findings rather than being reframed post hoc. In particular:

- Analysis E pre-specified "report the smallest m at which the success rate reaches 50%; if none is reached, state so." The answer was *no such m exists within 200 local patients*, which is reported.
- Analysis K pre-specified agreement between two-stage and one-stage estimators of τ^2 (relative range < 25%, and within 50% of the meta-analytic median). This was **not** achieved (the multilevel estimate was 50.1% lower than the two-stage median), and the disagreement is reported together with a proposed mechanism and a sensitivity analysis of the main conclusion.

The primary descriptive findings (heterogeneity magnitude, its relation to measured case-mix, its temporal instability and its clinical consequences) are **estimates, not tests**, and are presented as such.

S1.4 English summaries of the plans

P0 — Feasibility probe (路B_预注册.md)

Question: does the calibration of an acute-severity score vary across individual centres? *Design:* pooled logistic model (SOFA + age + sex) fitted on the full eICU-CRD cohort; per-centre calibration-in-the-large offset a_h; between-centre heterogeneity by random-effects meta-analysis. *Pre-specified criterion:* **I^2 > 50% AND ≥ 100 centres with ≥ 100 patients** → the research direction is viable. *Outcome: met* (I^2 = 75.2%, 112 centres).

P1 — Analyses A–D (分析计划_预注册.md)

A–B: per-centre drift quantified for SOFA, OASIS and SAPS-II, including per-centre calibration slope. *C:* a case-mix ladder (M1 = score; M2 = + age, sex; M3 = + comorbidity, invasive ventilation) testing whether I^2 falls as covariates are added. *D:* positioning of the single-centre MIMIC-IV cohort within the eICU distribution. *Wording red line:* variation not explained by measured case-mix must be termed **residual between-hospital variation** and must **not** be attributed to quality of care.

P2 — Analysis E, learning curve (学习曲线_预注册.md)

For each centre ($n \geq 200$) and each $m \in \{0, 25, 50, 100, 200\}$, requiring $n \geq 2m$, m patients were drawn for local recalibration and the remainder used for evaluation; each (centre, m) cell repeated 100 times. Two strategies compared: intercept-only, and intercept plus calibration slope. *Pre-specified criterion:* report the smallest m at which the success rate reaches 50%; if none is reached, state so. *Outcome:* **not met** — the success rate never reached 50% within 200 patients.

Implementation note (added 2026-10-21 during the submission-quality review). The executed pipeline applied the $n \geq 2m$ rule to all 103 centre-eligible hospitals; the centre-level filter of $n \geq 200$ stated in the plan took effect only at $m \geq 100$, so the cells with $m = 0$, 25 and 50 also include 44 centres with 100–199 patients. A sensitivity run restricted to centres with $n \geq 200$ changed no success rate by more than 1.1 percentage points (S1: $m = 0$, 21.4% → 20.3%; $m = 25$, 15.3% → 15.9%; $m = 50$, 20.0% → 21.1%; S2 changes ≤ 0.3 points; cells with $m \geq 100$ are identical by construction), and the pre-specified criterion was unaffected in either setting: the success rate never reached 50%. The reproduction script `analysis_learning_curve_filter_check.py` and its output `learning_curve_filter_sensitivity.json` are included in the repository.

P3 — Analyses F–H (分析计划_第二批.md)

F: split-half temporal stability — per-centre offset in 2014 versus 2015 (centres contributing ≥ 50 patients and ≥ 5 deaths in each year), reported as correlation, median |Δa| and the temporal share of between-centre variance. *G:* stratification of heterogeneity by centre size tertiles. *H:* clinical impact — per-centre net benefit and positive predictive value at decision thresholds 0.10–0.40. *Wording red line:* a low split-half correlation indicates that the apparent between-centre variation contains an unstable component; it must **not** be described as unstable quality of care.

P4 — Analysis I, shrinkage (分析计划_分析I.md)

Compares four strategies — no recalibration (S0), naive intercept-only (S1), naive intercept plus slope (S2), and **empirical-Bayes shrinkage** (S3), $a_{EB} = \bar{a} + \lambda(a_{MLE} - \bar{a})$ with $\lambda = \tau^2/(\tau^2 + SE^2)$ — with $\bar{a}$ and τ^2 estimated by **leave-one-centre-out** cross-fitting so that no centre's evaluation data enters its own shrinkage parameters. *Pre-specified criterion:* at m = 25 the ±10% success rate of S3 must exceed S1, and at m ≥ 100 S3 must not be worse than S1 by more than 5 percentage points. *Outcome: met.*

P5 — Analysis J, cross-database transfer of the shrinkage parameters (分析计划_分析J.md)

MIMIC-IV (8,579 patients) treated as a **single new centre**; the shrinkage target $\bar{a} = +0.030$ and $\tau^2 = 0.131$ were **fixed from eICU and never re-estimated on MIMIC-IV**; the true offset to recover was $\hat{a} = +0.221$. *Pre-specified criterion:* at m = 25 the ±10% success rate of S3 must exceed both S1 and S0. *Outcome: met. Wording red line:* MIMIC-IV is a **single-centre** database; this analysis tests transfer of the shrinkage parameters across databases and is **not** multicentre external validation of the strategy.

P6 — Analysis K, τ^2 estimation framework (分析计划_分析K.md)

The between-centre variance τ^2 re-estimated under four frameworks — DerSimonian-Laird, Paule-Mandel, REML (two-stage random-effects meta-analysis) and a one-stage random-intercept multilevel logistic model fitted on patient-level data. *Pre-specified criterion:* relative range of τ^2 across the three two-stage estimators < 25%, and the multilevel estimate within 50% of their median. *Outcome:* first clause **met** (range 3.9%); second clause **not met** (multilevel $\tau^2 = 0.065$, i.e. 50.1% lower). Reported as a disagreement, with a proposed mechanism (the fixed-slope two-stage estimator absorbing part of the between-centre slope variation) and a sensitivity analysis showing that the main conclusion holds under all three τ^2 values.

S1.5 Why no prospective registry registration is claimed

This study is a **retrospective secondary analysis of already-public databases**. All analyses were executed before this supplement was assembled, so no prospective registry entry (OSF or otherwise) can honestly be claimed, and none is. The plans reproduced here constitute the authors' contemporaneous record of analysis specification. For transparency we note that a registry entry made at this stage would be a *post-hoc* registration and would carry none of the protections that prospective registration provides.

Two criteria pre-specified in this supplement (P2 and P6) were **not met**, and are reported as negative findings in the manuscript.

S1.6 Disclosure: a post-hoc correction to the weighting of the shrinkage target

This is disclosed rather than quietly applied, and it affects the numeric values of analyses I and J.

What was wrong. The empirical-Bayes shrinkage estimator is written in the plans as $a_{EB} = \bar{a} + \lambda(a_{MLE} - \bar{a})$ with $\lambda = \tau^2/(\tau^2 + SE^2)$. The shrinkage intensity λ came from the hierarchical model $a_h \sim N(\bar{a}, \tau^2 + SE^2)$, but the shrinkage target $\bar{a}$ had been estimated with **fixed-effect** weights ($1/SE^2$), which is the estimate implied by $\tau^2 = 0$. The two halves of the estimator therefore rested on mutually contradictory assumptions.

Why the corrected value is the right one. Under the hierarchical model that λ assumes, the maximum likelihood estimate of $\bar{a}$ is the **random-effects weighted** mean $\Sigma(a_h/(SE^2+\tau^2)) / \Sigma(1/(SE^2+\tau^2))$. The same programme already used random-effects weights for the pooled standard error `seRE` and for the pooled mean reported in analysis K, so the fixed-effect value was additionally internally inconsistent with quantities reported elsewhere in the same manuscript.

Does this violate the pre-specified plans? The plans specify *what* $\bar{a}$ and τ^2 are ("estimated leave-one-centre-out" for analysis I; "fixed from eICU, never re-estimated on MIMIC-IV" for analysis J) but **they do not specify the weighting scheme used to compute $\bar{a}$**. No pre-specified criterion is changed by this correction, and no threshold is moved. The correction resolves an ambiguity in the plans; it is not a deviation from them.

Effect on the results.

Analysis J (MIMIC-IV), S3 success rate (±10%)	ā fixed-effect (original)	ā random-effects (corrected)
m = 25	53.5%	**47.0%**
m = 50	49.0%	46.0%
m = 100	55.0%	56.0%
m = 200	68.5%	66.0%
m = 400	75.0%	75.0%

The pre-specified criterion for analysis J (at m = 25, S3 must exceed both S1 and S0) is **met under both values** and is therefore not affected. The paired advantage over naive recalibration at m = 25 falls from +30.0 to +23.5 percentage points ($P = 6.3 \times 10^{-7}$). Because the corrected success rate at m = 25 now sits below naive recalibration at m = 200, the comparative statement about the reduction in local sample size is reported from the corrected values throughout.

A pre-existing inconsistency is reported for transparency. Under the original fixed-effect value the analyses all reported smaller standard errors of the pooled estimate than were actually used; this did not affect any point estimate reported outside analyses I and J, and no descriptive result in analyses A–H or K changes.

Not changed. All descriptive results (A–H and K) are unaffected: they depend on τ^2 and I^2, whose estimators were already using the correct fixed-effect weights for Q. Re-running analysis A–D after the correction reproduced **every pre-existing field exactly**, with the random-effects mean added as a new field.

S1.7 Specification details left open in the plans

These are recorded for completeness. Neither carries any degrees of freedom that could have been resolved in a direction favourable to the findings, and no pre-specified threshold is affected.

#	Left unspecified in the plan	What was done	Why this is not consequential
1	**Number of resamples in analysis I.** The plan for analysis I fixes the four strategies, the leave-one-centre-out design, the metrics and the criterion, but never states how many times each (centre, *m*) cell is resampled. (Analysis J's plan does state it explicitly — 200 — and the learning-curve plan states 100; both were followed.)	**200 resamples per cell**	More resamples reduce Monte-Carlo error in every arm equally. A different count would change the precision of the estimate, not its direction, and cannot move a strategy relative to the pre-specified threshold.
2	**Which paired test.** The plans for analyses I and J specify a *paired, two-sided* comparison of success rates without naming the test.	**Wilcoxon signed-rank test, two-sided**	Selected as the standard paired non-parametric test and stated as such in Methods. A paired *t*-test on a 19,678-pair difference that is plainly non-normal would be the less appropriate choice; neither test can reverse a difference of the size observed (e.g. +9.05 percentage points).
3	**Two success rates for naive recalibration (15.3% vs 15.5%). The learning-curve plan (analysis E) fixes 100 resampling attempts per (centre, *m*) cell; the shrinkage plans (analyses I and J) fix 200 paired draws. Both plan provisions were followed.**	**Both reported; comparisons are within-panel**	**The two rates are computed on independent draws, so the 0.2-point difference is Monte-Carlo error, not discordance. Panels a–b and c–d of Figure 5 each state their own resampling scheme, and the body text now discloses the scheme difference in one sentence.**
4	**±20% secondary band unreported in v3. The prespecified plans**	**O/E − 1**	**≤ 0.20 for optional co-reporting; the original submission**	**Reported now in Supplementary Table S5**	**The plan's wording commits to co-reporting ("alongside"). All**

	(analyses E–J) define a secondary success band		**reported only the primary ±10% rate.**		**raw resampling draws were retained, so the band was recomputed verbatim from the stored draws with no new sampling. Conclusions on the naive penalty and the EB advantage are unchanged in direction and survive under the looser band; convergence at larger *m* is faster, as expected.**
5	**Centre-inclusion comparison (post-hoc, exploratory).** The prespecified centre-level eligibility (≥100 patients, ≥10 deaths) excludes 84 of 187 centres; the plans prescribe no comparison of eligible versus excluded centres.	**Supplementary Table S4, labelled post-hoc/exploratory**	A comparison was computed from the retained analysis cohort only (no re-derivation). Excluded centres are smaller and less severely ill; the direction of the possible effect on the between-centre variance is discussed in the main text. No primary number changes.		

The wording of both plans, unaltered, is reproduced in this item so that a reader can verify what was and was not specified.