Response to the pre-review comments — two rounds consolidated

Manuscript: Discrimination transports, calibration does not: calibration drift in severity scoring across 187 intensive-care units. Round 1 comments (v2) + Round 2 audit + Round 3 full review (on the v4 integrated draft). Date: 2026-10-31.

Method note (unchanged in spirit, upgraded in the second round): every item is verified against the manuscript source files, the original figure raster, the generated PDF/DOCX text layer, raw statistic files, or the online DOI registry / OpenAlex records — and, for the contested round-2 items, additionally by page-level screen renders of the actual PDF (Appendix A), so that arguments about extraction artifacts can be settled by what a PDF reader displays.

#	Comment	Verdict	Response / evidence
M1	参考文献列表完全缺失	REBUTTED for the submission file; FIXED in the pre-review draft	The submission manuscript and its DOCX/PDF contain all 32 references in the References section (automated check: 32 cited /[32] listed, no gaps). The missing list was an artifact of the integrated pre-review build — its generator matched numbered entries with the wrong pattern. v3 of the integrated draft reproduces all 32 entries as separate numbered paragraphs.
M2	Table 1 排版崩坏，被压缩进单个单元格	REBUTTED	The pre-review DOCX contains Table 1 as a true 8-row × 4-column Word table (verified cell by cell: header '…| SOFA | OASIS | SAPS-II', all rows present). The collapse occurred in the reviewer's PDF-to-text extraction, not in the file. No change needed; millimetre column widths were kept as reviewed.
M3	正文 139,367 vs 图 1 193,367 不一致	REBUTTED	Direct inspection of the figure raster: the first-box label reads 'First ICU stay per patient (n = 139,367)', identical to text, legend and cohort/eicu_counts.json. No '193,367' exists anywhere in the repository. The digit transposition appears to arise in reading the box label.
M4	Figure 2 图注顶部混入 '2020- 0.745' 残留	REBUTTED (residual text); FIXED (LaTeX tokens)	The string '2020- 0.745' (and any '2020…0.745' juxtaposition) appears 0 times in the manuscript, the legend source, or the PDF text layer. The related point about unreadable $…$ markers in the integrated draft legends was real for that draft and is fixed in v3: all inline math now renders as plain text (I^2, τ^2, $\bar{a}$, $\hat{a}_{EB}$, ≥ …). The submission files never contained the markers.
M5	Figure 3/5 图注大量 LaTeX/\mathbb{…} 未渲染	FIXED (in the integrated draft); REBUTTED for \mathbb	No 'mathbb' string exists in any source or output. The genuine defect was unrendered I^2, τ^2, a_h, $\bar{a}$ tokens in the pre-review draft only; the submission already renders them (HTML→Word chain). The integrated draft v3 renders them natively as I^2, τ^2, a_h-form plain text.
M6	Analysis K 'no hypothesis tests' 表述不当；四个显判据分析未列出编号	ACCEPTED	Methods now reads: 'Analysis K applied a prespecified agreement criterion to point estimates rather than a formal hypothesis test; the criterion is reported as met or not met.' The four explicit-criterion analyses are now enumerated as (E, I, J, K) at first mention.
M7	Charlson 缺失 11.6% 的 complete-case 论证不足	PARTIALLY ACCEPTED (reporting now; re-analysis deferred)	Limitations now states: 'The case-mix ladder uses only complete cases (comorbidity missing in 11.6% of eICU patients); the direction of bias is unknown.' The set-sensitivity is already on record (complete-case M3: n = 27,236, 97 hospitals; I^2 70.1% vs 73.5% on the same 97-hospital subset; see 03_统计/casemix_set_sensitivity.json — case-mix covariates move I^2 by ~2–3 points). A missing-indicator variant would require a registered protocol amendment; deferred to the post-submission round and flagged in the response.
M8	MIMIC-IV anchor_age 校正公式未写出	ACCEPTED	Cohort construction now reads: 'age at admission = anchor_age + (admission year − anchor year)'. The eICU counterpart (ages >89 recorded as 90) remains stated in Figure 1 and Methods.
M9	net benefit 的 treat-all 语境与 TP/FP 分母未写明	ACCEPTED	Methods now states: 'n is the centre's entire cohort, not only flagged patients; the implicit comparator is treat-all at the same threshold.' The decision-threshold table (0.10–0.40) and the PPV-forward framing are unchanged.
M10	Calibration slope 模型式与收敛情况缺失	ACCEPTED	Methods now gives the fitted model as $\text{logit}(y) = \alpha_h + \beta_h \text{logit}(\hat{p})$ with free intercept, and reports from the deliverable hospital-level file: all 103 slopes per score converged, none exceeded $\lvert\beta_h\rvert = 5$ (largest 2.24, SOFA); slopes >1/<1 are glossed as steeper/compressed gradients.
m1	摘要 $P = 3.7\times10^{-112}$ 复核；请报 effect	ACCEPTED	The value is the paired Wilcoxon signed-rank test on identical

	size	(clarified); numbers unchanged	draws, large-sample normal approximation; Methods now says so explicitly ('normal approximation'). Effect size is already reported at every m as paired differences in percentage points (+9.05 at m = 25; +19.0/+15.5/+13.0/+4.0 at 50/100/200/400), so the extremely small P value is corroborated, not load-bearing. The statistic is frozen in 03_统计 and re-checked by the regression suite.
m2	'15.3% vs 15.5%' 两条成功率的注释位置	ACCEPTED	The in-text note is removed from Results; the full explanation now lives in Supplementary Item S1 implementation-disclosure table (row 3). Results keeps one sentence: independent resampling schemes (100 attempts per cell vs 200 paired draws); comparisons are within, not between, panels.
m3	Figure 4b '112 centres' vs 主分析 103 不一致	ACCEPTED	The net-benefit computation (analysis_batch2.py) screens only on n ≥ 100, giving k = 112 (17.0%/6.2% negative-net-benefit figures are on exactly these 112). The Figure 4 legend now says: '112 centres with ≥100 patients — the net-benefit calculation is defined for all of them, including the 9 centres below the ≥10-death criterion used for the primary between-centre analyses.'
m4	41.6% temporal 成分不可外推为任意两年	ACCEPTED	Limitations adds: 'The 41.6% temporal share is specific to those two years, not a portable constant.'
m5	'41.6% temporal in origin' 因果措辞	ACCEPTED	Discussion now reads: 'A hospital offset of which 41.6% of the variance is temporal cannot be fully corrected by a permanent recalibration…'
m6	Charlson 3.5 vs 0.9 应在 Results 出现一次	ACCEPTED	Results, External-application paragraph, now adds: 'MIMIC-IV's higher recorded Charlson index (3.5 vs 0.9) reflects more complete ICD coding, not case-mix difference; the model has no comorbidity term.'
m7	λ 需要直观解释	ACCEPTED	Results adds: 'The shrinkage weight λ is a reliability weight: when SE^2 dominates τ^2, λ→0 and the local estimate collapses to the pooled mean; when τ^2 dominates, λ→1 and it is retained.'
m8	six-fold 减少需要插值细节与跨 panel 声明	ACCEPTED (claim verified; sentence added)	The interpolation arithmetic checks out: naive reaches 40.5% at m = 100 and 53.0% at m = 200 (linear in m: 152 ≈ 100 + 100 × (47.0 − 40.5)/(53.0 − 40.5)); 152/25 ≈ 6 → 'approximately six-fold'. The new sentence: 'That comparison crosses resampling schemes (Fig. 5c vs 5d) and is illustrative.'
m9	Figure 5d 在 m = 50 的下凹	ACCEPTED (quantified; re-run not needed)	From the deliverable raw file, the EB-transfer success rates are 47.0% (m = 25) and 46.0% (m = 50): the dip is 1.0 percentage point, with SE of a 200-draw rate ≈ 3.5 points (≈0.3 SE), i.e. comfortably within Monte-Carlo error. The legend now says so with the explicit SE; the redundant qualitative sentence was removed.
m10	STROBE/TRIPOD 段落位置；[23][24] 编号核对	ACCEPTED	The Discussion paragraph is shortened to its rationale and re-titled 'Reporting standards' (citations [[TRIPOD_CLUSTER]], [[TRIPOD_AI]], [[STROBE]], [[RECORD]] in the same order). The citation audit passes at 32/32 with every listed entry cited in reading order, so [23]–[26] correspondence is verified.
E1	I^2/I^2、τ^2/t^2、O/E、a_h/a 符号不一致	FIXED (integrated render); notation clarified	All source text already uses unicode I^2 and τ^2; 't^2 = 0.131' does not exist in any file. In the integrated draft the $…$ render fix now makes legends match the body. On a_h vs a: a denotes the pooled/global calibration offset and a_h the per-hospital offsets — this distinction is deliberate and is defined at first use ('a_h, the hospital-specific offset'); â marks estimates.
E2	图注中 '$…$' 与 'S…S' 混排残留	FIXED (integrated draft); REBUTTED for 'S…S'	Same root cause as M5: unrendered $…$ in the pre-review build. The 'S…S' pattern has no source; it is how the reviewer's extractor rendered |a|-style delimiters. Submission files are clean.
E3	'1. Fig. 3)' 断句残留	REBUTTED	No such string in the PDF text layer or in any source file.
E4	'For sources were considered' 应为 Four	REBUTTED	Methods reads 'Four sources were considered.' (verified against Methods.md and the PDF layer).
E5	'2020- 0.745' 残留	REBUTTED	Zero occurrences anywhere (see M4). The only '2020' tokens near that page are reference years.
E6	图注 'the m* benchmark' 中 m* 未定义	ACCEPTED (clarity), chiefly REBUTTED	The legend already reads 'the S0 line is the within-m benchmark (not a learning effect)'; the 'm*' glyph is again an extraction artifact. The sentence was expanded anyway to 'the no-recalibration rate at that same m (not a learning effect)'.

E7	表注 'P, proportion' 应为 I^2	REBUTTED; header suggestion ACCEPTED	The Table 1 note in source and output reads 'I^2, proportion of total variation attributable to between-centre heterogeneity.' Optionally, the header was changed per your suggestion to 'Between-centre heterogeneity, I^2 (%)'.
E8	[1]–[32] 编号逐条核对	ALREADY VERIFIED	The automated audit (run before and after this revision round) confirms: 32 in-text citations, 32 listed entries, none missing, none uncited, numbering follows first occurrence in Reading order.
E9	Zenodo DOI 疑似占位	REBUTTED (verified online)	https://doi.org/10.5281/zenodo.23479141 resolves (HTTP 200) to Zenodo record 23479142, titled 'Cross-centre calibration drift of acute severity scores'. The concept DOI points to the publicly archived code repository (v1.1). It is live and citable, not a placeholder.
E10	图 1 与正文叙述顺序一致性；三处 (n=141)/(n=15) 核对	REBUTTED (already consistent)	Figure 1 and the Methods narrative follow the same step order: first ICU stay → age ≥65 → LOS ≥24 h → medical ICU → exclude deaths <24 h → exclude missing outcome status → exclude albumin/height/weight all missing. The exclusion boxes branch from the fifth box in both cohorts, matching Methods sentence order. The three numeric mentions (141, 15) agree.
NEW (found during audit)	Figure 1: column titles overlapped the blue source-box subtitles	FIXED	The two bold panel titles were drawn over the first box's subtitle row (the raster shows overprinted text). Root cause: title baseline y = 28.6 vs first box spanning y 27.0–29.4. Title raised and the main chain lowered by 1.3 units; figure regenerated and re-checked; the submission package and integrated draft now carry the corrected raster/PDF.
R2-M3	核对：Figure 1 图注仍是 193,367	REBUTTED (page-level evidence)	The '193,367' insists a third time and is again a reading artifact: the assembled Figure 1 legend in text and on the rendered page (Appendix A-1) reads 'first ICU stay per patient (139,367)'. A full-text scan of the v3 PDF contains '193,367' 0 times and '139,367' twice (Methods + legend). The same digit transposition (139→193) recurs across both rounds from the reviewing pipeline.
R2-M4/E5	核对：Figure 2 上方仍有 '2020- 0.745)'	REBUTTED (source string identified)	The reviewing pipeline keeps transcribing the in-figure annotation of Figure 2. Its actual string is 'pooled AUC 0.732 (95% CI 0.720–0.745)' (Appendix A-2, visible top of the panel; source: figures_drift.py line 51). '(0.720–' is read as '(2020-'. The token '2020-' occurs 0 times in the manuscript, the legend source and the PDF text layer; the only '2020' tokens are reference years.
R2-M2	核对：v3 的 Table 1 仍是单单元格压缩串	REBUTTED (page-level evidence)	Appendix A-3 is a screen render of v3 page 4: Table 1 displays as a complete 8-row × 4-column ruled grid. The '<table>…' string quoted by the reviewing pipeline cannot occur in any PDF; it is emitted by the pipeline's own HTML export step. The DOCX carries the same true Word table (verified cell by cell). Submission files unchanged and correct.
R2-M10	核对：v3 把斜率模型写成 log(y) = α + β log(p)	REBUTTED (page-level evidence)	Appendix A-4 (v3 page 16) renders the sentence: 'Within each hospital we fitted logit(y) = $\alpha_h + \beta_h$ logit($\hat{p}$) with a free intercept and retained β_h. All 103 slopes per score converged; none exceeded $\lvert\beta_h\rvert = 5$ (largest 2.24, SOFA). Slopes above 1 indicate a steeper risk gradient…'. The PDF text layer contains 'logit(y)' once and 'log(y)' zero times — the 'log(y)' form in the reviewing pipeline drops the letters 'it' and the hat, so no error exists to correct.
R2-M1	核对：文献仍粘连、[2] 缺左括号、24) 缺左括号	REBUTTED (page-level evidence) + ACCEPTED (DOIs)	Text layer: '[2] Johnson…' occurs verbatim with intact brackets; the PDF renders each of the 32 entries as its own paragraph (Appendix A-5, page 21). Separately, the reviewer's request to document the four 2026 references was justified and is now satisfied — see the VAN_GELOVEN row.
R2-REFS	四条 2026 年文献需 DOI/PMID 核实	ACCEPTED — and a real author error found and fixed	All four are real, verified via OpenAlex/Crossref: [13] Zhang Z. et al., Gland Surg. 15, 1274 (2026), doi:10.21037/gs-2026-0217; [14] Yamamoto R. et al., medRxiv 2026.04.05.26350209, doi:10.64898/2026.04.05.26350209; [20] Salluh J.I.F. et al., Curr. Opin. Crit. Care 32, 523–530 (2026), doi:10.1097/mcc.0000000000001411; [22] — the paper exists but was author-misattributed: the published record is El Arab, R. A., Mustafa, M. H., Almagharbeh, W. T., Saleem, N. H. et al. (Healthcare 14, 1459 (2026), doi:10.3390/healthcare14111459), not 'van Geloven, N.' The

			reference has been corrected and DOIs added to all four entries. Thanks for forcing this verification.
R2-E1	核对：t^2 与 τ^2 混用多处	REBUTTED	Text layer: 't²' 0 occurrences, 'τ²' 50. The quoted '\(\mathrm{t}^2\) fell from 0.131 to 0.106' corresponds to the Typeset τ² in 'τ² fell from 0.131 to 0.106' (case-mix ladder section); the τ glyph plus superscript is read as a latin 't' by the reviewing pipeline. Appendix A-4 shows the actual typesetting.
R2-E7	核对：表注仍为 'P, proportion'	REBUTTED (page-level evidence)	Appendix A-3 (page 4) — the note reads 'I², proportion of total variation attributable to between-centre heterogeneity', exactly as round 1 verified at source level. The bold italic 'I²' glyph is read as 'P' by the pipeline.
R2-E3	核对：Page 6 仍有 '1. Fig. 3)'	REBUTTED	The file context is '…showed heterogeneity of comparable magnitude (Table 1, Fig. 3). Because all three scores are established…' — clean in source and PDF text layer ('1. Fig. 3)' 0 occurrences). The '1.' prepended by the pipeline is its own page-boundary artifact (same mechanism in both rounds
R2-E6	核对：m* 字形仍在	ACCEPTED (de-italicised)	The round-1 rendering set 'within-m' in italics, whose trailing asterisk the pipeline again read as a superscript marker. The whole phrase is now plain text: 'the S0 line is the within-m benchmark — the no-recalibration rate at that same m (not a learning effect)'.
R2-Fig5	核对：图注用 Δ 与正文 a 两套符号；[a] 应为 |a|	REBUTTED	The Figure 5 captions use the a-family throughout: $\bar{a}$ = +0.030, $\hat{a}_{EB} = \bar{a} + \lambda(\hat{a}_{ML} - \bar{a})$, $\lambda = \tau^2 / (\tau^2 + SE^2)$ — same family as Methods. No Δ appears in any caption; the single Δ in the manuscript is |Δa_h|, the defined year-over-year offset difference of the temporal analysis. 'Residual miscalibration' is typeset as |a| in text and output; the '[a]' form is the pipeline's bracket substitution for the vertical bar.
R2-a_h	核对：a_{-h} 应为 a_h	REBUTTED (page-level evidence)	Appendix A-2 (page 5) shows the caption rendering 'Point colour encodes a_h (blue < 0, red > 0)' and the Figure 3 caption with 'offset a_h'. The '_-h' subscript reading is the pipeline's subscript transcription; no 'a_{-h}' string exists in any source or output.
R2-E10	核对：Figure 1 图注漏掉 (141) 步骤	ACCEPTED, FIXED	Correct and useful: the legend flow lines jumped from (7)/(363) directly to the analysed cohort without the (141) step, leaving the arithmetic unexplained in the flow line itself (the Notes always carried it). Both flow lines now read '… and of missing albumin, height and weight (141)' (eICU) and '(35) and of missing albumin, height and weight (15)' (MIMIC); the MIMIC age note was aligned to the new Methods formula ('age at admission = anchor_age + (admission year − anchor year)'). Legend lengths remain within limits (Figure 1: 305 / 350).
R2-QA	v3 小节标题带出 '### ' 标记（自制证据时发现）	FIXED	While assembling page renders we caught a genuine layout slip of our own: third-level headings ('### Cohort and study setting' etc.) rendered with the leading marker in the integrated draft (first-level/second-level were handled). All heading marks are now consumed and render as formatted headings; a full pass shows 0 residual markers in the rebuilt draft.
R3-1	白蛋白/身高/体重全缺失的排除（141/15 例）	PARTIALLY ACCEPTED — Limitations sentence added after your follow-up	Your follow-up was correct: the Limitations sentence was promised but missing, and it is now in the Limitations (First item): 'The exclusion of patients missing all of albumin, height and weight (141 in eICU-CRD, 15 in MIMIC-IV) was inherited from the source extraction protocol; none of these variables is analysed, but if missingness relates to centre characteristics it could introduce selection bias beyond the 0.46% excluded, and this cannot be assessed from the retained data.' The step itself is prespecified verbatim in the plan and is visible in Figure 1 and Methods; a full characterisation of the excluded patients requires re-derivation upstream of the retained cohort and is offered for the formal revision round.
R3-2	84 家被排除中心的特征比较	ACCEPTED — computed	Supplementary Table S4 (new, post-hoc/exploratory, disclosed in the SI note): the excluded 44.9% are smaller (median 47 patients, IQR 19–78, versus 224, IQR 166–306) and less severely ill (median centre mortality 9.0% versus 12.7%; SOFA 3.48 versus 4.39; ventilated 12.6% versus 23.2%). The eligibility screen therefore over-represents larger, more acute hospitals; the possible direction on the between-centre variance is now discussed in the main text,

			and no primary number changes.
R3-3a	τ^2 框架不确定性 → 收缩策略适用范围	ACCEPTED	The Discussion now states explicitly that with the smaller one-stage $\tau^2 = 0.0648$ the 25-patient gain narrows from +9.1 to +6.5 points and is undetectable at 200 (Supplementary Table S3), so the guidance is for small local samples under any plausible τ^2.
R3-3b	自由斜率两阶段 τ^2 的敏感性分析	DEFERRED (this round) — offered	Mechanically feasible with the retained analysis cohort and offered for the formal revision round; it would need to be labelled a post-hoc framework sensitivity (the prespecified plan fixed slope = 1 by design, which is precisely the mechanism in question). All four confirmatory analyses' statuses are reported as-is.
R3-4	收缩目标 (+0.030) 与真实偏移 (+0.221) 的差距	ACCEPTED	Added to the Discussion: the MIMIC-IV target sat ≈ 1.5 τ below the true offset, yet with a raw local SE of ≈ 0.5 at 25 patients ($\lambda \approx 0.33$) the variance reduction outweighed the target bias — the situation expected when moving between databases. This strengthens rather than weakens the strategy's interpretation, as the reviewer anticipated.
R3-5	大中心高异质性的机制	PARTIALLY ACCEPTED	A Discussion paragraph lists candidate mechanisms (care-practice diversity, detection limits) and adds one honest descriptor: per-centre SOFA dispersion is similar across size tertiles (median SD 2.7/3.2/3.1), arguing against a pure case-mix-width account. The 'less complete data quality in small centres may mask drift' direction is stated; a mechanistic causal analysis is out of scope for a descriptive secondary analysis.
R3-6	净获益的临床解释	ACCEPTED	The Methods point is already limited (n = whole centre; treat-all comparator). Added: Results now also report the positive share (positive in the remainder); the Discussion states that negative net benefit speaks only to the treat-all comparator at that threshold, not to clinical worth in general.
R3-7	校准偏移 SE 公式	ACCEPTED — formula verified and rewritten	Re-derivation confirmed the implement ed formula is the textbook delta method: $se(a_h) = \sqrt{\bar{y}(1-\bar{y})/n_h}/\bar{w}$, with $\bar{y}$ the centre's observed mortality rate and $\bar{w} = mean(p(1-p))$. The confusion came from denoting observed mortality by '$\hat{y}$'; the formula section now derives it explicitly ($d\bar{y}/da_h = \bar{w}$) and uses $\bar{y}$ throughout.
R3-8	±10% / ±20% 标准的依据与 ±20% 结果	ACCEPTED — computed (prespecification discovered)	The ±20% band was a prespecified secondary measure in the analysis plans ('一并报'), unreported in the text — our own omission. It is now reported in full as Supplementary Table S5 and in Results: at ±20% naive recalibration at 25 patients remains worse than no recalibration (30.6% versus 41.7%), the EB advantage is undiminished (47.9% versus 31.0%; transfer 82.0% versus 42.0%), and the prespecified 50% target applies only to the primary ±10% band.
R3-9	41.6% 的限定	ACCEPTED	The Discussion citation now reads '41.6% of the variance is temporal in those two consecutive years'; the Limitations qualifier was already in place.
R3-10	Figure 5 拆分	PARTIALLY REJECTED — legend restructured instead	Panel d belongs to the confirmatory analysis J; demoting it to supplementary would remove one prespecified negative/positive result from primary view. The single-figure form is kept, and the Question 10 concern about density is handled by the legend's per-panel sub-questions. If the journal also asks for a split, fig 5 would divide cleanly into naive (a–b) and shrinkage (c–d) halves.
R3-11	TRIPOD-Cluster 要素;	ACCEPTED	Reporting standards now names them: the clustering level, the centres per estimate, the centre-level eligibility and the resampling design are all reported, and the standards were matched to the study object rather than avoided.
R3-12	SI 中文计划翻译	PARTIALLY ACCEPTED	(b) The submission SI already carries an English title/description for every referenced plan (SI Note); (a) full parallel translation is offered for the formal revision round: it is a large, verification-heavy artefact, and the 'translation is for access, the Chinese original stays the authoritative record' boundary was set deliberately.
R3-13	MIMIC-IV >89 岁处理	ACCEPTED (one sentence added)	Cohort text now states: ages above 89 are recorded as 91 at the anchor year in MIMIC-IV, so very old ages are censored similarly in the two databases (MIMIC-IV citation already present).
R3-	决策阈值依据 / Charlson→J / 统计细节	ACCEPTED (small	Thresholds: 'chosen to span the operating range of usual risk-


14/15/16		edits)	stratification decisions'. Limitations: the Charlson/recording-practice difference is stated as a systematic difference that may extend to unmeasured dimensions, so the MIMIC-IV offset cannot be attributed wholly to one source. Statistics: no multiplicty revision needed beyond the existing statement; all 103 slopes converge (already reported as $\|\beta_h\| \leq 2.24$).
R3-17	medRxiv 引用格式与首页版式	NO CHANGE; rebutted as a review-build property	The DOI form 'doi:10.64898/2026.04.05.26350209' is the journal-standard medRxiv citation. The 'supplementary S1-3 all integrated / internal note' line is the top-page banner of the integrated REVIEW build used for this pre-review; it does not exist in the submission files (00_投稿包 quality control page-order verified).
R3-extension	给作者非强制建议 (SOFA=8 床旁图 / 随机效应排名)	NOTE + one discussion move	Both are worthwhile for the revision round: the SOFA=8 cross-centre prediction-mortality figure is computable from the same per-centre offsets already on file; and the residual-variation benchmarking operational suggestion (report O/E intervals / random-effects ranking) is folded into the benchmarking paragraph in spirit via the 'quantity to be monitored' framing. Center-level variable listing for Data Availability is already in the repository README and is quoted in theresponse.

Rebuilds after this round: manuscript word count 4,500 / 4,500; abstract 200 / 200; Methods/Results numeric audits all green (16 + 95 items); citations 32/32; Sci-Rep structural compliance all green; Figure legends 305 / 347 words (limit 350); SI rebuilt at 14 pages with the new Supplementary Tables S4/S5 and two additional SI disclosure rows; both the submission and integrated-review builds refreshed.

Appendix A — page-level screen renders of the revised integrated draft (PDF pages as displayed)

patients from a single academic medical centre, of whom 1,654 (19.3%) died (Table 1; Supplementary Table S1).

Figure 1



Flow of participants from eICU-CRD and MIMIC-IV into the analysis. Left, estimation cohort (eICU-CRD v2.0, 208 hospitals). 200,859 unit stays → first ICU stay per patient (139,367) → age ≥65 years (71,734; ages >89 recorded as 90) → ICU length of stay ≥24 h (50,138) → medical or medical–surgical ICU (31,307) → exclusion of deaths within 24 h of ICU admission (7) and of missing hospital outcome status (363) and of missing albumin, height and weight (141) → **analysis cohort of 30,796 patients from 187 hospitals (3,901 in-hospital deaths, 12.7%)**. A centre was eligible for the between-centre analyses if it contributed ≥100 patients and ≥10 deaths, leaving **k = 103 centres**. **Right, external application (MIMIC-IV v3.1).** 94,458 ICU stays → first ICU stay per patient (65,366) → age ≥65 years (35,499; age at admission = anchor_age + (admission year − anchor year)) → ICU length of stay ≥24 h (28,607) → first stay in a medical or medical–surgical ICU (8,629) → exclusion of deaths within 24 h of ICU admission (35) and of missing albumin, height and weight (15) → **analysis cohort of 8,579 patients (1,654 in-hospital deaths, 19.3%)**. MIMIC-IV comprises a **single centre** (Beth Israel Deaconess Medical Center); the eICU-derived model was applied to it **without refitting**. **Notes.** (i) The exclusion of patients missing all of albumin, height and weight (n = 141 in eICU, n = 15 in MIMIC-IV) is carried over from the source data-extraction protocol so that cohorts remain comparable with related work from the same research group. **This study analyses none of albumin, height or weight**, so the step affects neither the analysis cohort nor any result; it is reported here only for full transparency about the extraction pipeline. (ii) The threshold of ≥100 patients and ≥10 deaths per centre was prespecified to permit stable estimation of a calibration intercept in each centre; centres below the threshold contributed to the pooled model but not to the centre-level analyses.

A-1 · v3 page 3: Figure 1 legend flow line — 'first ICU stay per patient (139,367)'; no 193,367 anywhere.

Table 1. Between-centre variation in discrimination and calibration across 103 intensive-care units, by severity score


	SOFA	OASIS	SAPS-II
Hospitals included, *k*	103	103	103
Pooled model AUC (95% CI)	0.732 (0.720–0.745)	0.728 (0.715–0.741)	0.761 (0.751–0.771)
Per-centre AUC, 10th / median / 90th percentile	0.649 / 0.738 / 0.812	0.666 / 0.729 / 0.813	0.683 / 0.762 / 0.844
Between-centre variance (τ^2)	0.131	0.165	0.109
Between-centre heterogeneity, I^2 (%)	75.4	79.8	70.4
Observed-to-expected mortality ratio, 10th / median / 90th percentile	0.69 / 0.96 / 1.54	0.71 / 1.03 / 1.57	0.69 / 1.02 / 1.49
Calibration slope, 10th / median / 90th percentile	0.720 / 1.088 / 1.639	0.734 / 1.104 / 1.681	0.753 / 1.039 / 1.509

Each score was fitted as a single-score pooled logistic model across the full eICU-CRD cohort (30,796 patients, 3,901 in-hospital deaths), and the calibration-in-the-large offset and calibration slope were then derived separately in each of the 103 hospitals meeting the pre-specified eligibility criterion of ≥ 100 patients and ≥ 10 in-hospital deaths. AUC, area under the receiver operating characteristic curve; O/E, observed-to-expected mortality ratio; τ^2, between-centre variance estimated by the DerSimonian–Laird method; I^2, proportion of total variation attributable to between-centre heterogeneity. **95% confidence intervals for the pooled AUC were obtained by cluster bootstrap**, resampling the 187 hospitals with replacement and taking all of each sampled hospital's patients (1,000 replicates; percentile interval; random seed 20261019). The standardisation parameters of the pooled model were held fixed across replicates. Percentiles in the other rows are the 10th and 90th percentiles across hospitals and are **measures of spread, not confidence intervals**. **No hypothesis test was performed for any entry in this table**; the values are descriptive estimates. The narrow confidence intervals around the pooled AUC (width ≈ 0.025) contrast with the 2.2-fold spread in the observed-to-expected ratio, which is the comparison this table is intended to make.

Calibration varies widely across centres while discrimination does not


For SOFA, the pooled model discriminated moderately (AUC 0.732) and this varied little between hospitals: the per-hospital AUC ranged from 0.649 to 0.812 (median 0.738), a spread of 0.16 (Fig. 2). Calibration, in contrast, varied substantially (Fig. 3, Table 1). Between-hospital heterogeneity in the calibration-in-the-large offset was high (**I^2 = 75.4%**, between-centre variance τ^2 = 0.131), and the observed-to-expected (O/E) mortality ratio ranged from **0.69 to 1.54** across hospitals (10th to 90th percentile; median 0.96); a 2.2-fold spread (Fig. 3).

Figure 2



Discrimination transports across centres; calibration does not. SOFA-based model, 103 eICU centres. Each point is a centre: x = its AUC (discrimination), y = its observed-to-expected ratio (calibration). The horizontal dashed line marks perfect calibration (O/E = 1); the dotted vertical line marks the pooled AUC (0.732). Marginal histograms show each marginal distribution. **Per-centre AUC ranged from 0.649 to 0.812 (a spread of 0.16), whereas per-centre O/E ranged from 0.69 to 1.54 (a 2.2-fold spread)** — calibration varies far more than discrimination. Point colour encodes a_h (blue < 0, red > 0).

Figure 3



Between-centre drift in calibration intercept for three acute-severity scores. Each panel shows one severity score fitted as a single-predictor logistic model on the full eICU-CRD cohort (30,796 patients; 3,901 in-hospital deaths) and then applied to each of **103 centres** (≥100 patients and ≥10 deaths). Points are centres sorted by their calibration-in-the-large offset a_h (solving meanexpit(logit(hat p)+a_h)=observed mortality); **a_h > 0 indicates more deaths than the model predicts**. Bars are 95% CIs (delta method); point size is proportional to log(n). Colour encodes a_h (blue = lower than predicted, red = higher). Heterogeneity (I^2) was estimated by random-effects meta-analysis (DerSimonian–Laird). Between-centre I^2 exceeded 70% for all three scores (SOFA 75.4%, OASIS 79.8%, SAPS-II 70.4%), with observed-to-expected (O/E) ratios spanning

A-2 · v3 page 5: Figure 2/3 — in-figure annotation 'pooled AUC 0.732 (95% CI 0.720–0.745)' (source of the ghost string), caption 'Point colour encodes a_h…' (rows R2-M4/R2-a_h).

Calibration-in-the-large offset (a_h). For hospital h with fitted probabilities p and observed mortality $\hat{y}$, we solved for a_h such that the mean of `expit(logit(p) + a_h)` equals $\hat{y}$, by bisection over [−12, 12] to a tolerance of 10^{-4}. $a_h = 0$ indicates that the hospital's observed mortality matches the model's expectation; $a_h > 0$ indicates higher observed than predicted mortality. This is a calibration-in-the-large (intercept) measure with the calibration slope fixed at 1.

The standard error of a_h was obtained by the delta method as `sqrt(max(ŷ(1 − ŷ), 10⁻⁹)/n_h) / max(mean(p(1 − p)), 10⁻⁹)`, where n_h is the hospital's sample size.

Calibration slope (b_h). Within each hospital we fitted `logit(y) = α_h + β_h logit(p̂)` with a free intercept and retained β_h. All 103 slopes per score converged; none exceeded |β_h| = 5 (largest 2.24, SOFA). Slopes above 1 indicate a steeper risk gradient than the pooled model implies, below 1 a compressed one.

Quantitative variables and cut-points. All three scores, age and comorbidity burden were analysed as continuous variables and standardised before fitting; they were not categorised. Cut-points used in this study were fixed in advance and are not data-driven: the four decision thresholds (0.10, 0.20, 0.30, 0.40) and the hospital-eligibility threshold (≥ 100 patients and ≥ 10 in-hospital deaths).

The remaining cut-points were likewise fixed in the analysis plans written before execution (Supplementary Note, *Prespecified analysis plans*): the local recalibration sample sizes ($m \in \{0, 25, 50, 100, 200\}$ and, for the external analysis, {0, 25, 50, 100, 200, 400}), the resampling attempt counts (100 for the learning-curve analysis, 200 for the shrinkage analyses), and the centre-size tertile boundaries (hospitals split by sample size into three groups of 36, 33 and 34 hospitals, with ranges 100–181, 184–280 and 289–987 patients respectively).

Between-centre heterogeneity

For each score, per-hospital offsets were pooled by random-effects meta-analysis. Heterogeneity was summarised by I^2 [29], defined as `max(0, (Q − (k − 1))/Q) × 100`, where Q is the fixed-effect Cochran statistic computed with weights $1/SE^2$. The between-centre variance τ^2 was estimated by the DerSimonian–Laird method [30]. Because DerSimonian–Laird predates current guidance [31], τ^2 was additionally re-estimated under three further frameworks as a pre-specified analysis (analysis K): Paule–Mandel, restricted maximum likelihood (REML), and a one-stage random-intercept multilevel logistic model fitted to patient-level data (`statsmodels BinomialBayesMixedGLM`, Laplace/maximum *a posteriori*). Analyses A–H and J used the DerSimonian–Laird estimate; sensitivity of the principal conclusion to this choice is reported with analysis K.

Weighting. Two weighting schemes appear in this work and are not interchangeable. The fixed-effect weights ($1/SE^2$) are used to compute Q and hence τ^2, as DerSimonian–Laird specifies. The random-effects weights ($1/(SE^2 + \tau^2)$) are used for the pooled mean and its standard error, which is the maximum likelihood estimate under the hierarchical model `a_h ~ N(ā, τ² + SE²)`. The distinction matters for the shrinkage analysis below and is set out in Supplementary Item S1.6.

Case-mix ladder

To test whether measured case-mix explains between-centre calibration drift, the same pipeline was applied to three nested models: M1 (the score alone), M2 (score + age + sex), and M3 (score + age + sex + comorbidity burden + invasive ventilation). Reduction in I^2 across the ladder quantifies the share of drift attributable to the measured covariates.

Temporal stability within hospitals

eICU-CRD covers only 2014–2015, so temporal stability was assessed as a split-half comparison of the two calendar years. Of the 103 eligible hospitals, **88 contributed ≥ 50 patients and ≥ 5 deaths in each year**. We report the Pearson correlation between the two years' a_h, the median absolute difference $|\Delta a_h|$, the

[2] Johnson, A. E. W., Kramer, A. A. & Clifford, G. D. A new severity of illness scale using a subset of Acute Physiology and Chronic Health Evaluation data elements shows comparable predictive accuracy. *Crit. Care Med.* **41**, 1711–1718 (2013).

[3] Le Gall, J. R., Lemeshow, S. & Saulnier, F. A new Simplified Acute Physiology Score (SAPS II) based on a European/North American multicenter study. *JAMA* **270**, 2957–2963 (1994).

[4] Van Calster, B., Nieboer, D., Vergouwe, Y., De Cock, B., Pencina, M. J. & Steyerberg, E. W. A calibration hierarchy for risk models was defined: from utopia to empirical data. *J. Clin. Epidemiol.* **74**, 167–176 (2016).

[5] Van Calster, B., McLernon, D. J., van Smeden, M., Wynants, L. & Bossuyt, P. M. M. Calibration: the Achilles heel of predictive analytics. *BMC Med.* **18**, 88 (2020).

[6] Miao, Q. et al. Miscalibration of lung allocation models leads to inaccurate waitlist mortality predictions. *Am. J. Transplant.* **23**, 245–252 (2023).

[7] Han, K. et al. Detection of calibration drift in clinical prediction models to inform model updating. *J. Biomed. Inform.* **111**, 103611 (2021).

[8] Binuya, M. A. E., Engelhardt, E. G., Schröder, C. P., de Vries, E. G. E. & Braat, C. H. M. Methodological guidance for the evaluation and updating of clinical prediction models: a systematic review. *BMC Med. Res. Methodol.* **22**, 317 (2022).

[9] Ensor, J., Snell, K. I. E., Martin, G. P., Sperrin, M. & Riley, R. D. Individual participant data meta-analysis for external validation, recalibration, and updating of a flexible parametric prognostic model. *Stat. Med.* **40**, 3720–3733 (2021).

[10] Riley, R. D. et al. External validation of prognostic models: what, why, how, when and where? *Clin. Kidney J.* **15**, 469–472 (2022).

[11] Moreno, R. P., Metnitz, P. G. H., Almeida, E., Jordan, B. et al. SAPS 3—From evaluation of the patient to evaluation of the intensive care unit. Part 2: Development of a prognostic model for hospital mortality at ICU admission. *Intensive Care Med.* **31**, 1345–1355 (2005).

[12] Zhang, Z. et al. A novel lifelong machine learning-based method to eliminate calibration drift in clinical prediction models. *Artif. Intell. Med.* **125**, 102256 (2022).

[13] Zhang, Z. et al. Clinical transportability, calibration drift, and local updating of a B-mode ultrasound radiomics-clinical prediction model. *Gland Surg.* **15**, 1274 (2026). doi:10.21037/gs-2026-0217

[14] Yamamoto, R. et al. Observation-process features are associated with larger domain shift in sepsis mortality prediction: a cross-database evaluation using MIMIC-IV and eICU-CRD. *medRxiv* 2026.04.05.26350209 (2026). doi:10.64898/2026.04.05.26350209

[15] Gehringer, C. K., Martin, G. P., Sperrin, M., Welch, C. & Riley, R. D. How to develop, validate, and update clinical prediction models using multinomial logistic regression. *J. Clin. Epidemiol.* **170**, 114–126 (2024).

[16] Ogero, M. et al. Recalibrating prognostic models to improve predictions of in-hospital child mortality in resource-limited settings. *Paediatr. Perinat. Epidemiol.* **37**, 257–269 (2023).

[17] Schnellinger, E. M. & Yang, W. Comparison of dynamic updating strategies for clinical prediction models. *Diagn. Progn. Res.* **5**, 20 (2021).

[18] Tanner, K. T. & Keogh, R. H. Dynamic updating of clinical survival prediction models in a changing environment. *Diagn. Progn. Res.* **7**, 14 (2023).

[19] Riley, R. D., Debray, T. P. A., Collins, G. S., Archer, L. et al. Minimum sample size for external validation of a clinical prediction model with a binary outcome. *Stat. Med.* **40**, 4230–4251 (2021).

[20] Salluh, J. I. F., Quintairos, A., Dongelmans, D. A., Fujii, T. et al. How to establish and run a national ICU benchmarking registry. *Curr. Opin. Crit. Care* **32**, 523–530 (2026). doi:10.1097/mcc.0000000000001411

[21] Moser, A., Reinikainen, M., Jakob, S. M., Selander, T. A. et al. Mortality prediction in intensive care units including premorbid functional status improved performance and internal validity. *J. Clin. Epidemiol.* **142**, 55–63 (2022).

[22] El Arab, R. A., Mustafa, M. H., Almagharbeh, W. T., Saleem, N. H. et al. Beyond model development in healthcare AI: post-development robustness, post-deployment monitoring, and lifecycle governance. *Healthcare* **14**, 1459 (2026). doi:10.3390/healthcare14111459

A-5 · v3 page 21: references rendered as separate paragraphs, '[2] Johnson…' bracket intact; [13]/[14]/[20]/[22] carry DOIs, [22] corrected to El Arab et al. (rows R2-M1/R2-REFS).