Search Search Results

Searches and alerts

Search History

Marked List



Find PDF

Full Text Options



Add to Marked List

Tools

11 of 65

Optimally estimating the sample mean from the sample size, median, midrange, and/or mid-quartile range

 $\textbf{By:} \; \mathsf{Luo}, \mathsf{DH} \; (\mathsf{Luo}, \mathsf{Dehui})^{\texttt{[1]}}; \; \mathsf{Wan}, \; \mathsf{X} \; (\mathsf{Wan}, \mathsf{Xiang})^{\texttt{[2]}}; \; \mathsf{Liu}, \; \mathsf{JM} \; (\mathsf{Liu}, \mathsf{Jiming})^{\texttt{[2]}}; \; \mathsf{Tong}, \; \mathsf{TJ} \; (\mathsf{Tong}, \; \mathsf{Tiejun})^{\texttt{[1]}}; \; \mathsf{Tong}, \; \mathsf{TIejun})^{\texttt{[1]}}; \; \mathsf{Tong}, \; \mathsf{TIejun}, \; \mathsf{TIejun},$ View Web of Science ResearcherID and ORCID

STATISTICAL METHODS IN MEDICAL RESEARCH

Volume: 27 Issue: 6 Pages: 1785-1805 DOI: 10.1177/0962280216669183

Published: JUN 2018 Document Type: Article **View Journal Impact**

Abstract

The era of big data is coming, and evidence-based medicine is attracting increasing attention to improve decision making in medical practice via integrating evidence from well designed and conducted clinical research. Meta-analysis is a statistical technique widely used in evidence-based medicine for analytically combining the findings from independent clinical trials to provide an overall estimation of a treatment effectiveness. The sample mean and standard deviation are two commonly used statistics in meta-analysis but some trials use the median, the minimum and maximum values, or sometimes the first and third quartiles to report the results. Thus, to pool results in a consistent format, researchers need to transform those information back to the sample mean and standard deviation. In this article, we investigate the optimal estimation of the sample mean for meta-analysis from both theoretical and empirical perspectives. A major drawback in the literature is that the sample size, needless to say its importance, is either ignored or used in a stepwise but somewhat arbitrary manner, e.g. the famous method proposed by Hozo etal. We solve this issue by incorporating the sample size in a smoothly changing weight in the estimators to reach the optimal estimation. Our proposed estimators not only improve the existing ones significantly but also share the same virtue of the simplicity. The real data application indicates that our proposed estimators are capable to serve as rules of thumb and will be widely applied in evidence-based medicine.

Kevwords

Author Keywords: Median; meta-analysis; mid-range; mid-quartile range; optimal weight; sample mean; sample size KeyWords Plus: VITAMIN-D DEFICIENCY; PULMONARY TUBERCULOSIS; METAANALYSIS; CALCIUM

Author Information

Reprint Address: Tong, TJ (reprint author)

Hong Kong Baptist Univ, Dept Math, Hong Kong, Hong Kong, Peoples R China.

Reprint Address: Wan, X (reprint author)

Hong Kong Baptist Univ, Dept Comp Sci, Hong Kong, Hong Kong, Peoples R China.

🛨 [1] Hong Kong Baptist Univ, Dept Math, Hong Kong, Hong Kong, Peoples R China

🛨 [2] Hong Kong Baptist Univ, Dept Comp Sci, Hong Kong, Hong Kong, Peoples R China

E-mail Addresses: xwan@comp.hkbu.edu.hk; tongt@hkbu.edu.hk

Funding

Funding Agency	Show details	Grant Number
нкви		FRG2/14-15/077 FRG1/14-15/084 FRG2/15-16/019
Hong Kong Research Grants Council		HKBU12202114
HKBU Century Club Spons	orship Scheme	

View funding text

Publisher

SAGE PUBLICATIONS LTD, 1 OLIVERS YARD, 55 CITY ROAD, LONDON EC1Y 1SP, ENGLAND

Journal Information

Performance Trends: Essential Science Indicators Impact Factor: Journal Citation Reports

Citation Network

In Web of Science Core Collection

111

Highly Cited Hot Paper

Times Cited

Create Citation Alert

All Times Cited Counts

111 in All Databases

See more counts

21

Cited References

View Related Records

Most recently cited by:

Suarez, Sebastian; Conde-Agudelo, Agustin; Borovac-Pinheiro, Anderson; et al. Uterine balloon tamponade for the treatment of postpartum hemorrhage: a systematic review and meta-analysis AMERICAN JOURNAL OF OBSTETRICS AND GYNECOLOGY (2020)

Yan, Sen-bo; Wang, Xiao-yan; Shang, Guokai; et al.

Impact of Perioperative Levosimendan Administration on Risk of Bleeding After Cardiac Surgery: A Meta-analysis of Randomized Controlled Trials AMERICAN JOURNAL OF CARDIOVASCULAR

View All

Use in Web of Science

Web of Science Usage Count

21

Last 180 Days Since 2013

Learn more

This record is from: Web of Science Core Collection

View Record in Other Databases: View medical data (in MEDLINE®)

Suggest a correction

If you would like to improve the auality of the data in this record, please suggest a correction.