Web of Science



登入 ▼ 説明 ▼ 繁體中文 ▼

檢索 回到檢索結果











4 2 of 3 ▶

勾選的清單

Optimally estimating the sample mean from the sample size, median, mid-range, and/or mid-quartile range

作者: Luo, DH (Luo, Dehui)[1]; Wan, X (Wan, Xiang)[2]; Liu, JM (Liu, Jiming)[2]; Tong, TJ (Tong, Tiejun)[1] 檢視 Web of Science ResearcherID 和 ORCID

STATISTICAL METHODS IN MEDICAL RESEARCH

卷冊: 27 **期:** 6 **頁數:** 1785-1805 **DOI:** 10.1177/0962280216669183

出版: JUN 2018 文件類型: Article 檢視期刊影響力

摘要

The era of big data is coming, and evidence-based medicine is attracting increasing attention to improve decision making in medical practice via integrating evidence from well designed and conducted clinical research. Meta-analysis is a statistical technique widely used in evidence-based medicine for analytically combining the findings from independent clinical trials to provide an overall estimation of a treatment effectiveness. The sample mean and standard deviation are two commonly used statistics in meta-analysis but some trials use the median, the minimum and maximum values, or sometimes the first and third quartiles to report the results. Thus, to pool results in a consistent format, researchers need to transform those information back to the sample mean and standard deviation. In this article, we investigate the optimal estimation of the sample mean for meta-analysis from both theoretical and empirical perspectives. A major drawback in the literature is that the sample size, needless to say its importance, is either ignored or used in a stepwise but somewhat arbitrary manner, e.g. the famous method proposed by Hozo etal. We solve this issue by incorporating the sample size in a smoothly changing weight in the estimators to reach the optimal estimation. Our proposed estimators not only improve the existing ones significantly but also share the same virtue of the simplicity. The real data application indicates that our proposed estimators are capable to serve as rules of thumb and will be widely applied in evidence-based medicine.

關鍵字

作者關鍵字: Median; meta-analysis; mid-range; mid-quartile range; optimal weight; sample mean; sample size **KeyWords Plus**: VITAMIN-D DEFICIENCY; PULMONARY TUBERCULOSIS; METAANALYSIS; CALCIUM

作者資訊

涌訊作者地址:

Hong Kong Baptist University Hong Kong Baptist Univ, Dept Math, Hong Kong, Hong Kong, Peoples R China. Hong Kong Baptist University Hong Kong Baptist Univ, Dept Comp Sci, Hong Kong, Hong Kong, Peoples R China.

通訊地址: Tong, TJ (通訊作者)

Hong Kong Baptist Univ, Dept Math, Hong Kong, Hong Kong, Peoples R China.

通訊地址: Wan, X (通訊作者)

Hong Kong Baptist Univ, Dept Comp Sci, Hong Kong, Hong Kong, Peoples R China.

地址:

- 🛨 [1] Hong Kong Baptist Univ, Dept Math, Hong Kong, Hong Kong, Peoples R China
- + [2] Hong Kong Baptist Univ, Dept Comp Sci, Hong Kong, Hong Kong, Peoples R China

電子郵件地址: xwan@comp.hkbu.edu.hk; tongt@hkbu.edu.hk

贊助

贊助機構	顯示詳細資料	補助編號
НКВИ		FRG2/14-15/077 FRG1/14-15/084 FRG2/15-16/019
Hong Kong Research Grants Council		HKBU12202114
HKBU Century Club Sponso	orship Scheme	

檢視贊助文字

出版者

SAGE PUBLICATIONS LTD, 1 OLIVERS YARD, 55 CITY ROAD, LONDON EC1Y 1SP, ENGLAND

引用文獻網路

於 Web of Science 核心合輯

259

工具 ▼ 檢索與追蹤 ▼ 檢索歷史



被引用次數



所有被引用次數計數

260 於 所有資料庫

查看較多計數

21

參考文獻

檢視 Related Records

① 全新推出! 您可能也會喜歡... BETA

Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range.

BMC MEDICAL RESEARCH METHODOLOGY (2014)

Quantifying publication bias in metaanalysis.

BIOMETRICS (2018)

How Do Small and Medium-Sized Game Companies Use Analytics? An Attention-Based View of Game Analytics. INFORMATION SYSTEMS FRONTIERS (2020)

SIMEX estimation for single-index model with covariate measurement error.
ASTA-ADVANCES IN STATISTICAL ANALYSIS (2019)

Prevalence and impact of cardiovascular metabolic diseases on COVID-19 in China. CLINICAL RESEARCH IN CARDIOLOGY (2020

檢視所有建議

最近被以下文獻引用:

Wang, Min; Liu, Tianyu; Niu, Zheng; 等. Utility of venous-to-arterial carbon dioxide changes to arteriovenous oxygen content ratios in the prognosis of severe sepsis and septic shock: A systematic review and meta-analysis.

HONG KONG JOURNAL OF EMERGENCY MEDICINE (2021)

Lex, Johnathan R.; Edwards, Thomas C.; Packer, Timothy W.; 等.

Perioperative Systemic Dexamethasone Reduces Length of Stay in Total Joint Arthroplasty: A Systematic Review and Meta-Analysis of Randomized Controlled Trials.

JOURNAL OF ARTHROPLASTY (2021)