

Homework 3: Astr 5140 (Fall 2021)

Tien Vo

September 19, 2021

Problem 1 (Generalized Ohm's Law): The MHD force equation is derived by a linear combination of the electron and ion fluid equations. Generalized Ohm's law is, simply stated, a different linear combination.

We begin by multiplying the ion force equation by m_e and the electron force equation by m_i , then subtract

$$m_e m_i n \frac{D\mathbf{u}_i}{Dt} = -m_e \nabla P_i + n m_e e (\mathbf{E} + \mathbf{u}_i \times \mathbf{B}) - m_e m_i n \nu_{ie} (\mathbf{u}_i - \mathbf{u}_e) \quad (1.1a)$$

$$m_i m_e n \frac{D\mathbf{u}_e}{Dt} = -m_i \nabla P_e - n m_i e (\mathbf{E} + \mathbf{u}_e \times \mathbf{B}) - m_e m_i n \nu_{ei} (\mathbf{u}_e - \mathbf{u}_i) \quad (1.1b)$$

We use quasi-neutrality and bundle the convective derivative into the symbol D .

$$\begin{aligned} m_e m_i n \left(\frac{D\mathbf{u}_i}{Dt} - \frac{D\mathbf{u}_e}{Dt} \right) &= (-m_e \nabla P_i + m_i \nabla P_e) + en \mathbf{E} (m_e + m_i) \\ &\quad + en (m_e \mathbf{u}_i + m_i \mathbf{u}_e) \times \mathbf{B} - m_e m_i n (\nu_{ei} + \nu_{ie}) (\mathbf{u}_i - \mathbf{u}_e) \end{aligned} \quad (1.2)$$

(a) Divide each term by $en(m_i + m_e)$.

(b) Show that Term 1 can be re-written in the limit of $m_i \gg m_e$ as

$$\frac{m_e}{e^2} \frac{D(\mathbf{J}/n)}{Dt} \quad (1.3)$$

(c) Argue that since $m_i \gg m_e$ that, unless $P_i \gg P_e$ (very rare), Term 2 becomes

$$\frac{\nabla P_e}{en} \quad (1.4)$$

(d) Term 3 is trivial. Term 4 is tricky as it must be broken into two parts. Add $\mathbf{u} = (m_i \mathbf{u}_i + m_e \mathbf{u}_e)/(m_i + m_e)$, separate $\mathbf{u} \times \mathbf{B}$, then subtract $(m_i \mathbf{u}_i + m_e \mathbf{u}_e)/(m_i + m_e)$. Show that the remaining four terms ($m_i \gg m_e$) can be approximated as

$$\frac{-\mathbf{J} \times \mathbf{B}}{en} \quad (1.5)$$

(e) Show that Term 5 can be written as \mathbf{J}/σ . Define σ .

In the end, you should arrive at

$$\mathbf{E} + \mathbf{u} \times \mathbf{B} = \frac{\mathbf{J}}{\sigma} + \frac{\mathbf{J} \times \mathbf{B}}{en} - \frac{\nabla P_e}{en} + \frac{m_e}{ne^2} \frac{D\mathbf{J}}{Dt} + \text{small terms} \quad (1.6)$$

Note: If done exactly (full convective derivative and keeping small terms), n is not inside the derivative and furthermore, some of the "leftovers" from Term 1 cancel "leftovers" in Term 4.

Solution.

□

Problem 2 (Scale height in the solar corona): The surface gravity of the Sun is 274 m/s^2 . Assume that the corona is entirely protons with $T_{\text{corona}} = 10^6 \text{ }^\circ\text{K}$. Derive the isothermal scale height of the solar corona from the MHD equations assuming \mathbf{g} is constant (use 1D). How does this value compare with the radius of the Sun (what % of R_{sun} is H_0)? Does your answer agree with what is seen in UV or X-ray images?

Solution.

□

Problem 4 (Magnetic tension): This problem is to develop an intuition for magnetic tension. Examine the diagram, which shows a field line if $\mathbf{B} = B_0[(z/z_0)\hat{\mathbf{x}} + \hat{\mathbf{z}}]$. Examine the field line and convince yourself that the equation represents a curved magnetic field line. Derive the tension force at $z = 0$. What is the direction of the tension force? Argue that z_0 is the local radius of curvature.

Solution.

□