

Homework 2: Astr 5140 (Fall 2021)

Tien Vo

September 4, 2021

Problem 1 ($\mathbf{J} \times \mathbf{B}$ force): Using the two-fluid equations, calculate the Lorentz force per unit volume on a quasi-neutral (MHD) plasma using the definition of \mathbf{J} . Separate the electric force (\mathbf{F}_E) from the magnetic force (\mathbf{F}_B). Show that if the plasma is quasi-neutral, then \mathbf{F} reduces to the standard MHD result.

Problem 2 (EM review: Waves in a plasma): Using Maxwell's equations and setting the current to be the electron motion only

$$\frac{\partial \mathbf{J}}{\partial t} = \frac{ne^2 E}{m_e} \quad (2.1)$$

Show that the solution of a transverse light wave becomes:

$$\omega^2 = \omega_{pe}^2 + k^2 c^2 \quad (2.2)$$

where $\omega_{pe}^2 = ne^2/\epsilon_0 m_e$.

Problem 3 (Pick-up Ion at Mars): Suppose an O atom that escaped from Mars is at rest (in our frame) at $x = 0, y = 0$. It is photo-ionized (charge e) at $t = 0$ in the solar wind ($v_{sw} = 350$ km/s in the x direction) with a magnetic field $\mathbf{B} = 10$ nT in the z direction (see diagram). Assume that the photo-ionization does not move the O^+ ion.

(1) Describe the subsequent motion of the O^+ ion, $x(t), y(t), v_x(t), v_y(t)$, in our rest frame (not the plasma frame). *Hint*: What is the solar wind electric field? Calculate $v_x(t)$ and $v_y(t)$ then integrate. Apply boundary conditions, $x(t = 0) = 0, y(t = 0) = 0, v_x(t = 0) = 0, v_y(t = 0) = 0$ to get an exact solution.

(2) Sketch the O^+ path.

(3) The drift and gyration cause the O^+ ion to slow down and speed up in our rest frame. What is the drift speed? What is the gyration speed? What is the maximum velocity (km/s) and energy (keV) that the O^+ ion reaches?

(4) Using the gyration speed only, what is the perpendicular (to \mathbf{B}) temperature of that ion in $^\circ\text{K}$? (In 2D, temperature and energy are equal.)

Problem 4 (Current sheet): A current sheet is such that \mathbf{J} is in the y direction and \mathbf{B} is in the z direction. \mathbf{B}, \mathbf{J} , and P vary only with x (see diagram). Derive a solution for \mathbf{B} , and \mathbf{J} under the condition $\mathbf{J} \sim P$ that is valid for $-L < x < L$. Make sure that your

solution satisfies the boundary conditions of $\mathbf{B}(x = L) = -B_0\hat{\mathbf{z}}$, and $\mathbf{B}(x = 0) = \mathbf{0}$. L is a characteristic length. Sketch your results.

Hint: There is more than one possible solution – just give any valid solution. Be careful, the condition $\mathbf{J}^2 \sim P$ is NOT the same as in the Harris solution.

Problem 5 (Magnetic diffusion): Consider a magnetic field $\mathbf{B} = B_z(x, t)\hat{\mathbf{z}}$ where $B_z(x, t = 0) = B_0 \cos(k_1 x) + B_0 \cos(k_2 x)$ in a resistive plasma with $k_2 \gg k_1$.

(a) Find the solution for $B_z(x, t)$.

(b) Sketch (accurate plot not needed) the solution for $t = 0$ and $t > 0$. What happens to the high- k wave?