

The representation of peripheral neural activity in the middle-latency evoked field of primary auditory cortex in humans



Till Habersetzer, Stefan Uppenkamp, Andreas Spiegler

The representation of peripheral neural activity in the middle-latency evoked field of primary auditory cortex in humans(1)



André Rupp¹, Stefan Uppenkamp, Alexander Gutschalk, Roland Beucker, Roy D Patterson, Torsten Dau, Michael Scherg

Affiliations + expand



PMID: 12433393 DOI: [10.1016/s0378-5955\(02\)00614-7](https://doi.org/10.1016/s0378-5955(02)00614-7)



The representation of peripheral neural activity in the middle-latency evoked field of primary auditory cortex in humans



sub-01: feste Dipolorientierung nach SVD



The representation of peripheral neural activity in the middle-latency evoked field of primary auditory cortex in humans

Contraindications - Who can not participate?

To avoid magnetic interactions – especially with metals – because they reduce the measurement accuracy, **no metals** should be worn on your body during the measurements.

- No pacemakers, neuro-stimulators, insulin pumps or hearing devices.
- No metallic implants, they also have the possibility to be ferro magnetic (**Retainer!!**)
- Tattoos (also permanent make-up) can contain color particles with metallic parts in it.
- Also cosmetics such as mascaras, rouge or even styling gel can contain metal. Please avoid using those on the day of measurement.
- ...

We are planning to include an anatomical MRI of your brain in this project