

DARBOUX, MOSER AND WEINSTEIN THEOREMS FOR PREQUANTUM SYSTEMS

EVA MIRANDA AND JONATHAN WEITSMAN

4 **ABSTRACT.** We establish
analogues of the Darboux, Moser and Weinstein
theorems for pre- ∞
quantum systems. We show that two prequantum
systems on a manifold with vanishing H^1
first cohomology, with symplectic forms
defining the same cohomology class and homo- ∞
topic to each other within that class, differ only by
a symplectomorphism and a gauge

r transformation. As
an application, we
show that the Bohr-Sommerfeld quantization of pre-
quantum system on a
the connection.

Ap
manifold with trivial
first cohomology is independent of the choice of ∇

]

G

1. INTRODUCTION

Math.S

The Darboux theorem establishes that there are no local invariants in symplectic geometry. Namely, let M be a smooth manifold and let ω, ω' be two symplectic forms

geom-

on M ,

[m:en

then:

Theorem 1 (Darboux [1, 2]).

For every point $p \in$

M there exists a neighbourhood U of p and an embedding $\Phi : U \rightarrow$

M isotopic

to the inclusion and fixing p such that \triangleright

88

$\Phi^* \omega' = \omega|_U.$

9

The global invariants of a symplectic manifold

include the cohomology class defined by its symplectic form. This gives \mathbb{N}

a complete classification of symplectic
2-manifolds. In $\mathbb{C}P^2$ general,

there is
the following theorem due to Moser.

Theorem 2 (Moser [6]). *Let M be a compact manifold endowed with two symplectic forms ω and ω' . Assume that $[\omega] = [\omega']$, and that there exists a path ω_t of symplectic forms such that $[\omega_t] = [\omega_0]$ for all t , and with $\omega_0 = \omega$ and $\omega_1 = \omega'$. Then, there exists a diffeomorphism isotopic to the identity $\Phi : M \rightarrow M$ such that $\Phi^*\omega' = \omega$.*

iv:240
rX

This proof based
on Moser's method can be
adapted when symmetries are present. Un-
der the conditions above
when a
compact group acts on M preserving ω
and ω' , and where

the path ω_t also consists of
invariant forms, we have the following
result.

Theorem 3 (Weinstein [7]). *There exists an equivariant diffeomorphism $\Phi : M \rightarrow M$ such that $\Phi^*\omega' = \omega$.*

Recall that if (M, ω) is a symplectic
manifold, $\pi : L \rightarrow M$ is a (complex) line

bundle, ∇ is a connection on L , and $\text{curv}(\nabla) = \omega$, the quadruple (M, ω, L, ∇) is a *prequantum system*.

Both authors are partially supported by the Spanish State Research Agency grant

PID2019-103849GB-I00 of MICIU/AEI / 10.13039/501100011033. E. Miranda is supported by the Catalan Institution for Research and Advanced Studies via an ICREA Academia Prize 2021 and by the Alexander Von Humboldt foundation via a Friedrich Wilhelm Bessel Research Award. E. Miranda is also supported by the Spanish State Research Agency, through the Severo Ochoa and María de Maeztu Program for Centers and Units of Excellence in R&D (project CEX2020-001084-M) and partially supported by and by the AGAUR project 2021 SGR 00603.

J. Weitsman was supported in part by a Simons Collaboration Grant # 579801.

April 9, 2024.