

North West London Integrated Formulary

The Integrated Formulary (IF) is a list of medicines for adults and is for use in North West London.

The IF is intended to cover:

- prescriptions written in primary care;
- prescriptions or recommendations by hospital doctors in respect of outpatients or patients leaving hospital after an admission.

The IF should be taken fully into account by prescribers when exercising their clinical judgement. However, it does not override the individual responsibility of prescribers to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or guardian or carer and informed by the summary of product characteristics of any medicines they are considering.

Medicines recommended by NICE technology appraisal guidance (TAG) are regarded as included in the IF from the date the TAG is published, unless:

- the medicine will only be prescribed for hospital inpatients or
- it is deemed to be a medicine that should only be prescribed by a hospital specialist (in which case it will be added to the NWL 'red list' of medicines that a hospital doctor should not ask a GP to prescribe).

People currently receiving a medicine that is not included in the IF should have the option to continue until they and their clinicians consider it appropriate to stop.

Specialist initiation

A number of medicines in the IF are marked 'specialist initiation'.

These words should be interpreted as an attempt to reduce the risk of harm and taken into account by prescribers when exercising their clinical judgement.

Appropriate action in response to a 'specialist initiation' annotation may lie anywhere in a spectrum between:

1. The medicine should be prescribed by a relevant specialist for at least x months with monitoring of effectiveness and toxicity, until treatment is stable at which point the patient's GP can be asked whether s/he will take over prescribing, and
2. It is reasonable for a GP to write the first prescription after a telephone conversation with a relevant specialist clinician.

Different GPs might reasonably take different decisions regarding whether or when to prescribe, depending on their knowledge and experience.

Legal responsibility for prescribing lies with the doctor who signs the prescription.

The 'specialist initiation' annotations say nothing about who is or is not a relevant specialist in respect of a medicine. It will be reasonable for some of these medicines to be initiated by prescribers from more than one speciality.

Version: 64

Last updated: 03/09/2026

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
1. Gastro-intestinal system	1.1 Dyspepsia and gastro-oesophageal reflux disease	1.1.1 Antacids and simeticone	Co-magaldrox (prescribe Mucogel)	co-magaldrox 195/220 suspension	Unspecified	All prescribers	Low sodium. Mucogel costs less than Maalox.
1. Gastro-intestinal system	1.1 Dyspepsia and gastro-oesophageal reflux disease	1.1.2 Compound alginates	Peptac ®	suspension (peppermint or aniseed)	Unspecified	All prescribers	Contains alginate. Contains 6.2mmol sodium in 10ml.
1. Gastro-intestinal system	1.1 Dyspepsia and gastro-oesophageal reflux disease	1.1.2 Compound alginates	Gaviscon Infant®	oral powder;	Unspecified	All prescribers	Contains alginate and 0.92mmol sodium /dose.
1. Gastro-intestinal system	1.2 Antispasmodics and other drugs altering gastric motility	1.2	Mebeverine hydrochloride	tablets 135 mg	Unspecified	All prescribers	
1. Gastro-intestinal system	1.2 Antispasmodics and other drugs altering gastric motility	1.2	Peppermint oil (prescribe Mintec)	e/c capsules 0.2ml	Unspecified	All prescribers	Mintec capsules cost less than MR Colpermin capsules
1. Gastro-intestinal system	1.2 Antispasmodics and other drugs altering gastric motility	1.2	Hyoscine butylbromide	tablets 10mg; injection 20mg in 1ml	Unspecified	All prescribers	
1. Gastro-intestinal system	1.3 Anti-secretory drugs and mucosal protectants	1.3.3 Chelates and complexes	Pylera ®	capsules 140mg bismuth subcitrate potassium , 125mg metronidazole , 125mg tetracycline	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
1. Gastro-intestinal system	1.3 Anti-secretory drugs and mucosal protectants	1.3.3 Chelates and complexes	Sucralfate	oral suspension sugar free 1g/5ml	Unspecified	Specialist INITIATION	Only when PPI is not appropriate / sufficient.
1. Gastro-intestinal system	1.3 Anti-secretory drugs and mucosal protectants	1.3.5 Proton pump inhibitors	Esomeprazole	tablets 20mg, 40mg	Unspecified	Specialist INITIATION	
1. Gastro-intestinal system	1.3 Anti-secretory drugs and mucosal protectants	1.3.5 Proton pump inhibitors	Omeprazole	capsules 10mg, 20mg; dispersible tablets, film-coated, 10mg, 20mg	First line	All prescribers	Capsules cost much less than dispersible tablets
1. Gastro-intestinal system	1.3 Anti-secretory drugs and mucosal protectants	1.3.5 Proton pump inhibitors	Pantoprazole	tablets 20mg, 40mg	Unspecified	All prescribers	
1. Gastro-intestinal system	1.3 Anti-secretory drugs and mucosal protectants	1.3.5 Proton pump inhibitors	Lansoprazole	orodispersible tablet, 15mg, 30mg (Fastab ®); capsules 15mg, 30mg	First line	All prescribers	Capsules cost much less than orodispersible tablets
1. Gastro-intestinal system	1.3 Anti-secretory drugs and mucosal protectants	Test for Helicobacter pylori	diabact UBT		Unspecified	All prescribers	Arrangements for <i>H. pylori</i> testing differ in different parts of NWL
1. Gastro-intestinal system	1.4 Acute diarrhoea	1.4.2 Antimotility drugs	Loperamide	capsules 2mg	First line	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
1. Gastro-intestinal system	1.5 Chronic bowel disorders	1.5.1 Aminosalicylates	Mesalazine	e/c tablets 400mg (Octasa [®] - see comments), 800mg (Octasa [®]); MR tablets 500mg (Pentasa [®]); MR tablets 1.2g (Mezavant XL [®] - see comments) gastro-resistant prolonged release granules 500mg, 1g (Salofalk [®]); retention enema 1g in 100ml; enema 2g in 59ml; foam enema 1g/application; suppositories 500mg, 1g	First line	Specialist INITIATION	There is no evidence to show that any one oral preparation of mesalazine is more effective than another; however, the delivery characteristics of oral mesalazine preparations may vary. Newly initiated prescriptions should be for Octasa MR[®] tablets rather than Asacol MR[®] tablets. It is reasonable to switch patients from Asacol MR[®] tablets to Octasa MR[®] tablets (subject to the exception below). Patients should be advised to report any change of symptoms. If a patient's disease has been unstable on a maximum tolerated or licensed dose of Octasa MR[®] and they have been switched to the same dose of Asacol MR[®] which has resulted in worthwhile improvement (which would not be expected based on the pharmaceutical similarity of the products) they should not be switched back to Octasa MR[®]. Mezavant XL[®] should only be considered for select patients experiencing difficulty with a high pill burden (i.e. >2.4g/daily).
1. Gastro-intestinal system	1.5 Chronic bowel disorders	1.5.1 Aminosalicylates	Sulfasalazine	tablets 500mg; e/c tablets 500mg; suspension 250mg in 5ml; suppositories 500mg	Unspecified	Specialist INITIATION	To be used in line with a shared care protocol
1. Gastro-intestinal system	1.5 Chronic bowel disorders	1.5.1 Aminosalicylates	Olsalazine	capsules 250mg	Unspecified	Specialist INITIATION	
1. Gastro-intestinal system	1.5 Chronic bowel disorders	1.5.1 Aminosalicylates	Balsalazide	capsules 750 mg	Unspecified	Specialist INITIATION	
1. Gastro-intestinal system	1.5 Chronic bowel disorders	1.5.2 Corticosteroids	Beclometasone	5mg gastro-resistant modified-release tablets	Second line	Specialist INITIATION	To be used within its marketing authorisation. Hospitals should provide the full course (therapy cycles of not more than four weeks are recommended) but there may be instances during a pandemic when this is not possible and GPs may prescribe.
1. Gastro-intestinal system	1.5 Chronic bowel disorders	1.5.2 Corticosteroids	Budesonide	e/c MR capsules 3mg; 2mg/dose rectal foam enema (Budenofalk [®]); 4mg suppositories (Budenofalk [®])	Unspecified	Specialist INITIATION	First line if corticosteroid foam enema required.
1. Gastro-intestinal system	1.5 Chronic bowel disorders	1.5.2 Corticosteroids	Prednisolone	retention enema, short tube, 20mg in 100ml (as sodium phosphate); foam enema, 20mg per metered application (as sodium metasulphobenzoate); suppositories 5mg (as sodium phosphate)	Unspecified	Specialist INITIATION	Prednisolone foam enema costs much more than budesonide foam enema Budesonide 2mg foam enema (Budenofalk) is first line. Prednisolone (sodium phosphate) suppositories costs much more than Budesonide suppositories (Budenofalk [®])
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.1 Bulk-forming laxatives	Ispaghula husk	3.5g effervescent granules sachets gluten free sugar free	Unspecified	All prescribers	
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.2 Stimulant laxatives	Senna	tablets total sennosides 7.5mg; syrup total sennosides 7.5mg in 5ml	Unspecified	All prescribers	All prescribers
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.2 Stimulant laxatives	Glycerol (glycerin)	suppositories, 1g (infant), 2g (child), 4g (adult);	Unspecified	All prescribers	
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.2 Stimulant laxatives	Docusate sodium	capsules 100mg; oral solution 12.5mg in 5ml, 50mg in 5ml	Unspecified	All prescribers	
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.2 Stimulant laxatives	Co-danthramer	suspension 25/200 (dantron 25 mg, poloxamer '188' 200 mg) in 5ml; strong suspension 75/1000 (dantron 75 mg, poloxamer '188' 1 g) in 5ml	Unspecified	All prescribers	Only licensed for constipation in terminally ill patients.
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.2 Stimulant laxatives	Bisacodyl	e/c tablets 5mg; suppositories 10mg; paediatric suppositories 5mg	Unspecified	All prescribers	
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.4 Osmotic laxatives	Lactulose	solution	Unspecified	All prescribers	
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.4 Osmotic laxatives	Macrogol	Macrogol compound oral powder sachets NPF sugar free Macrogol compound half-strength oral powder sachets NPF sugar free	Unspecified	All prescribers	Do NOT prescribe as Macrogol 3350 oral powder 8.5g sachets sugar free (TransiSoft)
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.4 Osmotic laxatives	Phosphates	Cleen Ready-to-use 133ml enema	Unspecified	All prescribers	
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.4 Osmotic laxatives	Micralax Micro-enema®	enema	Unspecified	All prescribers	
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.6 Peripheral opioid-receptor antagonists	Naloxegol	tablets 12.5mg, 25mg	Unspecified	All prescribers	For use in line with NICE's guidance
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.7 Other drugs used in constipation	Linacotide	capsules 290 microgram	Unspecified	Specialist INITIATION	
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.7 Other drugs used in constipation	Prucalopride	tablets 1mg, 2mg	Unspecified	All prescribers	For use in line with NICE's guidance

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
1. Gastro-intestinal system	1.6 Laxatives - see BNF for comments on the role of laxatives	1.6.7 Other drugs used in constipation	Naldemedine	200microgram tablets	Unspecified	All prescribers	For use in line with NICE's guidance
1. Gastro-intestinal system	1.7 Local preparations for anal and rectal disorders	1.7.1 Soothing haemorrhoidal preparations	Anusol	Anusol Suppositories, Ointment, Cream	First line	All prescribers	Can be purchased over the counter.
1. Gastro-intestinal system	1.7 Local preparations for anal and rectal disorders	1.7.2 Compound haemorrhoidal preparations with corticosteroids	Xyloproct ®	Ointment (water-miscible), aluminium acetate 3.5%, hydrocortisone acetate 0.275%, lidocaine 5%, zinc oxide 18%	First line	All prescribers	First choice if corticosteroid required
1. Gastro-intestinal system	1.7 Local preparations for anal and rectal disorders	1.7.2 Compound haemorrhoidal preparations with corticosteroids	Scheriproct®	Suppositories, cinchocaine (dibucaine) hydrochloride 1mg, prednisolone hexanoate 1.3mg Ointment, cinchocaine (dibucaine) hydrochloride 0.5%, prednisolone hexanoate 0.19%.	Second line	All prescribers	Second choice if corticosteroid containing suppositories required
1. Gastro-intestinal system	1.7 Local preparations for anal and rectal disorders	1.7.4 Management of anal fissures	Diltiazem	rectal ointment 2% (unlicensed)	Second line	All prescribers	For treatment of anal fissure. Generally not suitable for repeat prescribing
1. Gastro-intestinal system	1.7 Local preparations for anal and rectal disorders	1.7.4 Management of anal fissures	Glyceryl trinitrate	rectal ointment 0.4% (Rectogesic ®)	First line	All prescribers	For treatment of anal fissure. Generally not suitable for repeat prescribing
1. Gastro-intestinal system	1.9 Drugs affecting intestinal secretions	1.9.1	Ursodeoxycholic acid	tablets 150mg; tablets 250mg	Unspecified	Specialist INITIATION	
1. Gastro-intestinal system	1.9 Drugs affecting intestinal secretions	1.9.4	Creon®	Creon® 10 000 Capsules,e/c granules of pancreatin (pork), providing: protease 600 units, lipase 10 000 units, amylase 8000 units; 25 000 capsules	Unspecified	Specialist INITIATION	Please refer to NW London Briefing document: Pancreatic Enzyme Replacement Therapy (PERT) shortages https://www.nwlondonicb.nhs.uk/professionals/medicines/medicines-shortages
2. Cardiovascular system	2.1 Positive inotropic drugs	2.1.1	Digoxin	tablets 62.5mcg, 125mcg, 250mcg; elixir 50mcg in 1ml;	Unspecified	All prescribers	
2. Cardiovascular system	2.2 Diuretics	2.2.1 Thiazides and related diuretics	Indapamide	tablets 2.5mg; MR tablets 1.5mg	First line	All prescribers	
2. Cardiovascular system	2.2 Diuretics	2.2.1 Thiazides and related diuretics	Bendroflumethiazide	tablets 2.5mg, 5mg	Unspecified	All prescribers	
2. Cardiovascular system	2.2 Diuretics	2.2.1 Thiazides and related diuretics	Metolazone	tablets 5mg (Xaqua ®)	Unspecified	Specialist INITIATION	For use oedema in heart failure only. Prescribe by brand name Specialist to prescribe the initial dose titration until optimised, which will usually be around after 4 weeks
2. Cardiovascular system	2.2 Diuretics	2.2.2 Loop diuretics	Furosemide	tablets 20mg, 40mg, 500mg; oral solution 20mg in 5ml, 40mg in 5ml, 50mg in 5ml;	Unspecified	All prescribers	
2. Cardiovascular system	2.2 Diuretics	2.2.2 Loop diuretics	Furosemide	solution for injection	Unspecified	Specialist INITIATION	Can be initiated in primary care following specialist recommendation as subcutaneous therapy for patients with advanced heart failure for palliative care purposes.
2. Cardiovascular system	2.2 Diuretics	2.2.2 Loop diuretics	Bumetanide	tablets 1mg, 5mg; Liquid 1mg in 5ml.	Unspecified	All prescribers	
2. Cardiovascular system	2.2 Diuretics	2.2.2 Loop diuretics	Torsemide	tablets	See comments column	Specialist INITIATION	
2. Cardiovascular system	2.2 Diuretics	2.2.3 Potassium-sparing diuretics and aldosterone antagonists	Eplerenone	tablets 25mg, 50mg	Unspecified	All prescribers	
2. Cardiovascular system	2.2 Diuretics	2.2.3 Potassium-sparing diuretics and aldosterone antagonists	Spirinolactone	tablets 25mg, 100mg; 50 mg	Unspecified	All prescribers	
2. Cardiovascular system	2.2 Diuretics	2.2.3 Potassium-sparing diuretics and aldosterone antagonists	Amiloride	tablets 5mg; oral solution 5mg in 5ml	Unspecified	All prescribers	
2. Cardiovascular system	2.2 Diuretics	2.2.3 Potassium-sparing diuretics and aldosterone antagonists	Finerenone	tablets 10mg, 20mg	Unspecified	Specialist INITIATION	To be used only in line with NICE Technology appraisal guidance [TA877]. Minimum six months' stabilisation by specialist before transfer of prescribing to primary care
2. Cardiovascular system	2.2 Diuretics	2.2.4 Potassium-sparing diuretics with other diuretics	Co-amilozide	tablets 2.5/25 (amiloride hydrochloride 2.5 mg, hydrochlorothiazide 25 mg), 5/50 (amiloride hydrochloride 5 mg, hydrochlorothiazide 50 mg)	Unspecified	All prescribers	
2. Cardiovascular system	2.2 Diuretics	2.2.4 Potassium-sparing diuretics with other diuretics	Co-amilofruse	tablets 2.5/20 (amiloride hydrochloride 2.5 mg, furosemide 20 mg), 5/40 (amiloride hydrochloride 5 mg, furosemide 40 mg)	Unspecified	All prescribers	
2. Cardiovascular system	2.3 Anti-arrhythmic drugs	2.3.2	Propafenone	tablets 150mg, 300mg	Unspecified	Specialist INITIATION	
2. Cardiovascular system	2.3 Anti-arrhythmic drugs	2.3.2	Flecainide	tablets 50mg, 100mg;	Unspecified	Specialist INITIATION	
2. Cardiovascular system	2.3 Anti-arrhythmic drugs	2.3.2	Disopyramide	capsules 100mg	Unspecified	Specialist INITIATION	
2. Cardiovascular system	2.3 Anti-arrhythmic drugs	2.3.2	Amiodarone	tablets 100mg, 200mg;	Unspecified	Specialist INITIATION	For use in line with a shared care protocol.
2. Cardiovascular system	2.3 Anti-arrhythmic drugs	2.3.2	Dronedarone	tablets 400mg	Unspecified	Specialist INITIATION	For use in line with the NICE Technology appraisal guidance [TA197] and a shared care protocol.
2. Cardiovascular system	2.4 Beta-adrenoceptor blocking drugs	2.4	Sotalol	tablets 40mg, 80mg, 160mg, 200mg	Unspecified	All prescribers	
2. Cardiovascular system	2.4 Beta-adrenoceptor blocking drugs	2.4	Nebivolol	tablets 5mg	Unspecified	All prescribers	
2. Cardiovascular system	2.4 Beta-adrenoceptor blocking drugs	2.4	Labetalol	tablets 50mg, 100mg, 200mg, 400mg	Unspecified	All prescribers	
2. Cardiovascular system	2.4 Beta-adrenoceptor blocking drugs	2.4	Bisoprolol	tablets 1.25mg, 2.5mg, 3.75mg, 5mg, 7.5mg, 10mg	First line	All prescribers	
2. Cardiovascular system	2.4 Beta-adrenoceptor blocking drugs	2.4	Carvedilol	tablets 3.125mg, 6.25mg, 12.5mg, 25mg	Second line	All prescribers	
2. Cardiovascular system	2.4 Beta-adrenoceptor blocking drugs	2.4	Propranolol	tablets 10mg, 40mg, 80mg,160mg; MR capsules 80mg, 160mg; oral solution 5mg in 5ml, 10mg in 5ml, 50mg in 5ml	Unspecified	All prescribers	
2. Cardiovascular system	2.4 Beta-adrenoceptor blocking drugs	2.4	Metoprolol	tablets 50mg, 100mg;	Unspecified	All prescribers	
2. Cardiovascular system	2.4 Beta-adrenoceptor blocking drugs	2.4	Atenolol	tablets 25mg, 50mg, 100mg; syrup 25mg in 5ml;	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.1 Vasodilator antihypertensive drugs	Hydralazine	tablets 25mg, 50mg;	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.1 Vasodilator antihypertensive drugs	Minoxidil	tablets 2.5mg, 5mg, 10mg	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.2 Centrally acting antihypertensive drugs	Moxonidine	tablets 200mcg, 300mcg, 400mcg	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.2 Centrally acting antihypertensive drugs	Methyldopa	tablets 125mg, 250mg, 500mg	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.4 Alpha-adrenoceptor blocking drugs	Phenoxybenzamine	capsules 10mg	Unspecified	Specialist INITIATION	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.4 Alpha-adrenoceptor blocking drugs	Doxazosin	tablets 1mg, 2mg, 4mg	First line	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5 Other drugs for heart failure	Dapagliflozin	tablets 5mg, 10mg	Unspecified	Specialist INITIATION	Generic dapagliflozin is the preferred SGLT-2i for the treatment of type 2 diabetes (with or without chronic kidney disease) and/or heart failure in line with its marketing authorisation. For use in line with NICE TA679 and NICE TA902
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5 Other drugs for heart failure	Empagliflozin	tablets 10mg, 25mg	Unspecified	Specialist INITIATION	For use in line with NICE TA773 and NICE TA929
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.1 Angiotensin-converting enzyme inhibitors	Ramipril	capsules or tablets 1.25mg, 2.5mg, 5mg, 10mg	First line	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.1 Angiotensin-converting enzyme inhibitors	Lisinopril	tablets 2.5mg, 5mg, 10mg, 20mg	First line	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.1 Angiotensin-converting enzyme inhibitors	Perindopril erbumine	tablets 2mg, 4mg, 8mg	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.1 Angiotensin-converting enzyme inhibitors	Enalapril	tablets 2.5mg, 5mg, 10mg, 20mg	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.1 Angiotensin-converting enzyme inhibitors	Captopril	tablets 12.5mg, 25mg, 50mg	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.2 Angiotensin-II receptor antagonists	Losartan	tablets 25mg, 50mg, 100mg, 12.5 mg	First line	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.2 Angiotensin-II receptor antagonists	Valsartan	capsules 40mg, 80mg, 160mg	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.2 Angiotensin-II receptor antagonists	Irbesartan	tablets 75mg, 150mg, 300mg	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.2 Angiotensin-II receptor antagonists	Candesartan	tablets 2mg, 4mg, 8mg, 16mg, 32mg	Unspecified	All prescribers	
2. Cardiovascular system	2.5 Hypertension and heart failure	2.5.5.2 Angiotensin-II receptor antagonists	Sacubitril/valsartan	tablets 24mg/26mg, 49mg/51mg, 97mg/103mg (Entresto®)	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.1 Nitrates	Isosorbide mononitrate	tablets 10mg, 20mg; MR capsules 25mg, 50mg; MR tablets 25mg, 40mg, 60mg	Unspecified	All prescribers	
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.1 Nitrates	Glycerol trinitrate	aerosol spray 400mcg/metered spray; tablets 500 micrograms	Unspecified	All prescribers	The tablets should be discarded after 8 weeks in use, so the spray generally provides better value for money
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.2 Calcium-channel blockers	Verapamil	tablets 40mg, 80mg, 120mg; MR capsules 120mg; MR tablets 240mg; oral solution 40mg in 5ml	Unspecified	All prescribers	
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.2 Calcium-channel blockers	Nifedipine	capsules 5mg, 10mg; Modified release preparations - prescribers should specify the brand to be dispensed (see BNF)	Unspecified	All prescribers	Potential confusion between MR brands probably means that a drug that does not require modified release formulations, such as amlodipine, is preferable for treatment of hypertension. If using MR nifedipine, prescribe by brand
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.2 Calcium-channel blockers	Diltiazem	Standard MR tablets 60mg; Longer-acting formulations - prescriber should specify the brand to be dispensed (see BNF).	Unspecified	All prescribers	Prescribe longer-acting formulations by brand name
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.2 Calcium-channel blockers	Amlodipine	tablets 5mg, 10mg	Unspecified	All prescribers	
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.3 Other antihypertensive drugs	Nicorandil	tablets 10mg, 20mg	Unspecified	All prescribers	
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.3 Other antihypertensive drugs	Ivabradine	tablets 5mg, 7.5mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.3 Other antihypertensive drugs	Ranolazine	tablets	Unspecified	Specialist INITIATION	
2. Cardiovascular system	2.6 Nitrates, Calcium-channel blockers and other antihypertensive drugs	2.6.4 Peripheral vasodilators and related drugs	Naftidrofuryl	capsules 100mg	Unspecified	All prescribers	For use in line with NICE's guidance
2. Cardiovascular system	2.7 Sympathomimetics	2.7.2 Vasoconstrictor sympathomimetics	Midodrine	tablets 2.5mg, 5mg and 10mg	Unspecified	Specialist INITIATION	For use in the treatment of severe orthostatic hypotension due to dysfunction of the autonomic nervous system when corrective factors have been ruled out (licensed indication). Clear monitoring requirements including monitoring for supine hypertension should be communicated to primary care when prescribing responsibility is transferred.

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.1 Parenteral anticoagulants	Tinzaparin	injection syringe 2500 units in 0.25ml, 3500 units in 0.35ml, 4500 units in 0.45ml, 8,000 units in 0.8ml, 10,000 units 0.5ml, 12,000 units in 0.6ml, 14,000 units in 0.7ml, 16,000 units in 0.8ml, 18,000 units in 0.9ml	Unspecified	Specialist INITIATION	
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.1 Parenteral anticoagulants	Heparin sodium or calcium	Subcutaneous injection 5,000units in 0.2ml; pre-filled syringes 5,000units in 0.2ml.	Unspecified	Specialist INITIATION	
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.1 Parenteral anticoagulants	Enoxaparin	injection syringe 20mg in 0.2ml, 40mg in 0.4ml, 60mg in 0.6ml, 80mg in 0.8ml, 100mg in 1ml, 120mg in 0.8ml, 150mg in 1ml	Unspecified	Specialist INITIATION	
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.1 Parenteral anticoagulants	Dalteparin	injection syringe 2500 units in 0.2ml, 5000 units in 0.2ml, 7500 units in 0.3ml, 10,000 units in 0.4ml, 12,500 units in 0.5ml, 15,000 units in 0.6ml, 18,000 units in 0.72ml	Unspecified	Specialist INITIATION	
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.2 Oral anticoagulants	Phenindione	tablets 10mg, 25mg, 50mg	Second line	Specialist INITIATION	
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.2 Oral anticoagulants	Apixaban	tablets 2.5mg, 5mg	First line	All prescribers	For patients commencing treatment for AF, generic apixaban or generic rivaroxaban are the 1st choice joint best value DOACs as per NHS England commissioning recommendations for national procurement for direct-acting oral anticoagulant(s). If this is contraindicated or not clinically appropriate for the specific patient then, subject to the criteria specified in the relevant NICE technology appraisal guidance, clinicians should then consider the next highest best value ranked DOAC. https://www.england.nhs.uk/long-read/commissioning-recommendations-for-national-procurement-for-doacs/ See NHS England commissioning recommendations for national procurement for direct-acting oral anticoagulant(s).
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.2 Oral anticoagulants	Edoxaban	tablets 15mg, 30mg, 60mg	Second line	All prescribers	https://www.england.nhs.uk/long-read/commissioning-recommendations-for-national-procurement-for-doacs/ For patients commencing treatment for AF, generic apixaban or generic rivaroxaban are the 1st choice joint best value DOACs as per NHS England commissioning recommendations for national procurement for direct-acting oral anticoagulant(s).
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.2 Oral anticoagulants	Rivaroxaban	tablets 2.5mg, 10mg, 15mg, 20mg	First line	All prescribers	https://www.england.nhs.uk/long-read/commissioning-recommendations-for-national-procurement-for-doacs/ For patients commencing treatment for AF, generic apixaban or generic rivaroxaban are the 1st choice joint best value DOACs as per NHS England commissioning recommendations for national procurement for direct-acting oral anticoagulant(s). If this is contraindicated or not clinically appropriate for the specific patient then, subject to the criteria specified in the relevant NICE technology appraisal guidance, clinicians should then consider the next highest best value ranked DOAC. https://www.england.nhs.uk/long-read/commissioning-recommendations-for-national-procurement-for-doacs/ See NHS England commissioning recommendations for national procurement for direct-acting oral anticoagulant(s).
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.2 Oral anticoagulants	Dabigatran etexilate	capsules, 110mg, 150mg	Unspecified	All prescribers	https://www.england.nhs.uk/long-read/commissioning-recommendations-for-national-procurement-for-doacs/ See NHS England commissioning recommendations for national procurement for direct-acting oral anticoagulant(s).
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.2 Oral anticoagulants	Warfarin	tablets 500mcg, 1mg, 3mg, 5mg	Unspecified	All prescribers	https://www.england.nhs.uk/long-read/commissioning-recommendations-for-national-procurement-for-doacs/
2. Cardiovascular system	2.8 Anticoagulants and protamine	2.8.2 Oral anticoagulants	Acenocoumarol	tablets 1mg	Unspecified	All prescribers	
2. Cardiovascular system	2.9 Antiplatelet drugs	2.9 Antiplatelet drugs	Prasugrel	tablets 5mg, 10mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
2. Cardiovascular system	2.9 Antiplatelet drugs	2.9 Antiplatelet drugs	Dipyridamole	tablets 25mg, 100mg; MR capsules 200mg; oral liquid 50mg in 5ml	Unspecified	All prescribers	For use in line with NICE's guidance
2. Cardiovascular system	2.9 Antiplatelet drugs	2.9 Antiplatelet drugs	Clopidogrel	tablets 75mg	Unspecified	All prescribers	For use in line with NICE's guidance
2. Cardiovascular system	2.9 Antiplatelet drugs	2.9 Antiplatelet drugs	Aspirin	tablets dispersible 75mg tablets 75mg	Unspecified	All prescribers	
2. Cardiovascular system	2.9 Antiplatelet drugs	2.9 Antiplatelet drugs	Ticagrelor	tablets 60mg, 90mg	See comments column	Specialist INITIATION	For use in line with NICE's guidance
2. Cardiovascular system	2.11 Antifibrinolytic drugs and haemostatics	2.11 Antifibrinolytic drugs and haemostatics	Tranexamic acid	tablets 500mg	Unspecified	All prescribers	
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Simvastatin	tablets 10mg, 20mg, 40mg, 80mg	Unspecified	All prescribers	
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Colestyramine (cholestyramine)	powder 4g per sachet (Questran Light ®); powder, 4g sachets (Questran ®)	Unspecified	All prescribers	
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Rosuvastatin	tablets 5mg, 10mg, 20mg, 40mg	Unspecified	All prescribers	Tablet formulation is more cost effective than capsules
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Colesevelam	tablets 625mg	Unspecified	All prescribers	Costs more than colestyramine, but the latter may not be tolerated.
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Atorvastatin	tablets 10mg, 20mg, 40mg, 80mg	Unspecified	All prescribers	
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Pravastatin	tablets 10mg, 20mg, 40mg	Unspecified	All prescribers	
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Fenofibrate	capsules (micronised) 67mg, 200mg, 267mg; tablets (micronised) 160mg	Unspecified	All prescribers	
	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Bempedoic acid	Bempedoic acid 180mg tablets; Bempedoic acid 180mg / Ezetimibe 10mg tablets	Unspecified	All prescribers	Only for use as per NICE Technology Appraisal Guidance [TA694] which recommends bempedoic acid with ezetimibe if: • statins are contraindicated or not tolerated, • ezetimibe alone does not control low-density lipoprotein cholesterol well enough
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Ezetimibe	tablets 10mg	Unspecified	All prescribers	
	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Incisiran	284mg/1.5ml solution for injection pre-filled syringes	Unspecified	All prescribers	For use in line with NICE's guidance [TA733] and NW London lipid guidelines: https://www.nwondonicb.nhs.uk/professionals/referral-guidelines-and-clinical-documents/lipids Not to be used in adults who have no history of cardiovascular events The Incisiran studies were not powered to show patient oriented outcomes such as cardiovascular events or death. A further study is underway to measure the benefit of Incisiran on cardiovascular outcomes.
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Bezafibrate	tablets 200mg; MR tablets 400mg	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Omega-3 fatty acid	capsules	Unspecified	Specialist INITIATION	For patients with hypertriglyceridaemia after any secondary causes have been addressed, capsules containing docosahexaenoic acid 380mg + eicosapentenoic acid 460mg can be used to reduce the risk of pancreatitis if the patient meets the following criteria: • Under the care of a lipid specialist and • Taking maximum tolerated fibrate and statin and • Fasting triglycerides remain >10mmol/l.
2. Cardiovascular system	2.12 Lipid-regulating drugs	2.12 Lipid-regulating drugs	Icosapent	capsules 998mg	Unspecified	Specialist INITIATION	For use in line with the NICE Technology appraisal guidance [TA805], for secondary prevention and when low-density lipoprotein cholesterol levels criteria is met (not to be used for primary prevention).
2. Cardiovascular system	2.14 Other miscellaneous	2.14 Other miscellaneous	Semaglutide	0.25mg/0.19ml solution for injection 1.5ml pre-filled disposable device; 0.5mg/0.37ml solution for injection 1.5ml pre-filled disposable device; 1mg/0.74ml solution for injection 3ml pre-filled disposable device; 1.7mg/0.75ml solution for injection 3ml pre-filled disposable device; 2.4mg/0.75ml solution for injection 3ml pre-filled disposable device	Unspecified	See comments column	For reducing the risk of major adverse cardiovascular events in people with cardiovascular disease and overweight or obesity (NICE TA 1152) Final formulary status is pending the development of a clinical pathway and agreement of the commissioning framework
3. Respiratory system	3.1 Bronchodilators	3.1.1.1 Selective beta2 agonists	Indacaterol	150 microgram inhalation powder, hard capsules; 300 microgram inhalation powder, hard capsules	Unspecified	Specialist INITIATION	
3. Respiratory system	3.1 Bronchodilators	3.1.1.1 Selective beta2 agonists	Salbutamol	Salamol 100micrograms/dose inhaler CFC Airomir 100micrograms/dose inhaler (Second line after Salamol MDI) Easyhaler Salbutamol sulfate 100micrograms/dose dry powder inhaler Ventolin 200micrograms/dose Accuhaler Airomir 100micrograms/dose Autohaler Salamol 100micrograms/dose Easi-Breathe inhaler Salbutamol 2.5mg/2.5ml nebuliser liquid unit dose vials Salbutamol 5mg/2.5ml nebuliser liquid unit dose vials	First line	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.1.1 Selective beta2 agonists	Terbutaline	Turbohaler 500mcg/inhalation; Respules 5mg in 2ml	Second line	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.1.1 Selective beta2 agonists	Salmeterol	aerosol inhalation 25mcg/metered inhalation; Accuhaler 50mcg/blister;	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.1.1 Selective beta2 agonists	Formoterol	Easyhaler (dry powder for inhalation) 12micrograms/metered inhalation; Alimos Modulite (aerosol inhalation) 12micrograms/metered inhalation; Oxis (turbohaler) 6mcg/inhalation, 12mcg/inhalation; Foradil (dry powder for inhalation) inhalation capsules 12mcg	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.2 Antimuscarinic bronchodilators	Tiotropium	Spiriva® (dry powder inhaler) 18microgram capsule; Braltus ® (dry powder inhaler) 10microgram capsule; Respimat® solution for inhalation 2.5micrograms/metered inhalation	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.2 Antimuscarinic bronchodilators	Acclidinium bromide	Dry powder inhaler 375 mcgs(= acclidinium 322 mcg)/inhalation (delivered dose) for use with Eklira Genuair® device	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.2 Antimuscarinic bronchodilators	Glycopyrronium bromide	Inhalation powder 50 mcg, hard capsule, for use with Seebri Breezhaler® device	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.2 Antimuscarinic bronchodilators	Ipratropium bromide	aerosol inhalation 20mcg/metered inhalation; nebuliser solution 250mcg in 1ml, 500mcg in 2ml unit dose vials	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.2 Antimuscarinic bronchodilators	Umeclidinium	inhalation powder 55 micrograms (Incruse Ellipta)	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.4 Compound bronchodilators	Anoro Ellipta ®	dry powder for inhalation umecclidinium 55micrograms + vilanterol 22micrograms	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.4 Compound bronchodilators	Duaklir Genuair ®	dry powder for inhalation acclidinium 340micrograms + formoterol 12micrograms	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.4 Compound bronchodilators	Spiolto Respimat ®	solution for inhalation tiotropium 2.5micrograms + olodaterol 2.5micrograms	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.4 Compound bronchodilators	Ultibro Breezhaler ®	dry powder for inhalation indacaterol 85micrograms + glycopyrronium 43micrograms	Unspecified	All prescribers	
3. Respiratory system	3.1 Bronchodilators	3.1.3 Theophylline	Theophylline	MR tablets (Uniphyllin Continus) 200mg, 300mg, 400mg; MR capsules 60mg, 125mg, 250mg (Slo-Phyllin)	Unspecified	All prescribers	Rate of absorption can vary between brands. Maintain the patient on the brand s/he has been stabilised on.
3. Respiratory system	3.1 Airways disease, obstructive	3.1 Phosphodiesterase Type-4 inhibitors	Roflumilast	tablets 500 micrograms	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
3. Respiratory system	3.1.5 Peak flow meters, inhaler devices and nebulisers	3.1.5	Drug delivery devices	spacer devices	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
3. Respiratory system	3.1.5 Peak flow meters, inhaler devices and nebulisers	3.1.5	Peak flow meters	Vitalograph® (standard & child)	Unspecified	All prescribers	
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Fostair ®	aerosol inhalation 100/6, 200/6 (beclometasone dipropionate 100mcg or 200mcg, formoterol fumarate 6mcg per metered inhalation). NEXThaler 100/6, 200/6 (dry powder inhaler containing beclometasone dipropionate 100mcg or 200mcg plus formoterol fumarate 6mcg per metered inhalation.)	Unspecified	All prescribers	Substitute Fostair 100/6, 200/6 pMDI with the like for like generic equivalent Bibecfo or Luforbec pMDI if Fostair NEXThaler (DPI) is not suitable. No new patients should be started on Fostair pMDI
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Luforbec®	aerosol inhalation 100/6, 200/6 (beclometasone dipropionate 100mcg or 200mcg, formoterol fumarate 6mcg per metered inhalation)	Unspecified	All prescribers	Luforbec pMDI is the generic like for like equivalent to Fostair pMDI and costs considerably less
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Bibecfo®	aerosol inhalation 100/6, 200/6 (beclometasone dipropionate 100mcg or 200mcg, formoterol fumarate 6mcg per metered inhalation)	Unspecified	All prescribers	Bibecfo pMDI is the generic like for like equivalent to Fostair pMDI and costs considerably less
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Symbicort	Turbohaler 100/6, 200/6, 400/12 dry powder for inhalation 160mcg/4.5mcg and 320mcg/9mcg. Containing budesonide (160mcg and 320mcg) and formoterol (4.5mcg and 9mcg)	Unspecified	All prescribers	
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	DuoResp Spiromax®	Budesonide 80micrograms/dose / Formoterol 4.5micrograms/dose dry powder inhaler	Unspecified	All prescribers	
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Fobumix Easyhaler ®	Budesonide 160micrograms/dose / Formoterol 4.5micrograms/dose dry powder inhaler Budesonide 320micrograms/dose / Formoterol 9micrograms/dose dry powder inhaler	Unspecified	All prescribers	Fobumix Easyhaler is the preferred option and most cost effective DPI for MART indication
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Flutiform ®	aerosol inhalation 50 and 125 (aerosol containing 5mcg formoterol plus 50/125 mcg fluticasone/metered inhalation. aerosol inhalation 250 (aerosol containing 10mcg formoterol plus 250 mcg fluticasone/metered inhalation)	Unspecified	All prescribers	
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Relvar Ellipta ®	dry powder for inhalation 92mcg/22mcg and 184mcg/22mcg. Containing vilanterol 22micrograms plus fluticasone furoate 92/184micrograms per delivered dose.	Unspecified	All prescribers	
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Trelegy Ellipta ®	Inhalation powder, pre-dispensed 92 micrograms/55 micrograms/22 micrograms	Unspecified	Specialist INITIATION	Seek specialist advice before escalating to triple therapy
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Trimbow ®	87micrograms/dose / 5micrograms/dose / 9micrograms/dose inhaler 172micrograms/dose / 5micrograms/dose / 9micrograms/dose inhaler NEXThaler 88micrograms/dose / 5micrograms/dose / 9micrograms/dose dry powder inhaler	Unspecified	Specialist INITIATION	Seek specialist advice before escalating to triple therapy
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Sereflo ®	Sereflo 25micrograms/dose / 125micrograms/dose inhaler Sereflo 25micrograms/dose / 250micrograms/dose inhaler Sereflo Ciphaler 50micrograms/dose / 250micrograms/dose dry powder inhaler Sereflo Ciphaler 50micrograms/dose / 500micrograms/dose dry powder inhaler	Unspecified	All prescribers	Costs significantly less than the Seretide (DPI and MDI) Ciphaler device is identical to the device of originator brand Accuhaler and this inhaler is authorised based on equivalence to reference product
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Seretide ®	Accuhaler 100, 250, 500 (dry powder breath-actuated disc inhaler containing 50mcg salmeterol plus 100/250/500mcg fluticasone) Evohaler 50, 125, 250 (aerosol containing 25mcg salmeterol plus 50/125/250mcg fluticasone/metered inhalation)	Unspecified	All prescribers	
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Fluticasone	Evohaler ® aerosol inhalation 50mcg/metered inhalation, 125mcg/metered inhalation, 250mcg/metered inhalation; Accuhaler ® 50mcg/blister, 100mcg/blister, 250mcg/blister, 500mcg/blister	Unspecified	All prescribers	
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Budesonide	Turbohaler 100mcg/inhalation, 200mcg/inhalation, 400mcg/inhalation; Respules 500mcg in 2ml, 1mq in 2ml	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Beclometasone dipropionate	Easyhaler Beclometasone 200micrograms/dose dry powder inhaler Kelhale 100micrograms/dose inhaler Clenil Modulite 50micrograms/dose inhaler Clenil Modulite 100micrograms/dose inhaler Clenil Modulite 200micrograms/dose inhaler Clenil Modulite 250micrograms/dose inhaler Qvar 50 Autohaler Qvar 100 Autohaler Qvar 50micrograms/dose Easi-Breathe inhaler Qvar 100micrograms/dose Easi-Breathe inhaler	Unspecified	All prescribers	Kelhale, Qvar and Fostair contain extrafine particle beclometasone which is approximately TWICE as potent as beclometasone in Clenil & Easyhaler Beclometasone.
3. Respiratory system	3.2 Corticosteroids	3.2 Corticosteroids	Mometasone	dry powder inhaler 200micrograms/metered inhalation, 400micrograms/metered inhalation	Exceptional case	All prescribers	
3. Respiratory system	3.03 Cromoglicate and related therapy and leukotriene receptor antagonists	3.3.3	Montelukast	tablets 10mg; chewable tablets 4mg, 5mg granules 4mg	Unspecified	All prescribers	
3. Respiratory system	3.4 Antihistamines, hyposensitisation, and allergic emergencies	3.4.1	Cetirizine	tablets 10mg; oral solution sugar free 1mg/ml	Unspecified	All prescribers	
3. Respiratory system	3.4 Antihistamines, hyposensitisation, and allergic emergencies	3.4.1	Loratadine	tablets 10mg; syrup 5mg in 5ml	Unspecified	All prescribers	
3. Respiratory system	3.4 Antihistamines, hyposensitisation, and allergic emergencies	3.4.1	Promethazine hydrochloride	tablets 10mg, 25mg; elixir 5mg in 5ml	Unspecified	All prescribers	
3. Respiratory system	3.4 Antihistamines, hyposensitisation, and allergic emergencies	3.4.1	Hydroxyzine	tablets 10mg, 25mg	Unspecified	All prescribers	
3. Respiratory system	3.4 Antihistamines, hyposensitisation, and allergic emergencies	3.4.1	Chlorphenamine	tablets 4mg; syrup 2mg in 5ml;	Unspecified	All prescribers	
3. Respiratory system	3.4 Antihistamines, hyposensitisation, and allergic emergencies	3.4.2	House dust mites extract (Acarizax ®) 12 SQ-HDM	oral lyophilisates	Unspecified	Specialist INITIATION	For use in line with NICE TA 1045 Specialist initiation with a transfer of prescribing to primary care after 6 months
3. Respiratory system	3.4 Antihistamines, hyposensitisation, and allergic emergencies	3.4.2	White Birch (Betula verrucosa) pollen allergen extract (Itulazax®) 12 SQ-Bet	oral lyophilisates sugar free	Unspecified	Specialist INITIATION	For use in line with NICE TA 1087 Specialist initiation and prescribing for 6 months and continuation in primary care
3. Respiratory system	3.4 Antihistamines, hyposensitisation, and allergic emergencies	3.4.3	Adrenaline/ epinephrine	pre-filled auto-injector delivering 150 micrograms (EpiPen Junior, Jext), 300 micrograms (EpiPen, Jext)	Unspecified	All prescribers	Prescribe using the brand name
3. Respiratory system	3.7 Mucolytics	3.7	Carbocisteine	capsules 375mg; oral liquid 250mg in 5ml	Unspecified	All prescribers	For use in line with NICE guidance
3. Respiratory system	3.7 Mucolytics	3.7	Domase alfa	nebuliser solution	Unspecified	Specialist INITIATION	GPs should not agree to take on any new prescribing of domase alfa as it is now covered by NHS England specialised commissioning.
3. Respiratory system	3.7 Mucolytics	3.7	Hypertonic sodium chloride	nebuliser solution 3%, 6%, 7%	Unspecified	All prescribers	
3. Respiratory system	3.9 Cough preparations	3.9.1	Codeine	linctus 15mg in 5ml	Unspecified	All prescribers	
3. Respiratory system	3.9 Cough preparations	3.9.2	Simple	linctus, linctus paediatric	Unspecified	All prescribers	
4. Central nervous system	4.1 Hypnotics and anxiolytics	4.1.1 Hypnotics	Zopiclone	tablets 3.75mg, 7.5mg	Unspecified	All prescribers	
4. Central nervous system	4.1 Hypnotics and anxiolytics	4.1.1 Hypnotics	Zolpidem tartrate	tablets 5mg, 10 mg	Unspecified	All prescribers	
4. Central nervous system	4.1 Hypnotics and anxiolytics	4.1.1 Hypnotics	Temazepam	tablets 10mg, 20mg; oral solution 10mg in 5ml	Unspecified	All prescribers	For use in line with NICE guidance. When a short acting hypnotic is indicated NICE recommend the cheapest of short-acting benzodiazepine, zolpidem or zopiclone. Zolpidem and zopiclone are currently the lowest cost drugs.
4. Central nervous system	4.1 Hypnotics and anxiolytics	4.1.2	Lorazepam	tablets 1mg, 2.5mg	Unspecified	All prescribers	
4. Central nervous system	4.1 Hypnotics and anxiolytics	4.1.2	Diazepam	tablets 2mg, 5mg, 10mg; rectal tubes 2.5mg in 1.25ml, 5mg in 2.5ml, 10mg in 2.5ml	Unspecified	All prescribers	
4. Central nervous system	4.1 Hypnotics and anxiolytics	4.1.2	Chlordiazepoxide	capsules 5mg, 10mg	Unspecified	All prescribers	
4. Central nervous system	4.1 Hypnotics and anxiolytics	4.1.2	Buspiron	tablets 5mg, 10mg	Unspecified	All prescribers	
	4.1 Hypnotics and anxiolytics	4.1.4	Daridorexant	tablets 25mg, 50mg	Unspecified	All prescribers	Only for use in line with the NICE TA 922: Daridorexant for treating long-term insomnia 1.1 Daridorexant is recommended for treating insomnia in adults with symptoms lasting for 3 nights or more per week for at least 3 months, and whose daytime functioning is considerably affected, only if: • cognitive behavioural therapy for insomnia (CBTI) has been tried but not worked, or • CBTi is not available or is unsuitable. 1.2 The length of treatment should be as short as possible. Treatment with daridorexant should be assessed within 3 months of starting and should be stopped in people whose long-term insomnia has not responded adequately. If treatment is continued, assess whether it is still working at regular intervals
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Trifluoperazine	tablet 1mg, 5mg; oral solution 5mg in 5ml	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Sulpiride	tablets 200mg, 400mg; oral solution 200mg in 5ml.	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Quetiapine	tablets 25mg, 100mg 150mg, 200mg, 300mg; MR tablets 50mg, 150mg, 200mg, 300mg, 400mg	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance Modified release (MR) tablets cost significantly more than non-MR tablets. Modified release quetiapine is a reasonable choice when: • Nurses visit the patient's home to administer medicines • The patient is known to take quetiapine erratically (as there are likely to be more dose-related side effects from the immediate release formulation). • The patient refuses to take medicines more than once a day. • The patient cannot tolerate their dose when it is given using the immediate release formulation (e.g. because of significant hypotension). • For initial dose titration in hospital (because titration to 600mg/day takes around 7 days with the immediate release formulation but only 2 days with the MR formulation). After titration and before discharge from inpatient care patients should generally be changed from modified release to immediate release quetiapine.
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Olanzapine	tablets 2.5mg, 5mg, 7.5mg, 10mg, 15mg; 20mg orodispersible tablets sugar free 5mg, 10mg, 15mg, 20mg	Unspecified	All prescribers	Avoid lyophilisates sugar free formulation For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Chlorpromazine hydrochloride	tablets 25mg, 50mg, 100mg; oral solution 25mg in 5ml, 100mg in 5ml	Unspecified	Specialist INITIATION	
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Aripiprazole	tablets 5mg, 10mg, 15mg, 30mg; orodispersible tablets 10mg, 15mg oral solution 1mg in 1ml	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Amisulpride	tablets 50mg, 100mg, 200mg, 400mg; oral solution 100mg in 1ml	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Risperidone	tablets 500micrograms, 1mg, 2mg, 3mg, 4mg, 6mg; orodispersible tablets 500micrograms, 1mg, 2mg, 4mg; oral solution 1mg/ml sugar free	Unspecified	All prescribers	Orodispersible tablets cost much more than standard tablets or liquid. For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Haloperidol	tablets 1.5mg, 5mg, 10mg, 20mg; oral liquid 10mg in 5ml; injection 5mg in 1ml	Unspecified	All prescribers	Haloperidol 500microgram tablets costs significantly more than other haloperidol formulations. Consider haloperidol 5mg/5ml oral solution sugar free liquid instead as appropriate
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Flupentixol	tablets 3mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Zuclopenthixol	tablets 2mg, 10mg, 25mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.1 Antipsychotic drugs	Levomepromazine	tablets 25mg, injection 25mg/ml	Unspecified	All prescribers	For use in palliative care
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.2 Antipsychotic depot injections	Aripiprazole	depot injection 400mg	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.2 Antipsychotic depot injections	Flupentixol decanoate	depot injection 20mg/ml, 100mg/ml, 200mg/ml	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.2 Antipsychotic depot injections	Haloperidol (as decanoate)	depot injection 50mg/ml, 100mg/ml	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.2 Antipsychotic depot injections	Paliperidone	monthly injection (Xepion) 50mg, 75mg, 100mg, 150mg; 3 monthly injection (Trevicta) 175mg, 263mg, 350mg, 525mg	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.2 Antipsychotic depot injections	Risperidone	depot injection 25mg, 37.5mg, 50mg	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.2 Antipsychotic depot injections	Zuclopenthixol decanoate	depot injection 200mg/ml, 500mg/ml	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of Prescribing of Antipsychotics Guidance

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
							Do not start valproate for the first time in people (male or female) younger than 55 years, unless 2 specialists independently agree and document that there is no other effective and tolerated treatment, or there are compelling reasons that the reproductive risks do not apply. Ensure the pregnancy prevention programme is in place if valproate is used in women and girls of childbearing potential. Follow the MHRA safety advice on the use of valproate. NICE Bipolar disorder guidance [CG 185] states: If a woman or girl of childbearing potential is already taking valproate, advise her to gradually stop the drug under medical supervision because of the risk of fetal malformations and adverse neurodevelopmental outcomes after any exposure in pregnancy. MHRA Guidance: Valproate use by women and girls Drug Safety Update September 2024: Valproate use in men: as a precaution, men and their partners should use effective contraception Drug Safety Update January 2024: Valproate (Belvo, Convulex, Depakote, Dyzantil, Epilim, Epilim Chrono or Chronosphere, Episenta, Epival, and Syonell▼); new safety and educational materials to support regulatory measures in men and women under 55 years of age Drug Safety Update October 2023: Valproate: dispense full packs of valproate-containing medicines Drug Safety Update August 2023: Valproate: re-analysis of study on risks in children of men taking valproate Drug Safety Update February 2020: Valproate (Epilim▼, Depakote▼) pregnancy prevention programme: updated educational materials Drug Safety Update April 2019: Valproate medicines and serious harms in pregnancy: new Annual Risk Acknowledgement Form and clinical guidance from professional bodies to support compliance with the Pregnancy Prevention Programme Drug Safety Update April 2018: Valproate medicines (Epilim▼, Depakote▼): contraindicated in women and girls of childbearing potential unless conditions of Pregnancy Prevention Programme are met
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.3 Antimanic drugs	Valproic acid	e/c tablets 250mg, 500mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.3 Antimanic drugs	Lithium citrate	liquid 520mg in 5ml (Priadel ®); oral solution 509mg in 5ml (LI-liquid ®);	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of lithium prescribing guideline
4. Central nervous system	4.2 Drugs used in psychoses and related disorders	4.2.3 Antimanic drugs	Lithium carbonate	MR tablets 200mg, 400mg (Priadel ®, Camcolit ®, Liskonum®)	Unspecified	Specialist INITIATION	For use in line with the NW London Transfer of lithium prescribing guideline
4. Central nervous system	4.03 Antidepressant drugs	4.3.1 Tricyclic and related antidepressant drugs	Amitriptyline hydrochloride	tablets 10mg, 25mg, 50mg; oral solution 25mg in 5ml, 50mg in 5ml	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.1 Tricyclic and related antidepressant drugs	Clomipramine	capsules, 10mg, 25mg, 50mg	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.1 Tricyclic and related antidepressant drugs	Trazodone	capsules 50mg, 100mg; tablets 150mg; liquid 50mg in 5ml	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.1 Tricyclic and related antidepressant drugs	Lofepramine	tablets 70mg; oral suspension 70mg in 5ml	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.1 Tricyclic and related antidepressant drugs	Imipramine	tablets 10mg, 25mg	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.1 Tricyclic and related antidepressant drugs	Nortriptyline	tablets 10mg, 25mg	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.2 Monoamine-oxidase inhibitors	Phenelzine	tablets 15mg	Unspecified	Specialist INITIATION	Initiation, discontinuation and dose changes should only be done by a mental health specialist, or under the supervision of a mental health specialist
4. Central nervous system	4.03 Antidepressant drugs	4.3.2 Monoamine-oxidase inhibitors	Moclobemide	tablets 150mg	Unspecified	Specialist INITIATION	Initiation, discontinuation and dose changes should only be done by a mental health specialist, or under the supervision of a mental health specialist
4. Central nervous system	4.03 Antidepressant drugs	4.3.3 Selective serotonin re-uptake inhibitors	Sertraline	tablets 50mg, 100mg	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.3 Selective serotonin re-uptake inhibitors	Fluoxetine	capsules 20mg; liquid 20mg in 5ml dispersible tablets 20mg	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.3 Selective serotonin re-uptake inhibitors	Citalopram	tablets 10mg, 20mg, 40mg; oral solution 40mg/ml sugar free	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.3 Selective serotonin re-uptake inhibitors	Paroxetine	tablets 20mg, 30mg	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.4 Other antidepressant drugs	Duloxetine	capsules 30mg, 60mg	Unspecified	All prescribers	For use in neuropathic pain e.g. In palliative care. Not included in the Integrated Formulary for treatment of depression or generalised anxiety disorder.
4. Central nervous system	4.03 Antidepressant drugs	4.3.4 Other antidepressant drugs	Venlafaxine	tablets 37.5mg, 75mg MR tablets 75mg, 150mg MR capsules 225 mg	Unspecified	All prescribers	Plain tablets cost less than MR formulations.
4. Central nervous system	4.03 Antidepressant drugs	4.3.4 Other antidepressant drugs	Mirtazapine	tablets 15mg, 30mg, 45mg; Orodispersible (soluble) tablets 15mg, 30mg, 45mg	Unspecified	All prescribers	
4. Central nervous system	4.03 Antidepressant drugs	4.3.4 Other antidepressant drugs	Vortioxetine	tablets 5mg, 10mg, 20mg	Unspecified	All prescribers	For use in line with NICE's guidance
4. Central nervous system	4.4 Central nervous system stimulants	4.4 CNS stimulants and drugs used for ADHD	Methylphenidate hydrochloride	tablets 5mg, 10mg; MR capsules 10mg, 20mg, 30mg (Equasym XL ®); MR capsules 10mg, 20mg, 30mg, 40mg (Medikinet XL ®); MR tablets 18mg, 27mg, 36mg, 54mg (Delmosart®); MR tablets 18mg, 27mg, 36mg, 54mg (Affenid ® XL)	Unspecified	Specialist INITIATION	For use in line with NICE's guidance and a shared care protocol. Affenid® XL is the preferred brand for modified-release tablets
4. Central nervous system	4.4 Central nervous system stimulants	4.4 CNS stimulants and drugs used for ADHD	Atomoxetine	Capsules, 10mg, 18mg, 25mg, 40mg, 60mg, 80mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance and a shared care protocol.

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
4. Central nervous system	4.4 Central nervous system stimulants	4.4 CNS stimulants and drugs used for ADHD	Dexamfetamine sulphate	tablets 5mg, 10mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance and a shared care protocol.
4. Central nervous system	4.4 Central nervous system stimulants	4.4 CNS stimulants and drugs used for ADHD	Lisdexamfetamine mesilate	capsules 30mg, 50mg, 70mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance and a shared care protocol.
4. Central nervous system	4.4 Central nervous system stimulants	4.4 CNS stimulants and drugs used for ADHD	Modafinil	tablets 100mg, 200mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.4 Central nervous system stimulants	4.4 CNS stimulants and drugs used for ADHD	Guanfacine	modified-release tablets 1mg, 2mg, 3mg, 4mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance and a shared care protocol.
4. Central nervous system	4.5 Drugs used in the treatment of obesity	4.5.1	Orlistat	capsules 120mg	Unspecified	All prescribers	
4. Central nervous system	4.6 Drugs used in nausea and vertigo	4.6	Prochlorperazine	tablets 5mg, buccal tablets 3mg	Unspecified	All prescribers	
4. Central nervous system	4.6 Drugs used in nausea and vertigo	4.6	Metoclopramide	tablets 10mg; oral solution 5mg in 5ml; injection 10mg in 2ml	Unspecified	All prescribers	
4. Central nervous system	4.6 Drugs used in nausea and vertigo	4.6	Domperidone	tablets 10mg; suspension 5mg in 5ml; tablets 50 mg; injection 50mg in 1ml	Unspecified	All prescribers	
4. Central nervous system	4.6 Drugs used in nausea and vertigo	4.6	Cyclizine	tablets 500micrograms	Unspecified	All prescribers	
4. Central nervous system	4.6 Drugs used in nausea and vertigo	4.6	Hyoscine hydrobromide	tablets 300micrograms	Unspecified	All prescribers	
4. Central nervous system	4.6 Drugs used in nausea and vertigo	4.6	Bethahistine dihydrochloride	tablets 8mg 16mg	Unspecified	All prescribers	
4. Central nervous system	4.6 Drugs used in nausea and vertigo	4.6	Ondansetron	tablets 4mg, 8mg; oral solution 4mg/5ml; injection 2mg/ml	Unspecified	Specialist INITIATION	
4. Central nervous system	4.6 Drugs used in nausea and vertigo	4.6	Doxylamine/Pyridoxine	gastro-resistant tablets 10mg/10mg (Xonvea ®)	Third line	Specialist INITIATION	Third line option (after trial of at least 2 regular anti-emetics) for the management of moderate and severe nausea and vomiting in pregnancy
4. Central nervous system	4.7 Analgesics	4.7.1 Non-opioid analgesics and compound analgesic preparations	Paracetamol	tablets 500mg; effervescent tablets 500mg; oral suspension sugar free 120mg in 5ml, 250mg in 5ml	Unspecified	All prescribers	Suppositories cost much more than tablets and liquids
4. Central nervous system	4.7 Analgesics	4.7.1 Non-opioid analgesics and compound analgesic preparations	Co-dydramol	tablets 10/500	Unspecified	All prescribers	
4. Central nervous system	4.7 Analgesics	4.7.1 Non-opioid analgesics and compound analgesic preparations	Co-codamol	tablets 8/500, 30/500; dispersible tablets 8/500, 30/500	Unspecified	All prescribers	Dispersible tablets cost more than plain tablets
4. Central nervous system	4.7 Analgesics	4.7.1 Non-opioid analgesics and compound analgesic preparations	Aspirin	dispersible tablets 300mg	Unspecified	All prescribers	
4. Central nervous system	4.7 Analgesics	4.7.1 Non-opioid analgesics and compound analgesic preparations	Nefopam	tablets 30mg	Unspecified	All prescribers	
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Fentanyl	patches 12mcg, 25mcg, 37.5mcg, 50mcg, 75mcg and 100mcg/hour all for 72 hours tablets (Sevredol ®) 10mg, 20mg, 50mg; MR tablets (Morphgesic SR ®) 10mg, 30mg, 60mg, 100mg; MR capsules (Zomorph ®) 10mg, 30mg, 60mg, 100mg, 200mg ; MR capsules (MXL ®) 30mg, 60mg, 90mg, 120mg, 150mg, 200mg ; MR tablets (MST ®) 5mg; oral solution 10mg in 5ml, 100mg in 5ml; injections 10mg/ml, 15mg/ml, 20mg/ml, 30mg/ml	Unspecified	All prescribers	Mezolar® is currently the only brand available in the 37.5mcg/hr strength.
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Morphine sulphate	tablets 5mg; injection 10mg/ml capsules 5mg, 10mg, 20mg; MR tablets 5mg, 10mg, 20mg, 40mg, 80mg; oral solution 5mg in 5ml, 50mg in 5ml; injection 10mg in 1ml; 50mg in 1ml	First line	All prescribers	First line oral strong opioid. Reserve MR suspension sachets for people who cannot take MR tablets
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Diamorphine	injection 5mg, 10mg, 30mg, 100mg	Unspecified	All prescribers	
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Methadone	tablets 5mg; injection 10mg/ml	Unspecified	Specialist INITIATION	
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Oxycodone	capsules 5mg, 10mg, 20mg; MR tablets 5mg, 10mg, 20mg, 40mg, 80mg; oral solution 5mg in 5ml, 50mg in 5ml; injection 10mg in 1ml; 50mg in 1ml	Second line	All prescribers	
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Tapentadol	50mg, 75mg; MR tablets 50mg, 100mg, 150mg, 200mg, 250mg	Unspecified	Specialist INITIATION	Included in the IF for restricted use: it should be initiated in secondary care and only used in line with the protocol approved by that hospital's New Drugs Panel or equivalent. For MR preparations, Tadomon ® MR brand is the preferred option as it costs less than other MR tablets
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Tramadol hydrochloride	Capsule 50mg MR tablets 100mg, 150mg, 200mg, MR capsules, 50mg, 100mg, 150mg, 200mg	Unspecified	All prescribers	
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Dihydrocodeine	tablets 30 mg	Unspecified	All prescribers	
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Codeine phosphate	Tablets codeine phosphate 15 mg, 30 mg, 60 mg	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
4. Central nervous system	4.7 Analgesics	4.7.2 Opioid analgesics	Buprenorphine	Patches: '5' patch (releasing 5 micrograms/hour for 7 days), '10' patch (releasing 10 micrograms/hour for 7 days) '15' patch (releasing 10 micrograms/hour for 7 days) '20' patch (releasing 20 micrograms/hour for 7 days) 35 micrograms / hour for 96 hours 52.5 micrograms / hour for 96 hours 70 micrograms / hour for 96 hours Sublingual tablets: 200micrograms 400micrograms	Unspecified	All prescribers	Prescribe by brand. For 7-day patches, Sevodyne® brand is the preferred option as it costs less than other transdermal patches.
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Amitriptyline hydrochloride	tablets 10mg, 25mg, 50mg; oral solution 25mg in 5ml, 50mg in 5ml	Unspecified	All prescribers	Approved for use in migraine prophylaxis For use in line with NW London Headache Primary Care Clinical Pathway (Adults)
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Atocepan	tablets 10mg, 60mg	Unspecified	See comments column	<u>Acute treatment:</u> Approved for the management of acute migraine in line with NICE TA 1172 Suitable for initiation in primary and secondary care. <u>Preventative treatment:</u> Approved for migraine prophylaxis in line with NICE TA 973 Initiate in primary care following specialist clinical review and recommendation only Primary care review at 12 weeks Episodic migraine (<15 headache days/month): Discontinue atocepan if <50% reduction in migraine frequency. Refer to neurology for alternative options Chronic migraine (≥15 headache days/month, including ≥8 migraine days) Discontinue atocepan if <30% reduction in migraine frequency. Refer to neurology for alternative options For use in line with NW London Headache Primary Care Clinical Pathway (Adults) due to be updated
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Candesartan	tablets 2mg, 4mg, 8mg, 16mg	Unspecified	All prescribers	Approved for use in migraine prophylaxis (off-label) For use in line with NW London Headache Primary Care Clinical Pathway (Adults)
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Propranolol	tablets 10mg, 40mg, 80mg, 160mg; MR capsules 80mg, 160mg; oral solution 5mg in 5ml, 10mg in 5ml, 50mg in 5ml	Unspecified	All prescribers	Approved for use in migraine prophylaxis For use in line with NW London Headache Primary Care Clinical Pathway (Adults)
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Rimegepan	oral lyophilisates sugar free 75mg	Unspecified	See comments column	<u>Acute treatment:</u> Approved for the management of acute migraine in line with NICE TA 919 Suitable for initiation in primary and secondary care <u>Preventative treatment:</u> Approved for migraine prophylaxis in line with NICE TA 906 Initiate in primary care following specialist clinical review and recommendation only Primary care review at 12 weeks Episodic migraine (<15 headache days/month): Discontinue rimegepan if <50% reduction in migraine frequency. Refer to neurology for alternative options For use in line with NW London Headache Primary Care Clinical Pathway (Adults)
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Sumatriptan	tablets 50mg, 100mg; 3mg/0.5ml solution for injection pre-filled disposable devices; 6mg/0.5ml solution for injection pre-filled disposable devices; nasal spray 10mg in 0.1ml, 20mg in 0.1ml	First line	All prescribers	For use in line with NW London Headache Primary Care Clinical Pathway (Adults)

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Topiramate	tablets 25mg, 50mg, 100mg; sprinkle capsules 15mg, 25mg, 50mg	Unspecified	All prescribers	Approved for use in migraine prophylaxis For use in line with NW London Headache Primary Care Clinical Pathway (Adults) Drug Safety Update June 2024: Topiramate (Topamax): introduction of new safety measures, including a Pregnancy Prevention Programme
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Naratriptan	tablets 2.5mg	Unspecified	All prescribers	For use in line with NW London Headache Primary Care Clinical Pathway (Adults)
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Frovatriptan	tablets 2.5mg	Unspecified	All prescribers	For use in line with NW London Headache Primary Care Clinical Pathway (Adults)
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Rizatriptan	orodispersible tablets sugar free 5mg, 10mg	Unspecified	All prescribers	For use in line with NW London Headache Primary Care Clinical Pathway (Adults)
4. Central nervous system	4.7 Analgesics	4.7.4 Antimigraine drugs	Zolmitriptan	nasal spray unit dose 5mg/0.1ml	Unspecified	All prescribers	For use in line with NW London Headache Primary Care Clinical Pathway (Adults)
4. Central nervous system	4.7 Analgesics	4.7.4.3 Cluster headaches	Verapamil	tablets 40mg, 80mg,120mg	Unspecified	Specialist INITIATION	Approved for use in cluster headache prophylaxis (off-label use) Specialist to prescribe until dose stabilised and then continued in primary care For use in line with NW London Headache Primary Care Clinical Pathway (Adults)
4. Central nervous system	4.8 Antiepileptics	4.8.1	Brivaracetam	tablets 25mg, 50mg, 100mg oral solution sugar free 50mg/5mL	Second line	Specialist INITIATION	To be used in line with NICE Guidance NG 217 as a second line option for generalised epilepsy, myoclonic epilepsy and focal epilepsy where levetiracetam is not tolerated or behavioural side effects are a concern
4. Central nervous system	4.8 Antiepileptics	4.8.1	Vigabatrin	tablets 500mg; powder 500mg/sachet	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Topiramate	tablets 25mg, 50mg, 100mg, 200mg; sprinkle capsules 15mg, 25mg, 50mg	Unspecified	All prescribers	Drug Safety Update June 2024: Topiramate (Topamax): introduction of new safety measures, including a Pregnancy Prevention Programme
4. Central nervous system	4.8 Antiepileptics	4.8.1	Tiagabine	tablets 5mg, 10mg, 15mg	Unspecified	Specialist INITIATION	Do not start valproate for the first time in people (male or female) younger than 55 years, unless 2 specialists independently agree and document that there is no other effective and tolerated treatment, or there are compelling reasons that the reproductive risks do not apply. Ensure the pregnancy prevention programme is in place if valproate is used in women and girls of childbearing potential. Follow the MHRA safety advice on the use of valproate. NICE Bipolar disorder guidance [CG 185] states: If a woman or girl of childbearing potential is already taking valproate, advise her to gradually stop the drug under medical supervision because of the risk of fetal malformations and adverse neurodevelopmental outcomes after any exposure in pregnancy. MHRA Guidance: Valproate use by women and girls Drug Safety Update September 2024: Valproate use in men: as a precaution, men and their partners should use effective contraception Drug Safety Update January 2024: Valproate (Belvo, Convulex, Depakote, Dyzzantil, Epilim, Epilim Chrono or Chronosphere, Episenta, Epival, and Syonell▼): new safety and educational materials to support regulatory measures in men and women under 55 years of age Drug Safety Update October 2023: Valproate: dispense full packs of valproate-containing medicines Drug Safety Update August 2023: Valproate: re-analysis of study on risks in children of men taking valproate Drug Safety Update February 2020: Valproate (Epilim▼, Depakote▼) pregnancy prevention programme: updated educational materials Drug Safety Update April 2019: Valproate medicines and serious harms in pregnancy: new Annual Risk Acknowledgement Form and clinical guidance from professional bodies to support compliance with the Pregnancy Prevention Programme Drug Safety Update April 2018: Valproate medicines (Epilim▼, Depakote▼): contraindicated in women and girls of childbearing potential unless conditions of Pregnancy Prevention Programme are met
4. Central nervous system	4.8 Antiepileptics	4.8.1	Sodium valproate	ec tablets 200mg, 500mg; MR tablets 200mg, 300mg, 500mg; crushable tablets 100mg; liquid (sugar-free) 200mg in 5ml; solution	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Primidone	tablet 250 mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Pregabalin	capsules 25mg, 50mg, 75mg, 100mg, 150mg, 200mg, 225mg, 300mg; capsules phenytoin sodium 25mg, 50mg, 100mg, 300mg;	Unspecified	All prescribers	For use in line with NICE's guidance
4. Central nervous system	4.8 Antiepileptics	4.8.1	Phenytoin	tablets phenytoin sodium 100mg; chewable tablets phenytoin 50mg; suspension phenytoin 30mg in 5ml	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Phenobarbital	tablets 15mg, 30mg, 60mg; elixir 15mg/5ml	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Oxcarbazepine	tablets 150mg, 300mg, 600mg; oral suspension 300mg in 5ml	Unspecified	Specialist INITIATION	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
4. Central nervous system	4.8 Antiepileptics	4.8.1	Levetiracetam	tablets 250mg, 500mg, 750mg, 1g; oral liquid 500mg in 5ml; solution for infusion	Unspecified	Specialist INITIATION	Solution for infusion can be initiated in primary care for palliative care on specialist recommendation with clear communication on dosing.
4. Central nervous system	4.8 Antiepileptics	4.8.1	Lamotrigine	tablets 25mg, 50mg, 100mg, 200mg; dispersible tablets 5mg, 25mg, 100mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Perampanel	tablets 2mg, 4mg, 6mg, 8mg, 10mg, 12mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Gabapentin	capsules 100mg, 300mg, 400mg; Tablets 600mg	Unspecified	All prescribers	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Ethosuximide	capsules 250 mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Clonazepam	tablet 500micrograms	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Clobazam	tablets 10mg;	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Carbamazepine	tablets 100mg, 200mg, 400mg; MR tablets 200mg, 400mg; tablets chewable 100mg, 200mg; liquid 100mg in 5ml	Unspecified	All prescribers	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Cenobamate	tablets, 12.5mg, 25mg, 50mg, 100mg, 150mg, 200mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance [TA753]. Specialist to prescribe the initial dose titration until optimised, which will usually be after around 3 months.
4. Central nervous system	4.8 Antiepileptics	4.8.1	Lacosamide	tablets 50mg, 100mg, 150mg, 200mg; Syrup, 50mg in 5ml	Unspecified	Specialist INITIATION	
4. Central nervous system	4.8 Antiepileptics	4.8.1	Zonisamide	capsules 25mg, 50mg, 100mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Selegiline	tablets 5mg, 10mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Rotigotine	patches 2mg/24hr, 4mg/24hr, 6mg/24hr, 8mg/24hr	Unspecified	Specialist INITIATION	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Ropinirole	tablets 250microgram, 500micrograms, 1mg, 2mg, 5mg; MR tablets 2mg, 4mg, 8mg; starter pack; follow-on pack	Unspecified	Specialist INITIATION	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Rasagiline mesilate	tablet 1 mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Pramipexole	tablets 88microgram, 180micrograms, 350micrograms, 700micrograms; MR tablets 260micrograms, 520micrograms, 1.05mg, 1.57mg, 2.1mg, 2.62mg, 3.15mg	Unspecified	Specialist INITIATION	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Entacapone	tablets 200mg	Unspecified	Specialist INITIATION	Entacapone-containing preparations (levodopa/carbidopa/entacapone), entacapone, and opicapone may be considered as first-line catechol-O-methyltransferase (COMT) inhibitor adjunctive treatment options for patients with Parkinson's disease experiencing 'end-of-dose' motor fluctuations. The choice of treatment should be made collaboratively between the patient and their Parkinson's disease specialist, taking into account individual patient factors and clinical suitability.
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Co-careldopa	tablets 12.5/50, contain levodopa 50mg and carbidopa 12.5mg (Sinemet 62.5 ®); tablets 10/100, contain levodopa 100mg and carbidopa 10mg; tablets 25/250, contain levodopa 250mg and carbidopa 25mg; tablets 25/100, contain levodopa 100mg and carbidopa 25mg; tablets 25/100, modified-release, contain levodopa 100mg and carbidopa 25mg (Half Sinemet CR ®); tablets 50/200, modified-release, contain levodopa 200mg and carbidopa 50mg (Sinemet CR ®);	Unspecified	All prescribers	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Co-beneldopa	capsules 12.5/50 contain levodopa 50mg and benserazide 12.5mg; 25/100 contain levodopa 100mg and benserazide 25mg; 50/200 contain levodopa 200mg and benserazide 50mg; dispersible tablets 12.5/50, 25/100; MR capsules 25/100 (Madopar CR ®) contain levodopa 100mg and benserazide 25mg	Unspecified	All prescribers	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Staneke •	Entacapone, levodopa and carbidopa tablets (all strengths)	Unspecified	Specialist INITIATION	Entacapone-containing preparations (levodopa/carbidopa/entacapone), entacapone, and opicapone may be considered as first-line catechol-O-methyltransferase (COMT) inhibitor adjunctive treatment options for patients with Parkinson's disease experiencing 'end-of-dose' motor fluctuations. The choice of treatment should be made collaboratively between the patient and their Parkinson's disease specialist, taking into account individual patient factors and clinical suitability.
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Amantadine	capsules 100mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.1	Opicapone	capsules 50mg	Unspecified	Specialist INITIATION	Entacapone-containing preparations (levodopa/carbidopa/entacapone), entacapone, and opicapone may be considered as first-line catechol-O-methyltransferase (COMT) inhibitor adjunctive treatment options for patients with Parkinson's disease experiencing 'end-of-dose' motor fluctuations. The choice of treatment should be made collaboratively between the patient and their Parkinson's disease specialist, taking into account individual patient factors and clinical suitability.
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.2	Trihexyphenidyl	tablets 2mg, 5mg; syrup 5mg in 5ml	Unspecified	All prescribers	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.2	Procyclidine	tablets 5mg; sugar free syrup 2.5mg in 5ml, 5mg in 5ml	Unspecified	All prescribers	
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.3 Drugs used in essential tremor, chorea, tics, and related disorders	Riluzole	tablets 50mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance and a shared care protocol.
4. Central nervous system	4.9 Drugs used in parkinsonism and related disorders	4.9.3 Drugs used in essential tremor, chorea, tics, and related disorders	Tetrabenazine	tablets 25mg	Unspecified	All prescribers	
4. Central nervous system	4.10 Drugs used in substance dependence	4.10.1 Alcohol dependence	Acamprosate calcium	tablet 333 mg	Unspecified	All prescribers	
4. Central nervous system	4.10 Drugs used in substance dependence	4.10.1 Alcohol dependence	Disulfiram	tablets 200mg	Unspecified	All prescribers	
4. Central nervous system	4.10 Drugs used in substance dependence	4.10.1 Alcohol dependence	Nalmefene	tablets 18mg	Unspecified	All prescribers	For use in line with NICE's guidance
4. Central nervous system	4.10 Drugs used in substance dependence	4.10.2 Nicotine dependence	Varenicline	Tablets, f/c, varenicline (as tartrate) 500 micrograms , 1 mg	Unspecified	All prescribers	For use in line with NICE's guidance
4. Central nervous system	4.10 Drugs used in substance dependence	4.10.2 Nicotine dependence	Nicotine	Nicotine 1.5mg lozenges sugar free Nicotine 10mg/16hours transdermal patches Nicotine 14mg/24hours transdermal patches Nicotine 15mg inhalation cartridges with device Nicotine 15mg/16hours transdermal patches Nicotine 1mg/dose oromucosal spray sugar free Nicotine 21mg/24hours transdermal patches Nicotine 25mg/16hours transdermal patches Nicotine 2mg lozenges sugar free Nicotine 2mg medicated chewing gum sugar free Nicotine 2mg sublingual tablets sugar free Nicotine 4mg lozenges sugar free Nicotine 4mg medicated chewing gum sugar free Nicotine 500micrograms/dose nasal spray Nicotine 7mg/24hours transdermal patches Nicotine bitartrate 1mg lozenges sugar free Nicotine bitartrate 2mg lozenges sugar free	Unspecified	All prescribers	
4. Central nervous system	4.10 Drugs used in substance dependence	4.10.2 Nicotine dependence	Cytisinicline (Cytisine)	tablets 1.5mg	Unspecified	All prescribers	It is recommended that patients receive support from a local smoking cessation service throughout the entire treatment period.
4. Central nervous system	4.10 Drugs used in substance dependence	4.10.3 Opioid dependence	Methadone	sugar-free oral solution 1mg in 1ml	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
4. Central nervous system	4.10 Drugs used in substance dependence	4.10.3 Opioid dependence	Buprenorphine	sublingual tablets 400micrograms, 2mg, 8mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
4. Central nervous system	4.10 Drugs used in substance dependence	4.10.3 Opioid dependence	Naltrexone	tablets 50mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
4. Central nervous system	4.11 Dementia	4.11 Dementia	Rivastigmine	capsules 1.5mg, 3mg, 4.5mg, 6mg; oral solution 2mg/ml; patches 4.6mg/24 hours, 9.5mg/24 hours, 13.3mg/24 hours	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
4. Central nervous system	4.11 Dementia	4.11 Dementia	Memantine	tablets 10mg, 20mg; treatment initiation pack; oral solution 5mg/actuation	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
4. Central nervous system	4.11 Dementia	4.11 Dementia	Galantamine	MR capsules 8mg, 16mg, 24mg oral solution 4mg in 1ml	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
4. Central nervous system	4.11 Dementia	4.11 Dementia	Donepezil	tablets 5mg, 10mg	Unspecified	Specialist INITIATION	For use in line with NICE's guidance Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.1 Penicillins	Phenoxyethylpenicillin (penicillin V)	tablets 250mg; sugar-free oral solution 125mg in 5ml, 250mg in 5ml	Unspecified	All prescribers	The following alternative sugar containing presentations may be used if there is an individual clinical need or supply issues with the sugar-free formulary alternatives: Phenoxyethylpenicillin 125mg/5mL oral solution Phenoxyethylpenicillin 250mg/5mL oral solution

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
5. Infections	5.1 Antibacterial drugs	5.1.1 Penicillins	Flucloxacillin	capsules 250mg, 500mg; sugar-free oral solution 125mg in 5ml, 250mg in 5ml	Unspecified	All prescribers	Refer to local anti-bacterial guidelines The following alternative sugar containing presentations may be used if there is an individual clinical need or supply issues with the sugar –free formulary alternatives: Flucloxacillin 125mg/5mL oral solution Flucloxacillin 250mg/5mL oral solution The Pharmacy First PGD states: Flucloxacillin sugar free oral solution may have a poor taste potentially leading to reduced compliance. After discussion with individual/carer/parent/guardian consider sugar containing preparation
5. Infections	5.1 Antibacterial drugs	5.1.1 Penicillins	Co-amoxiclav	Tablets 375mg (amoxicillin 250mg and clavulanic acid 125mg); 625mg (amoxicillin 500mg and clavulanic acid 125mg); suspension 125/31 in 5ml, 250/62 in 5ml, 400/57 in 5ml	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.1 Penicillins	Benzylpenicillin	injection 600mg, 1.2g	Unspecified	All prescribers	For suspected meningococcal disease. Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.1 Penicillins	Amoxicillin	capsules 250mg, 500mg; sugar-free oral suspension 125mg in 5ml; sugar-free oral suspension 250mg in 5ml; sachets 3g	Unspecified	All prescribers	Refer to local anti-bacterial guidelines The following alternative sugar containing presentations may be used if there is an individual clinical need or supply issues with sugar-free formulary alternatives: Amoxicillin 125mg/5mL oral suspension Amoxicillin 250mg/5mL oral suspension Amoxicillin 500mg/5mL SF oral suspension may be used if there is an individual clinical need or supply issues with formulary alternatives
5. Infections	5.1 Antibacterial drugs	5.1.1 Penicillins	Pivmecillinam hydrochloride	tablets 200mg	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.2 Cephalosporins	Cefalexin (cephalexin)	capsules 250mg, 500mg; oral liquid/suspension 125mg in 5ml, 250mg in 5ml	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.2 Cephalosporins	Cefixime	tablets 200mg	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.3 Tetracyclines	Oxytetracycline	tablets 250 mg	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.3 Tetracyclines	Lymecycline	capsules 408 mg	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.3 Tetracyclines	Doxycycline	capsules 50mg, 100mg; dispersible tablets 100mg;	Unspecified	All prescribers	Capsules cost less than dispersible tablets. Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.3 Tetracyclines	Demeclocycline hydrochloride	capsules 150mg	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.3 Tetracyclines	Minocycline	capsules 100mg	Unspecified	Specialist INITIATION	Refer to local anti-bacterial guidelines. Included in the Formulary for respiratory infections on the advice of a specialist, not for acne. Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.5 Macrolides	Erythromycin	e/c tablets 250mg; sugar-free suspension 125mg in 5ml, 250mg in 5ml, 500mg in 5ml	Unspecified	All prescribers	The following alternative sugar containing presentations may be used if there is an individual clinical need or supply issues with the sugar –free formulary alternatives: Erythromycin 125mg/5mL oral suspension Erythromycin 250mg/5mL oral suspension Erythromycin 500mg/5mL oral suspension'
5. Infections	5.1 Antibacterial drugs	5.1.5 Macrolides	Clarithromycin	tablets 250mg, 500mg; suspension 125mg in 5ml, 250mg in 5ml	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.5 Macrolides	Azithromycin	tablets 250mg, 500mg; oral suspension 200mg in 5ml	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.6 Clindamycin	Clindamycin	capsules 150mg	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.7 Some other antibacterials	Vancomycin	capsules 125mg, 250mg	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.7 Some other antibacterials	Fidaxomicin	200mg tablets, 40mg/ml oral suspension sugar free	Unspecified	Specialist INITIATION	For use in line with NICE/ North West London Antimicrobial Prescribing Guidelines for Primary Care. GP initiation permissible following microbiology advice.
5. Infections	5.1 Antibacterial drugs	5.1.8 Sulfonamides and trimethoprim	Co-trimoxazole	tablets 480mg, 960mg; suspension 240mg in 5ml, 480mg in 5ml;	Unspecified	Specialist INITIATION	Restricted use - see BNF. Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.8 Sulfonamides and trimethoprim	Trimethoprim	tablets 100mg, 200mg; suspension 50mg in 5ml	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.9 Antituberculosis drugs	Rifampicin	capsules 150mg, 300mg; combined preparations: Rifater, Rifinah 150/100, Rifinah 300/150	Unspecified	Specialist INITIATION	Red Listed for the treatment of TB: the norm is for TB treatment to be prescribed by the team supervising the treatment
5. Infections	5.1 Antibacterial drugs	5.1.9 Antituberculosis drugs	Pyrazinamide	tablets 500mg	Unspecified	Specialist INITIATION	Red Listed for the treatment of TB: the norm is for TB treatment to be prescribed by the team supervising the treatment
5. Infections	5.1 Antibacterial drugs	5.1.9 Antituberculosis drugs	Isoniazid	tablets 50mg, 100mg	Unspecified	Specialist INITIATION	Red Listed for the treatment of TB: the norm is for TB treatment to be prescribed by the team supervising the treatment
5. Infections	5.1 Antibacterial drugs	5.1.9 Antituberculosis drugs	Ethambutol	tablets 100mg, 400mg	Unspecified	Specialist INITIATION	Red Listed for the treatment of TB: the norm is for TB treatment to be prescribed by the team supervising the treatment
5. Infections	5.1 Antibacterial drugs	5.1.9 Antituberculosis drugs	Rifabutin	capsules 150mg	Unspecified	Specialist INITIATION	Red Listed for the treatment of TB: the norm is for TB treatment to be prescribed by the team supervising the treatment
5. Infections	5.1 Antibacterial drugs	5.1.11 Metronidazole	Metronidazole	tablets 200mg, 400mg; suspension 200mg in 5ml; suppositories 500mg, 1g	Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.12 Quinolones	Ciprofloxacin	tablets 100mg, 250mg, 500mg, 750mg suspension 250mg in 5ml	Unspecified	All prescribers	Refer to local anti-bacterial guidelines Drug Safety Update January 2024: Fluoroquinolone antibiotics: must now only be prescribed when other commonly recommended antibiotics are inappropriate
5. Infections	5.1 Antibacterial drugs	5.1.12 Quinolones	Levofloxacin	tablets 250mg, 500mg	Unspecified	Specialist INITIATION	Refer to local anti-bacterial guidelines Drug Safety Update January 2024: Fluoroquinolone antibiotics: must now only be prescribed when other commonly recommended antibiotics are inappropriate

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
5. Infections	5.1 Antibacterial drugs	5.1.12 Quinolones	Moxifloxacin	tablets 400mg	Unspecified	Specialist INITIATION	Refer to local anti-bacterial guidelines Drug Safety Update January 2024: Fluoroquinolone antibiotics: must now only be prescribed when other commonly recommended antibiotics are inappropriate
5. Infections	5.1 Antibacterial drugs	5.1.12 Quinolones	Ofloxacin	tablets 200mg, 400mg	Unspecified	Specialist INITIATION	Refer to local anti-bacterial guidelines Drug Safety Update January 2024: Fluoroquinolone antibiotics: must now only be prescribed when other commonly recommended antibiotics are inappropriate
5. Infections	5.1 Antibacterial drugs	5.1.13 Urinary-tract infections	Nitrofurantoin	capsules 50mg, 100mg; capsules 100mg; tablets 50mg, 100mg; suspension 25mg in 5ml granules 3g	MR Unspecified	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.13 Urinary-tract infections	Fosfomycin	granules 3g	Second line	All prescribers	Refer to local anti-bacterial guidelines
5. Infections	5.1 Antibacterial drugs	5.1.13 Urinary-tract infections	Methenamine hippurate	tablets 1g	Unspecified	Specialist INITIATION	For use in line with NICE's guidance Not to be continued after 6 months unless reviewed
5. Infections	5.2 Antifungal drugs	5.2	Terbinafine	tablets 250mg	Unspecified	All prescribers	
5. Infections	5.2 Antifungal drugs	5.2	Nystatin	suspension 100000units in 1ml	Unspecified	All prescribers	
5. Infections	5.2 Antifungal drugs	5.2	Itraconazole	capsules 100mg; oral liquid 10mg in 1ml	Unspecified	All prescribers	
5. Infections	5.2 Antifungal drugs	5.2	Fluconazole	capsules 50mg, 150mg, 200mg; suspension 50mg in 5ml, 200mg in 5ml;	Unspecified	All prescribers	
5. Infections	5.3 Antiviral drugs	5.3.2 Herpesvirus infections	Valaciclovir	tablets 250mg, 500mg; tablets 200mg, 400mg, 800mg; suspension 200mg in 5ml, 400mg in 5ml; dispersible tablets 200mg, 400mg, 800mg;	Unspecified	All prescribers	Dispersible tablets cost less than oral suspension.
5. Infections	5.3 Antiviral drugs	5.3.2 Herpesvirus infections	Aciclovir		Unspecified	All prescribers	For Pharmacy First the PGD states: The 800mg tablets should be supplied in the first instance and 400mg or 200mg tablets supplied only if 800mg tablets are not available
5. Infections	5.3 Antiviral drugs	5.3.4 Influenza	Zanamivir	dry powder for inhalation 5mg/blister	Unspecified	All prescribers	For use in line with NICE's guidance
5. Infections	5.3 Antiviral drugs	5.3.4 Influenza	Oseltamivir	capsules 30mg, 45mg, 75mg; suspension 6mg/ml	Unspecified	All prescribers	For use in line with NICE's guidance
5. Infections	5.4 Antiprotozoal drugs	5.4.1	Quinine sulphate	tablets 200mg, 300mg	Unspecified	All prescribers	
5. Infections	5.5 Anthelmintics	5.5.1	Mebendazole	chewable tablets 100mg; suspension, 100mg in 5ml	Unspecified	All prescribers	
5. Infections	5.5 Anthelmintics	5.5.6	Ivermectin	tablets 3mg	Third line	All prescribers	To be used when: Scabies that has not been resolved after using first line topical treatments or if topical treatments are contra-indicated (permethrin and malathion) For hyperkeratotic (crusted) Scabies
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.1 Short-acting insulins	Insulin lispro	Humalog [®] - all formulations 100 units/ml, 200 units/ml	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.1 Short-acting insulins	Insulin lispro (Lyumjev [®])	Lyumjev [®] - 100 units/mL KwikPen solution for injection in pre-filled pen 200 units/mL KwikPen solution for injection in pre-filled pen 100 units/ml Solution for injection vial (10ml) 100 units/mL Solution for injection cartridge (3ml)	Second line	Specialist initiation	Ultra rapid-acting analogue formulation. Please note this formulation is not directly equivalent to other short acting insulins. Insulins must be prescribed by brand name. Second-line fast acting insulin in adults, restricted to use where the first line fast acting insulin does not provide adequate post-prandial plasma glucose (PPG) control. Initiation, dose titration and stabilisation will be undertaken by the Diabetes specialist team with the first 4-6 weeks' supply prescribed. Prescribing can subsequently continued by prescribers in primary care.
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.1 Short-acting insulins	Insulin glulisine	Apidra - all formulations	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.1 Short-acting insulins	Insulin aspart (Trurapi)	All formulations	Unspecified	All prescribers	Prescribe by brand name All new initiations should be the lower cost biosimilar, Trurapi, instead of NovoRapid For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.1 Short-acting insulins	Insulin aspart (Novorapid)	All formulations	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.1 Short-acting insulins	Insulin	Actrapid, Humulin S - all formulations	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.2 Intermediate- and long-acting insulins	Isophane (NPH) insulin	Insulatard, Humulin I - all formulations	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.2 Intermediate- and long-acting insulins	Insulin glargine (Abasaglar)	100units/ml	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.2 Intermediate- and long-acting insulins	Insulin glargine (Lantus)	100units/ml	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.2 Intermediate- and long-acting insulins	Insulin glargine (Semglee [®])	100units/ml	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.2 Intermediate- and long-acting insulins	Insulin detemir	all formulations	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.2 Intermediate- and long-acting insulins	Biphasic isophane insulin	Humulin M3 - all formulations	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.2 Intermediate- and long-acting insulins	Biphasic insulin lispro	Humalog Mix25, Humalog Mix50 - all formulations	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.1.2 Intermediate- and long-acting insulins	Biphasic insulin aspart	NovoMix 30 - all formulations	Unspecified	All prescribers	Prescribe by brand name For use in line with NICE guidance or NW London Diabetes guidelines
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.1 Sulphonylureas	Glimepiride	tablets 1mg, 2mg, 3mg, 4mg	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.1 Sulphonylureas	Gliclazide	tablets 40mg, 80mg	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.2 Biguanides	Metformin	tablets 500mg, 850mg; MR tablets 500mg, 750mg and 1g Metformin 500mg oral powder sachets sugar free	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Linagliptin	tablets 5mg	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Alogliptin	tablets 6.25mg, 12.5mg, 25mg	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Vildagliptin	tablets 50mg; tablets vildagliptin + metformin 850mg (Eucreas 50mg/850mg) or metformin 1g (Eucreas 50mg/1g)	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Sitagliptin	tablets, 25mg, 50mg, 100mg; tablets sitagliptin 50mg + metformin 1g	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Saxagliptin	tablets, 2.5mg, 5mg	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Repaglinide	tablets 500micrograms, 1mg, 2mg	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Pioglitazone	tablets 15mg, 30mg, 45mg	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Canagliflozin	tablets 100mg, 300mg	Unspecified	All prescribers	For use in line with NICE's guidance
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Dapagliflozin	tablets 5mg, 10mg	Unspecified	All prescribers	Generic dapagliflozin is the preferred SGLT-2i for the treatment of type 2 diabetes (with or without chronic kidney disease) and/or heart failure in line with its marketing authorisation. For use in line with NICE TA288, NICE TA390, NICE TA418, NICE TA1075
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Empagliflozin	tablets 10mg, 25mg	Unspecified	All prescribers	For use in line with NICE TA942, NICE TA336 and NICE TA390
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Ertugliflozin	tablets 5mg, 15mg	Unspecified	All prescribers	For use in line with NICE's guidance
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Dulaglutide	injection 0.75mg, 1.5mg, 3mg and 4.5mg	Unspecified	All prescribers	For use in line with NICE type 2 diabetes in adults [NG28] Stop GLP-1 mimetic prescriptions if NICE's criteria for continuing have not been met • For use in line with NICE type 2 diabetes in adults [NG28] • Routine switching to oral semaglutide from injectable GLP1s is not recommended as cardiovascular benefits of oral semaglutide have not been proven • Rybelsus® tablets are being updated to a new formulation with improved bioavailability, leading to the following dose changes:
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Semaglutide	0.25mg/0.19ml solution for injection 1.5ml pre-filled disposable device; 0.5mg/0.37ml solution for injection 1.5ml pre-filled disposable device; 1mg/0.74ml solution for injection 3ml pre-filled disposable device; 1.5mg tablets, 3mg tablets, 4mg tablets, 7mg tablets, 9mg tablets and 14mg tablets	Unspecified	All prescribers	3 mg (original starting dose) → 1.5 mg (new starting dose) 7 mg (original maintenance dose) → 4 mg (new maintenance dose) 14 mg (original maintenance dose) → 9 mg (new maintenance dose) Semaglutide (Wegovy®) solution for injection is on the Red List in line with NICE guidelines for overweight and obesity NG 246 and NICE TA 875 For use in line with NICE type 2 diabetes in adults [NG28] and NICE TA 924 Initiation by a specialist or a suitably trained healthcare professional in primary care.
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Tirzepatide	solution for injection 2.4ml pre-filled disposable devices (all strengths)	Unspecified	See comments column	Each KwikPen device contains four doses, hence a single pen contains 4 weeks supply.
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Tirzepatide	solution for injection 2.4ml pre-filled disposable devices (all strengths)	Unspecified	See comments column	For use in line with NICE TA 1026 for prescribing in overweight and obesity To be prescribed in line with the NHS England cohorts and timelines. Please see link below for further information: https://www.england.nhs.uk/long-read/interim-commissioning-guidance-nice-ta1026-tirzepatide/
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Liraglutide	6mg/ml solution for injection 3ml pre-filled disposable devices	Unspecified	All prescribers	For use in line with NICE type 2 diabetes in adults [NG28] and NICE TA 924
6. Endocrine system	6.1 Drugs used in diabetes	6.1.2.3 Other antidiabetic drugs	Acarbose	tablets 50mg, 100mg	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.4 Treatment of hypoglycaemia	Diazoxide	tablets 50mg	Unspecified	Specialist INITIATION	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.4 Treatment of hypoglycaemia	Glucagon	1mg powder and solvent for solution for injection vials (GlucaGen Hypokit 1mg powder and solvent for solution for injection)	Unspecified	All prescribers	
6. Endocrine system	6.1 Drugs used in diabetes	6.1.6 Diagnostic and monitoring devices for diabetes mellitus	Diagnostic and monitoring devices for diabetes mellitus		Unspecified	All prescribers	Please refer to North West London Preferred Blood Glucose and Ketone Meters, Test Strips and Lancets guidance North West London Preferred Blood Glucose and Ketone Meters, Test Strips and Lancets– November 2025
6. Endocrine system	6.1 Drugs used in diabetes	6.1.6 Glucose interstitial fluid detection sensors	Freestyle Libre 2 Plus		Unspecified	All prescribers	For information on continuous glucose monitoring (CGM) devices, please refer to the individual device entries. For CGM access criteria please refer to: Pan London implementation document for type 2 diabetes A-pan-London-implementation-document-for-continuous-glucose-sensors-for-people-living-with-type-2-diabetes.pdf and NICE Guidance for Type 1 Diabetes (NG17)
6. Endocrine system	6.1 Drugs used in diabetes	6.1.6 Glucose interstitial fluid detection sensors	Dexcom One+		Unspecified	All prescribers	For CGM access criteria please refer to: Pan London implementation document for type 2 diabetes A-pan-London-implementation-document-for-continuous-glucose-sensors-for-people-living-with-type-2-diabetes.pdf and NICE Guidance for Type 1 Diabetes (NG17)
6. Endocrine system	6.1 Drugs used in diabetes	6.1.6 Glucose interstitial fluid detection sensors	GlucoRx Aidex		Unspecified	All prescribers	For CGM access criteria please refer to: Pan London implementation document for type 2 diabetes A-pan-London-implementation-document-for-continuous-glucose-sensors-for-people-living-with-type-2-diabetes.pdf and NICE Guidance for Type 1 Diabetes (NG17)

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
6. Endocrine system	6.1 Drugs used in diabetes	6.1.6 Glucose interstitial fluid detection sensors	CareSens Air		Unspecified	All prescribers	For CGM access criteria please refer to: Pan London implementation document for type 2 diabetes A-pan-London-implementation-document-for-continuous-glucose-sensors-for-people-living-with-type-2-diabetes.pdf and NICE Guidance for Type 1 Diabetes (NG17) Should only be used in line with the following guidance:
6. Endocrine system	6.2 Thyroid and anti-thyroid drugs	6.2.1 Thyroid hormones	Liothyronine sodium	Tablets 5microgram, 10microgram, 20microgram Capsules 5microgram, 10microgram, 20microgram	Unspecified	Specialist INITIATION	NHS England » Items which should not routinely be prescribed in primary care: policy guidance NHS England » Liothyronine – advice for prescribers Liothyronine capsules cost less than tablets
6. Endocrine system	6.2 Thyroid and anti-thyroid drugs	6.2.1 Thyroid hormones	Levothyroxine (thyroxine sodium)	tablets 25micrograms, 50micrograms, 100microgram	Unspecified	All prescribers	Liquid formulations of levothyroxine are expensive. The tablets are small.
6. Endocrine system	6.2 Thyroid and anti-thyroid drugs	6.2.2 Antithyroid drugs	Propylthiouracil	tablets 50mg	Unspecified	All prescribers	
6. Endocrine system	6.2 Thyroid and anti-thyroid drugs	6.2.2 Antithyroid drugs	Carbimazole	tablets 5mg, 20mg	Unspecified	All prescribers	
6. Endocrine system	6.3 Corticosteroids	6.3.1	Fludrocortisone acetate	tablets 100micrograms	Unspecified	Specialist INITIATION	
6. Endocrine system	6.3 Corticosteroids	6.3.2	Prednisolone	tablets 1mg, 5mg; tablets soluble 5mg	Unspecified	All prescribers	Soluble prednisolone tablets cost more than plain tablets.
6. Endocrine system	6.3 Corticosteroids	6.3.2	Hydrocortisone	tablets 10mg, 20mg	Unspecified	All prescribers	
6. Endocrine system	6.3 Corticosteroids	6.3.2	Dexamethasone	tablets 500micrograms, 2mg	Unspecified	All prescribers	
6. Endocrine system	6.3 Corticosteroids	6.3.2	Betamethasone	soluble tablets 500micrograms	Unspecified	All prescribers	
6. Endocrine system	6.3 Corticosteroids	6.3.2	Methylprednisolone	tablets 100mg	Unspecified	Specialist INITIATION	Only for treating acute relapse of multiple sclerosis (0.5 g daily for 5 days) in line with the NICE [CG186] and local guidelines.
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Raloxifene	tablets 60mg	Unspecified	All prescribers	For use in line with NICE's guidance
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Tibolone	tablets 2.5mg	Unspecified	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Premique ®	Premique Low Dose;	Fourth line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Kliefem ®	tablets	First line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Femoston ®	1/10 tablets, 2/10 tablets	Second line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Femoston-conti ®	tablets	Second line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Evorel ®	'25' patch, 50 patch, 75 patch, 100 patch	First line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Evorel Sequi ®	patches	First line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Evorel Conti ®	patches	First line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Estradot ®	'25' patch, 37.5 patch, 50 patch, 75 patch, 100 patch	Second line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Elleste-Duet ®	Elleste-Duet 1mg tablets; Elleste-Duet 2mg tablets	First line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Elleste-Solo ®	Elleste-Solo 1mg tablets; Elleste-Solo 2mg tablets	First line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Premarin ®	tablets 300micrograms, 625micrograms, 1.25mg	Third line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Zumenon ®	1mg tablets, 2mg tablets	Second line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Oestrogel ®	Gel 0.06% (64 dose pump pack)	First line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Sandrena ®	0.5mg gel, 1mg gel	First line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Lenzetto ®	1.53mg/dose transdermal spray	Second line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.1 Oestrogens and HRT	Kliovance ®	tablets	First line	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.2 Progestogens and progesterone receptor modulators	Dydrogesterone	tablets 10mg	Second line	All prescribers	To be used in combination with oral or transdermal oestrogen, as a second line option for menopausal women with a uterus requiring HRT in the following cohorts: - For patients who are unable to tolerate the adverse effects of micronised progesterone. - For patients who have previously been on Femoston ® (estradiol/dydrogesterone) and develop venous thromboembolism risk factors during the treatment period.
6. Endocrine system	6.4 Sex hormones	6.4.1.2 Progestogens and progesterone receptor modulators	Progesterone	8% vaginal gel 1.125g applicators gel 90mg/application pessaries 400mg	Unspecified	All prescribers	Red listed for use associated with IVF.
6. Endocrine system	6.4 Sex hormones	6.4.1.2 Progestogens and progesterone receptor modulators	Norethisterone	tablets 5mg	Unspecified	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.2 Progestogens and progesterone receptor modulators	Medroxyprogesterone acetate	tablets 2.5mg, 5mg, 10mg	Unspecified	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.1.2 Progestogens and progesterone receptor modulators	Dienogest	tablets 2mg	Unspecified	Specialist INITIATION	
6. Endocrine system	6.4 Sex hormones	6.4.1.2 Progestogens and progesterone receptor modulators	Progesterone (micronised)	100mg capsules, 200mg capsules	Unspecified	All prescribers	
6. Endocrine system	6.4 Sex hormones	6.4.2 Male sex hormones and antagonists	Testosterone undecanoate	injection (oily) 1g in 4ml	Unspecified	Specialist INITIATION	
6. Endocrine system	6.4 Sex hormones	6.4.2 Male sex hormones and antagonists	Testosterone	16.2mg/g transdermal gel (20.25mg per actuation) [Testogel] 2% transdermal gel (10mg per actuation) [Tostan] 20mg/g transdermal gel (23mg per actuation) [Testavan] 40.5mg/2.5g transdermal gel unit dose sachets	Unspecified	Specialist INITIATION	
6. Endocrine system	6.4 Sex hormones	6.4.2 Male sex hormones and antagonists	Finasteride	tablets 5mg	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
6. Endocrine system	6.4 Sex hormones	6.4.2 Male sex hormones and antagonists	Cyproterone acetate	tablets 50mg	Unspecified	Specialist INITIATION	
6. Endocrine system	6.5 Hypothalamic and pituitary hormones and anti-oestrogens	6.5.1	Somatropin	all formulations in BNF	Unspecified	Specialist INITIATION	For use in line with NICE's guidance.
6. Endocrine system	6.5 Hypothalamic and pituitary hormones and anti-oestrogens	6.5.1	Clomifene	tablets 50mg	Unspecified	Specialist INITIATION	
6. Endocrine system	6.5 Hypothalamic and pituitary hormones and anti-oestrogens	6.5.2	Desmopressin	tablets 100mcg, 200mcg; nasal spray 10 micrograms per metered spray	Unspecified	All prescribers	
6. Endocrine system	6.6 Drugs affecting bone metabolism	6.6.2	Sodium clodronate	capsules 400mg; tablets 800mg	Unspecified	Specialist INITIATION	
6. Endocrine system	6.6 Drugs affecting bone metabolism	6.6.2	Ibandronic acid	tablets 50mg	Unspecified	Specialist INITIATION	
6. Endocrine system	6.6 Drugs affecting bone metabolism	6.6.2	Alendronic acid	tablets 10mg, 70mg; effervescent tablets 70mg	Unspecified	All prescribers	For use in line with NICE's guidance. Effervescent tablets cost much more than plain tablets.
6. Endocrine system	6.6 Drugs affecting bone metabolism	6.6.2	Risedronate sodium	tablets 5mg, 30mg, 35 mg	Unspecified	All prescribers	For use in line with NICE's guidance
6. Endocrine system	6.6 Drugs affecting bone metabolism	6.6.2	Denosumab	injection 60mg in 1ml	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
6. Endocrine system	6.7 Other endocrine drugs	6.7.1	Cabergoline	tablets 500micrograms	Unspecified	Specialist INITIATION	
6. Endocrine system	6.7 Other endocrine drugs	6.7.1	Bromocriptine	tablets, 1mg, 2.5mg	Unspecified	Specialist INITIATION	
6. Endocrine system	6.7 Other endocrine drugs	6.7.2	Quinagolide	tablets 75 micrograms; starter pack	Unspecified	Specialist INITIATION	
6. Endocrine system	6.7 Other endocrine drugs	6.7.2	Triptorelin	11.25mg powder and solvent for suspension for injection vials	Unspecified	Specialist INITIATION	
6. Endocrine system	6.7 Other endocrine drugs	6.7.2	Triptorelin embonate	22.5mg powder and solvent for suspension for injection vials	Unspecified	Specialist INITIATION	
6. Endocrine system	6.7 Other endocrine drugs	6.7.2	Triptoreline acetate	3.75mg powder and solvent for suspension for injection pre-filled syringes 3mg powder and solvent for suspension for injection vials	Unspecified	Specialist INITIATION	
6. Endocrine system	6.7 Other endocrine drugs	6.7.2	Nafarelin	nasal spray 200mcg/metered spray; see section 8.03.4	Unspecified	Specialist INITIATION	
6. Endocrine system	6.7 Other endocrine drugs	6.7.2	Leuprorelin acetate		Unspecified	Specialist INITIATION	
6. Endocrine system	6.7 Other endocrine drugs	6.7.2	Goserelin	implant 3.6mg syringe (as acetate); implant 10.8mg syringe (as acetate); see section 8.3.4	Unspecified	Specialist INITIATION	
6. Endocrine system	6.7 Other endocrine drugs	6.7.2	Relugolix–estradiol–norethisterone acetate	Ryeqo 40mg/1mg/0.5mg tablets	Unspecified	Specialist INITIATION	Only for use in line with NICE TA 832 and NICE TA 1057 Specialist initiation with a transfer of prescribing to primary care after 3 months Ryeqo® is the preferred option for non-peptide GNRH receptor antagonist
7. Obstetrics, gynaecology, and urinary-tract disorders	7.1 Drugs used in obstetrics	7.1.1	Syntometrine ®	Injection, 1ml contains ergometrine maleate 500 micrograms and oxytocin 5 units.	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.1 Drugs used in obstetrics	7.1.1	Ergometrine	injection 500mcg in 1ml	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.2 Treatment of vaginal and vulval conditions	7.2.1 Preparations for vaginal and vulval changes	Estriol	intravaginal cream 0.01%, 0.1%; pessaries 500micrograms	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.2 Treatment of vaginal and vulval conditions	7.2.1 Preparations for vaginal and vulval changes	Estradiol	vaginal tablets10micrograms, vaginal ring 7.5mcg/24hr (Estring)	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.2 Treatment of vaginal and vulval conditions	7.2.1 Preparations for vaginal and vulval changes	ReplensMD	Non hormonal vaginal moisturiser	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.2 Treatment of vaginal and vulval conditions	7.2.2 Vaginal and vulval infections	Balance Activ Rx ®	gel 5ml applicator tubes	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.2 Treatment of vaginal and vulval conditions	7.2.2 Vaginal and vulval infections	Sylk	Non hormonal vaginal moisturiser	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.2 Treatment of vaginal and vulval conditions	7.2.2 Vaginal and vulval infections	Metronidazole	vaginal gel 0.75%	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.2 Treatment of vaginal and vulval conditions	7.2.2 Vaginal and vulval infections	Econazole nitrate	topical cream 1%; pessaries 150mg	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.2 Treatment of vaginal and vulval conditions	7.2.2 Vaginal and vulval infections	Clotrimazole	pessaries 100mg, 200mg, 500mg; vaginal cream 10%; topical cream 1%, 2%; 500mg + cream (topical) 2%	pessaries Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.2 Treatment of vaginal and vulval conditions	7.2.2 Vaginal and vulval infections	Clindamycin	vaginal cream 2%	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Yasmin ®; Lucette®		Unspecified	All prescribers	Lucette® costs less than Yasmin®.
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Synphase		Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Norimin ®		Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Microgynon 30 ED ®		Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Microgynon 30 ®; Ovranette ®; Rigevidon®		Unspecified	All prescribers	Rigevidon® costs less than Microgynon 30 ® or Ovranette ®
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Mercilon ®; Gedarel ® 20/150		Unspecified	All prescribers	Gedarel ® 20/150 costs less than Mercilon ®

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Marvelon ®; Gedarel ® 30/150		Unspecified	All prescribers	Gedarel ® 30/150 costs less than Marvelon ®
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Logynon ED ®		Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Logynon ®; TriRegol ®		Unspecified	All prescribers	TriRegol ® costs less than Logynon ®
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Femodette®; Millinette ® 20/75		Unspecified	All prescribers	Millinette ® 20/75 costs less than Femodette ®
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Femodene ®; Millinette ® 30/75		Unspecified	All prescribers	Millinette ® 30/75 costs less than Femodene ®
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Clikue ®		Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Lizina ®		Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Brevinor ®		Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.1 Combined hormonal contraceptives	Evra®		Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.2 Progestogen-only contraceptives	Norethisterone	tablets 350micrograms; injection (oil) norethisterone enantate 200mg/ml	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.2 Progestogen-only contraceptives	Medroxyprogesterone acetate	104mg/0.65ml suspension for injection pre-filled disposable devices; 150mg/1ml suspension for injection pre-filled syringes	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.2 Progestogen-only contraceptives	Levonorgestrel	tablets 30micrograms; intra-uterine system 20micrograms/24hours	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.2 Progestogen-only contraceptives	Etonogestrel	implant 68mg	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.2 Progestogen-only contraceptives	Desogestrel	tablets, 75 micrograms	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.2 Progestogen-only contraceptives	Drospirenone	tablets 4mg	Second line	All prescribers	For use as a second line progestogen-only contraceptive after desogestrel, for women without risk factors for hyperkalaemia
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.4 Contraceptive devices	Intra-uterine contraceptive devices	see BNF for range	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.5 Emergency contraception	Levonorgestrel	tablets 1.5mg (Levonelle ® 1500)	First line	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.3 Contraceptives	7.3.5 Emergency contraception	Ulipristal acetate	tablets 30mg	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.1 Drugs for urinary retention	Tamsulosin	MR capsules 400mcg	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.1 Drugs for urinary retention	Indoramin	tablets 20mg	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.1 Drugs for urinary retention	Doxazosin	tablets 1mg, 2mg, 4mg	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.1 Drugs for urinary retention	Alfuzosin hydrochloride	tablets 2.5mg; M/R tablets 10mq	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.2 Drugs for urinary frequency, enuresis, and incontinence	Duloxetine	capsules 20mg, 40mg	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.2 Drugs for urinary frequency, enuresis, and incontinence	Vibegron	tablets 75mg	Unspecified	All prescribers	For use in line with NICE TA 999: Vibegron for treating the symptoms of overactive bladder syndrome in adults. Only recommended if antimuscarinic medicines are not suitable, do not work well enough or have unacceptable side effects
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.2 Drugs for urinary frequency, enuresis, and incontinence	Mirabegron	M/R tablets 25mg, 50mg	Unspecified	All prescribers	For use in line with NICE's guidance
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.2 Drugs for urinary frequency, enuresis, and incontinence	Oxybutynin	tablets 2.5mg, 5mg; elixir 2.5mg in 5ml	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.2 Drugs for urinary frequency, enuresis, and incontinence	Tolterodine	tablets 1mg, 2mg; MR tablets 2mg, 4mg	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.2 Drugs for urinary frequency, enuresis, and incontinence	Darifenacin	tablets 7.5mg, 15mg M/R	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.2 Drugs for urinary frequency, enuresis, and incontinence	Solifenacin	tablets 5mg, 10mg	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.3 Drugs used in urological pain	Potassium citrate	soluble tablets (Effercitrate - 1 tablet equivalent to 5ml potassium citrate); mixture 1.5q in 5ml	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.4 Bladder instillations	Sodium chloride	irrigation 0.9% 100ml	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.4 Bladder instillations	Chlorhexidine	Catheter patency solution 0.02%	Unspecified	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.5 Drugs for erectile dysfunction	Sildenafil	tablets 25mg, 50mg, 100mg	First line	All prescribers	
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.5 Drugs for erectile dysfunction	Vardenafil	tablets 5mg, 10mg, 20mg	Unspecified	All prescribers	See Drug Tariff for circumstances under which this can be prescribed on the NHS.
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.5 Drugs for erectile dysfunction	Tadalafil	tablets 5mg, 10mg, 20mg	Unspecified	See comments column	See Drug Tariff for circumstances under which this can be prescribed on the NHS.

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.5 Drugs for erectile dysfunction	Alprostadil	Injection 5microgram, 10microgram, 20 microgram, 40microgram (Caverject ®); cartridges 10microgram, 20microgram (Caverject ® Dual cartridge); urethral application 125micrograms, 250micrograms, 500micrograms, 1mg (Muse ®), Cream; 300microgram single-use container	Unspecified	Specialist INITIATION	See Drug Tariff for circumstances under which this can be prescribed on the NHS
7. Obstetrics, gynaecology, and urinary-tract disorders	7.4 Drugs for genito-urinary disorders	7.4.5 Drugs for erectile dysfunction	Aviptadil with phentolamine mesilate	Solution for injection 25micrograms/2mg/0.35ml ampoules (Invicorp®)	Unspecified	Specialist INITIATION	See Drug Tariff for circumstances under which this can be prescribed on the NHS
7. Obstetrics, gynaecology, and urinary-tract disorders	7.5 Other drugs for gynaecological conditions	7.5.1 Other drugs for gynaecological conditions	Fezolinetant	tablets 45mg	Unspecified	All prescribers	Approved for treating moderate to severe vasomotor symptoms associated with menopause in accordance with NICE TA 1143 Monitoring requirements: Liver function tests should be performed before initiation of treatment and monitored during therapy in all patients receiving fezolinetant, in line with the recommendations in the Summary of Product Characteristics. Restrictions and precautions: - Fezolinetant is not recommended for use in patients with known liver disease. - Fezolinetant is not recommended for women receiving active oncological treatment for breast cancer or other oestrogen-dependent malignancies. - In women with a history of breast cancer or other oestrogen-dependent malignancies who have completed oncological treatment, use of fezolinetant should be considered on an individual basis following a specialist benefit-risk assessment.
8. Malignant disease and immunosuppression	8.1 Cytotoxic drugs	8.1.3	Mercaptopurine	tablets 50mg	Unspecified	Specialist INITIATION	To be prescribed in line with a shared care protocol
8. Malignant disease and immunosuppression	8.1 Cytotoxic drugs	8.1.5	Hydroxycarbamide	capsules 500mg	Unspecified	Specialist INITIATION	To be prescribed in line with a shared care protocol
8. Malignant disease and immunosuppression	8.2 Drugs affecting the immune response	8.2.1	Mycophenolate mofetil	capsules 250mg; tablets 500mg; suspension 1g in 5ml;	Unspecified	Specialist INITIATION	Red listed for new transplant patients To be prescribed in line with a shared care protocol
8. Malignant disease and immunosuppression	8.2 Drugs affecting the immune response	8.2.1	Azathioprine	tablets 25mg, 50mg	Unspecified	Specialist INITIATION	To be prescribed in line with a shared care protocol for non transplant indications
8. Malignant disease and immunosuppression	8.2 Drugs affecting the immune response	8.2.2	Tacrolimus	capsules 500micrograms, 1mg, 3mg, 5mg (prescribe by brand name)	Unspecified	Specialist INITIATION	Red listed for new transplant patients
8. Malignant disease and immunosuppression	8.2 Drugs affecting the immune response	8.2.2	Ciclosporin	capsules 10mg, 25mg, 50mg, 100mg; oral solution 100mg/ml	Unspecified	Specialist INITIATION	To be prescribed in line with a shared care protocol for non transplant indications
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.2 Progestogens	Megestrol acetate	tablets 160mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.2 Progestogens	Medroxyprogesterone acetate	tablets 100mg, 200mg, 400mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.2 Progestogens	Norethisterone	tablets 5mg	Unspecified	All prescribers	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4.3 Somastatin analogues	Octreotide	Octreotide 100micrograms/1ml solution for injection ampoules Octreotide 100micrograms/1ml solution for injection pre-filled syringes Octreotide 100micrograms/1ml solution for injection vials Octreotide 1mg/5ml solution for injection vials Octreotide 500micrograms/1ml solution for injection ampoules Octreotide 500micrograms/1ml solution for injection pre-filled syringes Octreotide 500micrograms/1ml solution for injection vials Octreotide 50micrograms/1ml solution for injection ampoules	Unspecified	Specialist INITIATION	GPs should only be asked to prescribe this if a local shared care protocol has been agreed OR in palliative care
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Triptorelin	injection 3mg, 3.75mg, 11.25mg, 22.5mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Tamoxifen	tablets 10mg, 20mg, 40mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Leuprorelin	injection 3.75, 11.25mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Letrozole	tablets 2.5mg	Unspecified	Specialist INITIATION	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Goserelin	implant 3.6mg, 10.8mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Flutamide	tablets 250mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Exemestane	tablets 25mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Cyproterone acetate	tablets 50mg, 100mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Bicalutamide	tablets 50mg, 150mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Anastrozole	tablets 1mg	Unspecified	Specialist INITIATION	
8. Malignant disease and immunosuppression	8.3 Sex hormones & hormone antagonists in malignant disease	8.3.4 Hormone antagonists	Relugolix	tablets 120mg	Unspecified	Specialist INITIATION	To be used in line with NICE TA995. Minimum 3 months' stabilisation by specialist before transfer of prescribing to primary care. Specialist to provide complete counselling (which includes contraception and information on managing missed doses)
9. Nutrition and blood	9.1 Anaemias and other blood disorders	9.1.1 Iron-deficiency anaemias	Ferrous fumarate	140mg/5ml oral solution sugar free 210mg tablets 305mg capsules 322mg tablets	First line	All prescribers	
9. Nutrition and blood	9.1 Anaemias and other blood disorders	9.1.1 Iron-deficiency anaemias	Sodium feredetate (sodium iron edetate)	Elixir 190mg in 5ml (27.5mg iron in 5ml).	Unspecified	All prescribers	
9. Nutrition and blood	9.1 Anaemias and other blood disorders	9.1.1 Iron-deficiency anaemias	Ferrous sulphate	tablets 200mg (65mg iron)	Unspecified	All prescribers	
9. Nutrition and blood	9.1 Anaemias and other blood disorders	9.1.1 Iron-deficiency anaemias	Ferrous gluconate	tablets 300mg (35mg iron)	Unspecified	All prescribers	
9. Nutrition and blood	9.1 Anaemias and other blood disorders	9.1.2 Drugs used in megaloblastic anaemias	Hydroxocobalamin	injection 1mg in 1ml	Unspecified	All prescribers	
9. Nutrition and blood	9.1 Anaemias and other blood disorders	9.1.2 Drugs used in megaloblastic anaemias	Folic acid	tablets 400micrograms, 5mg; sugar-free syrup 2.5mg in 5ml	Unspecified	All prescribers	
9. Nutrition and blood	9.1 Anaemias and other blood disorders	9.1.2 Drugs used in megaloblastic anaemias	Cyanocobalamin	tablets 50 micrograms	Unspecified	All prescribers	
9. Nutrition and blood	9.2 Fluids and electrolytes	9.2.1	Sodium chloride	MR tablets 600mg contain approximately 10mmol each of sodium and chloride ions	Unspecified	Specialist INITIATION	
9. Nutrition and blood	9.2 Fluids and electrolytes	9.2.1	Sodium bicarbonate	capsules 500mg	Unspecified	Specialist INITIATION	
9. Nutrition and blood	9.2 Fluids and electrolytes	9.2.1	Potassium; Sando-K ®	Tablets effervescent, potassium bicarbonate and chloride, containing 12mmol potassium and 8mmol chloride per tablet.	Unspecified	All prescribers	
9. Nutrition and blood	9.2 Fluids and electrolytes	9.2.1	Patiromer calcium	8.4g and 16.8g oral powder sachets	Unspecified	Specialist INITIATION	For use in line with NICE's guidance [TA623]. Specialist to prescribe the maintenance treatment until optimised, which will usually be after around 3 months. Once treatment is optimised, specialist to send transfer of care documentation to patient's GP practice detailing the diagnosis, current and ongoing dose, any relevant test results and when the next monitoring is required, as well as specialist's contact information.
9. Nutrition and blood	9.2 Fluids and electrolytes	9.2.1	Sodium zirconium cyclosilicate	5g and 10g oral powder sachets	Unspecified	Specialist INITIATION	For use in line with NICE's guidance [TA599]. Specialist to prescribe the maintenance treatment until optimised, which will usually be after around 3 months. Once treatment is optimised, specialist to send transfer of care documentation to patient's GP practice detailing the diagnosis, current and ongoing dose, any relevant test results and when the next monitoring is required, as well as specialist's contact information.
9. Nutrition and blood	9.2 Fluids and electrolytes	9.2.1	Oral rehydration salts	Dioralyte ® oral powder sachets	Unspecified	All prescribers	
9. Nutrition and blood	9.5 Minerals	9.5.1	Calcium salts	tablets, film-coated, calcium carbonate 1.25g, providing 500mg or 12.5mmol calcium per tablet (Calcium-500); tablets, chewable, calcium carbonate 1.25g, providing 500mg or 12.5mmol calcium per tablet (Calcichew ®); tablets, chewable, calcium carbonate 2.5g, providing 1g or 25mmol calcium per tablet (Calcichew Forte ®); tablets, effervescent, calcium salts providing 1g or 25mmol calcium per tablet (Calvive 1000 ®)	Unspecified	All prescribers	
9. Nutrition and blood	9.5 Minerals	9.5.1	Magnesium aspartate	10mmol magnesium oral powder sachets (Magnaspartate ®)	Unspecified	All prescribers	
9. Nutrition and blood	9.5 Minerals	9.5.1	Cinacalcet	tablets 30mg, 60mg, 90mg	Unspecified	Specialist INITIATION	GPs should not prescribe cinacalcet for dialysis patients. For use in line with NICE's guidance.
9. Nutrition and blood	9.5 Minerals	9.5.2	Phosphate-Sandoz ®	effervescent tablets containing 16.1mmol phosphate, 20.4mmol sodium and 3.1mmol potassium per tablet (Phosphate Sandoz ®).	Unspecified	Specialist INITIATION	
9. Nutrition and blood	9.5 Minerals	9.5.4	Zinc sulphate	effervescent tablets 125mg (45mg zinc)	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.1 Vitamin A	Vitamins A and D	capsules vitamin A 4000 units, vitamin D 400 units	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
9. Nutrition and blood	9.6 Vitamins	9.6.2 Vitamin B group	Vitamin B compound strong	tablets containing nicotinamide 20mg, pyridoxine hydrochloride 2mg, riboflavin 2mg and thiamine hydrochloride 5mg.	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.2 Vitamin B group	Thiamine (Vitamin B1)	tablets 50mg, 100mg	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.2 Vitamin B group	Pyridoxine hydrochloride (Vitamin B6)	tablets 10mg, 20mg, 50mg	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.4 Vitamin D	Ergocalciferol (calciferol, vitamin D2)	injection 300,000 units in 1ml	Unspecified	All prescribers	Refer to local guidance on vitamin D products.
9. Nutrition and blood	9.6 Vitamins	9.6.4 Vitamin D	Colecalciferol (Cholecalciferol, vitamin D3)	Capsules: 800 units, 3,200 units, 10,000 units, 20,000 units, 50,000 units Tablets: 1000 units, 25,000 units, Colecalciferol 25,000units/1ml oral solution unit dose ampoules sugar free Colecalciferol 50,000units/1ml oral solution unit dose ampoules sugar free	Unspecified	All prescribers	Also refer to local guidance on vitamin D products.
9. Nutrition and blood	9.6 Vitamins	9.6.4 Vitamin D	Calcitriol (1,25-dihydroxycolecalciferol)	Capsules, 250 nanograms, 500 nanograms	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.4 Vitamin D	Calceos ®	chewable tablets, contain calcium carbonate 1.25 g (calcium 500 mg/12.5 mmol), colecalciferol 10 micrograms (400 units), 1000mg/1,000unit chewable tablets (calcium carbonate 2.5 gram and colecalciferol 1000 iu)	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.4 Vitamin D	Calci-D ®	capsules 250 nanograms, 500 nanograms, 1mcg; oral drops 2mca/ml	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.4 Vitamin D	Alfacalcidol (1-alpha-hydroxycolecalciferol)	caplets, calcium carbonate 750 mg (calcium 300 mg/7.5 mmol), colecalciferol 5 micrograms (200 units),	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.4 Vitamin D	Adcal-D3 ®	tablets, calcium carbonate 1.5g (calcium 600mg/15mmol), colecalciferol 10 micrograms (400 units)	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.5 Vitamin E	Alpha tocopheryl acetate	suspension 500mg in 5ml	Unspecified	Specialist INITIATION	
9. Nutrition and blood	9.6 Vitamins	9.6.6 Vitamin K	Phytomenadione	injection 10mg in 1ml, 2mg in 0.2ml	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.6 Vitamin K	Menadiol sodium phosphate	tablets 10mg	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.7 Multivitamin preparations	Vitamins	capsules, ascorbic acid 15 mg, nicotinamide 7.5 mg, riboflavin 500micrograms, thiamine hydrochloride 1mg, vitamin A 2500 units, vitamin D 300 units	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.7 Multivitamin preparations	Ketovite	tablets; liquid	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.7 Multivitamin preparations	Forceval ®	capsules, soluble tablets	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.7 Multivitamin preparations	Dalivit ®	oral drops	Unspecified	All prescribers	
9. Nutrition and blood	9.6 Vitamins	9.6.7 Multivitamin preparations	Abidec ®	oral drops	Unspecified	All prescribers	Includes arachis (peanut) oil
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.1 Non-steroidal anti-inflammatory drugs	Ibuprofen	tablets 200mg, 400mg; MR tablets 800mg; sugar free syrup 100mg in 5ml	First line	All prescribers	Of NSAIDs, ibuprofen is associated with the lowest risk of GI adverse events. MR tablets cost more than conventional tablets.
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.1 Non-steroidal anti-inflammatory drugs	Naproxen	tablets 250mg, 500mg	Second line	All prescribers	
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.1 Non-steroidal anti-inflammatory drugs	Celecoxib	capsules 100mg, 200mg	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.2 Corticosteroids	Methylprednisolone acetate + lidocaine	injection methylprednisolone acetate 40mg + lidocaine 10mg in 1ml, injection methylprednisolone acetate 80mg + lidocaine 20mg in 2ml	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.2 Corticosteroids	Methylprednisolone acetate	injection 40mg in 1ml, 80mg in 2ml	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.2 Corticosteroids	Dexamethasone	injection 3.8mg in 1ml	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.3 Drugs that suppress the rheumatic disease process	Penicillamine	tablets 125mg, 250mg	Unspecified	Specialist INITIATION	
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.3 Drugs that suppress the rheumatic disease process	Methotrexate	tablets 2.5mg; pre-filled disposable injection for subcutaneous use 7.5mg, 10mg, 12.5mg, 15mg, 17.5mg, 20mg, 22.5mg, 25mg, 27.5mg, 30mg	Unspecified	Specialist INITIATION	Ensure NPSA guidance is followed. Cancer indications are Red listed. For use in line with a shared care protocol.
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.3 Drugs that suppress the rheumatic disease process	Leftunomide	tablets 10mg and 20mg	Unspecified	Specialist INITIATION	For use in line with a shared care protocol.
10. Musculoskeletal and joint diseases	10.1 Drugs used in rheumatic disease and gout	10.1.3 Drugs that suppress the rheumatic disease process	Hydroxychloroquine sulphate	tablets 200mg	Unspecified	Specialist INITIATION	For use in line with a shared care protocol.
10. Musculoskeletal and joint diseases	10.1.4 Gout and cytotoxic-induced hyperuricaemia	10.1.4 Gout and cytotoxic-induced hyperuricaemia	Febuxostat	tablets 80mg, 120mg	Unspecified	All prescribers	For use in line with NICE'S guidance
10. Musculoskeletal and joint diseases	10.1.4 Gout and cytotoxic-induced hyperuricaemia	10.1.4 Gout and cytotoxic-induced hyperuricaemia	Colchicine	tablets 500micrograms	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.1.4 Gout and cytotoxic-induced hyperuricaemia	10.1.4 Gout and cytotoxic-induced hyperuricaemia	Allopurinol	tablets 100mg, 300mg	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.2 Drugs used in neuromuscular disorders	10.2.1 Drugs that enhance neuromuscular transmission	Pyridostigmine bromide	tablets 60mg	Unspecified	Specialist INITIATION	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
10. Musculoskeletal and joint diseases	10.2 Drugs used in neuromuscular disorders	10.2.2 Skeletal muscle relaxants	Dantrolene	capsules 25mg, 100mg	Unspecified	Specialist INITIATION	
10. Musculoskeletal and joint diseases	10.2 Drugs used in neuromuscular disorders	10.2.2 Skeletal muscle relaxants	Baclofen	tablets 10mg; oral solution 5mg in 5ml	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.2 Drugs used in neuromuscular disorders	10.2.2 Skeletal muscle relaxants	Tizanidine	tablets 2mg,4mg	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.3 Drugs for the relief of soft-tissue inflammation and topical pain relief	10.3.2	Diclofenac	gel 1.16%	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.3 Drugs for the relief of soft-tissue inflammation and topical pain relief	10.3.2	Ibuprofen	gel 5%	Unspecified	All prescribers	
10. Musculoskeletal and joint diseases	10.3 Drugs for the relief of soft-tissue inflammation and topical pain relief	10.3.2	Capsaicin	cream 0.025%, 0.075%	Unspecified	All prescribers	
11. Eye	11.3 Anti-infective eye preparations	11.3.1 Antibacterials	Chloramphenicol	eye drops 0.5%; eye ointment 1%; Single Use Drops Minims® 0.5% 20x0.5ml - see comments	First line	All prescribers	In primary care, reserve single use drops (Minims®) for when a preservative free preparation is required.
11. Eye	11.3 Anti-infective eye preparations	11.3.1 Antibacterials	Fusidic acid	MR eye drops 1%	Second line	All prescribers	
11. Eye	11.3 Anti-infective eye preparations	11.3.1 Antibacterials	Ofloxacin	eye drops 0.3%	Second line	Specialist INITIATION	
11. Eye	11.3 Anti-infective eye preparations	11.3.1 Antibacterials	Levofloxacin	Preservative free drops Single use units 0.5% 30x 0.5ml only on formulary- see comments	Third line	Specialist INITIATION	Reserved as third line for occasions when a preservative free quinolone preparation is required.
11. Eye	11.3 Anti-infective eye preparations	11.3.1 Antibacterials	Gentamicin	drops 0.3%	Unspecified	Specialist INITIATION	
11. Eye	11.3 Anti-infective eye preparations	11.3.1 Antibacterials	Moxifloxacin	drops 0.5% (as hydrochloride)	Unspecified	Specialist INITIATION	
11. Eye	11.3 Anti-infective eye preparations	11.3.3 Antivirals	Aciclovir	eye ointment 3%	Unspecified	Specialist INITIATION	
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.1 Corticosteroids	Betamethasone	drops (for ear, eye or nose) 0.1%; eye ointment 0.1%;	Unspecified	Specialist INITIATION	Licensed preservative free product not available
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.1 Corticosteroids	Dexamethasone	eye drops 0.1%; Single Use Drops Minims® 0.1% 20x0.5ml	Unspecified	Specialist INITIATION	In primary care, reserve single use drops (Minims®) for when a preservative free preparation is required.
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.1 Corticosteroids	Fluorometholone	eye drops 0.1%	Unspecified	Specialist INITIATION	Licensed preservative free product not available
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.1 Corticosteroids	Hydrocortisone sodium phosphate	single-use eye drops 3.35 mg/ml	Unspecified	Specialist INITIATION	
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.1 Corticosteroids	Loteprednol	eye drops 0.5%	Unspecified	Specialist INITIATION	Licensed preservative free product not available
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.1 Corticosteroids	Prednisolone	drops (for ear or eye) prednisolone sodium phosphate 0.5%; Single Use Drops Minims® 0.5% 20x0.5ml eye drops prednisolone acetate 1%	Unspecified	Specialist INITIATION	In primary care, reserve single use drops (Minims®) for when a preservative free preparation is required.
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.1 Corticosteroids - with antibacterial	Dexamethasone with neomycin and polymyxin B sulphate	drops dexamethasone 0.1%, neomycin 0.35%, polymyxin B sulphate 6000 units/ml (Maxitrol®)	Unspecified	Specialist INITIATION	
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.1 Corticosteroids - with antibacterial	Dexamethasone with tobramycin	drops dexamethasone 0.1%, tobramycin 0.3%	Unspecified	Specialist INITIATION	
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.2 Other anti-inflammatory preparations	Sodium cromoglicate	eye drops 2% 13.5ml, 2% preservative free single use drops	First line	All prescribers	13.5ml bottles cost less than 10ml bottles in primary care.
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.2 Other anti-inflammatory preparations	Olopatadine	eye drops 1mg in 1ml	Second line	All prescribers	
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.2 Other anti-inflammatory preparations	Ketotifen	250micrograms/ml eye drops 0.4ml unit dose preservative free	Second line	All prescribers	When a preservative free product is required
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.2 Other anti-inflammatory preparations	Lodoxamide	eye drops 0.1%	Third line	All prescribers	
11. Eye	11.4 Corticosteroids & other anti-inflammatory preparations	11.4.2 Other anti-inflammatory preparations	Ciclosporin	eye drops 0.1% (Ikervis®) 0.3ml unit dose, eye drops 0.09% (Cequa®) 0.25mL unit dose	Unspecified	Specialist INITIATION	For use in line with NICE's guidance. Cequa® is reserved for the treatment of severe dry eye disease in adults who do not tolerate Ikervis® (1mg/mL) or have complex eye disease requiring more than once daily administration of Ikervis®
11. Eye	11.5 Mydriatics and cycloplegia	11.5 Mydriatics and cyclopegics	Atropine sulphate	Single use drops (Minims®) 1%	Unspecified	Specialist INITIATION	
11. Eye	11.5 Mydriatics and cycloplegia	11.5 Mydriatics and cyclopegics	Cyclopentolate	eye drops 0.5%, 1% Single use drops (Minims®) 0.5%, 1%.	Unspecified	Specialist INITIATION	
11. Eye	11.5 Mydriatics and cycloplegia	11.5 Mydriatics and cyclopegics	Phenylephrine	Single use drops (Minims®) 2.5%, 10%	Unspecified	Specialist INITIATION	
11. Eye	11.5 Mydriatics and cycloplegia	11.5 Mydriatics and cyclopegics	Tropicamide	Single use drops (Minims®) eye drops, 0.5%, 1%.	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma- beta blockers	Timolol	eye drops 0.25%, 0.5%; unit dose eye drops 0.25% - preservative free; eye gel 0.25%, 0.5%; unit dose eye gel 0.1% - preservative free	First line	Specialist INITIATION	Reserve eye gel preparation (once daily preparation) where compliance issues .
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma- beta blockers	Betaxolol	eye drops 0.5%	Second line	Specialist INITIATION	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-prostaglandin analogues and prostamides	Latanoprost	eye drops 50micrograms/1ml, 50micrograms/1ml, preservative free single use drops	First line	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-prostaglandin analogues and prostamides	Bimatoprost	eye drops 100micrograms/ml, 300micrograms/ml preservative free single use drops	Second line	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-prostaglandin analogues and prostamides	Travoprost	eye drops 40micrograms/ml; 40micrograms/ml eye drops 0.1ml unit dose preservative free	Third line	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-prostaglandin analogues and prostamides	Tafuprost	Unit dose eye drops 15 micrograms/ml	Third line	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-combination preparation	Latanoprost and timolol	eye drops, latanoprost 50 micrograms, timolol 5mg/ml.	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-combination preparation	Bimatoprost and Timolol	eye drops, bimatoprost 300micrograms/ml, timolol 5mg/ml preservative free single use drops	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-combination preparation	Travoprost and Timolol	eye drops, travoprost 40micrograms, timolol 5mg/ml	Unspecified	Specialist INITIATION	
	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-combination preparation	Latanoprost and Netarsudil	eye drops latanoprost 50micrograms/ml, netarsudil 200micrograms/ml	Unspecified	Specialist INITIATION	For use in line with NICE TA1009
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-sympathomimetics	Brimonidine	eye drops 0.2%	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-combination preparation	Combigan ®	eye drops, brimonidine tartrate 0.2%, timolol 0.5%	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-carbonic anhydrase inhibitors and systemic drugs	Dorzolamide	eye drops 2%	First line	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-carbonic anhydrase inhibitors and systemic drugs	Brinzolamide	eye drops 10mg/ml	Second line	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-combination preparation	Cosopt ®	eye drops, dorzolamide 2%, timolol 0.5%	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-combination preparation	Taptiqom ®	eye drops, tafuprost 15micrograms/ml, timolol 5mg/ml preservative free single use drops	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-carbonic anhydrase inhibitors and systemic drugs	Azarga ®	eye drops, brinzolamide 10mg/ml, timolol 5mg/ml	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-carbonic anhydrase inhibitors and systemic drugs	Simbrinza ®	eye drops, 10mg/ml brinzolamide, 2mg/ml brimonidine tartrate	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-carbonic anhydrase inhibitors and systemic drugs	Acetazolamide	tablets 250mg	Unspecified	Specialist INITIATION	
11. Eye	11.6 Treatment of glaucoma	11.6 Treatment of glaucoma-miotics	Pilocarpine	eye drops 1%, 2%, 4% Unit dose Minims ® eye drops pilocarpine nitrate 2%	Unspecified	Specialist INITIATION	
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Hypromellose	eye drops 0.3%; eye drops 0.5%; Preservative free unit dose eye drops 0.3% (e.g. Lumecare, Hydromoor)	First line	All prescribers	Preservative free drops cost more than preservative-containing drops. Reserve the former for when it is important to avoid preservative or reduce preservative load.
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Acetylcysteine	eye drops 5% with hypromellose 0.35%	Second line	All prescribers	
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Polyvinyl alcohol	eye drops 1.4% Liquifilm Tears® 15ml Sno Tears® 10ml	Second line	All prescribers	
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Yellow Soft Paraffin	Xaillin Night eye ointment preservative free	Second line	All prescribers	
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Camellose	0.5% eye drops 0.5% eye drops preservative free 1% eye drops preservative free	Third line	All prescribers	Unit dose preparation is a licensed preservative free preparation. Reserve for when it is important to avoid preservative or reduce preservative load.
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Hydroxymethylcellulose	Single Use Drops Minims® Artificial Tears 20x0.5ml	Third line	All prescribers	Licensed preservative free preparation. Reserve for when it is important to avoid preservative or reduce preservative load.
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Sodium chloride	Single use Minims ® eye drops 0.9%	Third line	All prescribers	
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Carbomer 980	eye drops 0.2% (GelTears®, Viscotears®)	Second line	All prescribers	Viscotears® brand preserved with cetrimide not benzalkonium chloride so may be considered where benzalkonium intolerance or toxicity is suspected
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Sodium hyaluronate	0.1% eye drops preservative free 0.2% eye drops preservative free	Third line	All prescribers	
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.1 Tear deficiency, ocular lubricants, and astringents	Optive® Fusion	eye drops, sodium hyaluronate 0.1%, carmellose sodium 0.5%, glycerol 0.9%	Unspecified	All prescribers	
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.2 Ocular peri-operative preparations and photodynamic treatment	Fluorescein sodium	Single use Minims ® eye drops 1%, 2%	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.2 Ocular peri-operative preparations and photodynamic treatment	Ketorolac	eye drops 0.5%	Unspecified	Specialist INITIATION	
11. Eye	11.8 Miscellaneous ophthalmic preparations	11.8.2 Ocular peri-operative preparations and photodynamic treatment	Sodium chloride- hypertonic	eye drops 5%; eye ointment 5%	Unspecified	Specialist INITIATION	Used for corneal oedema.
12. Ear, nose, and oropharynx	12.1 Drugs acting on the ear	12.1.1 Otitis externa	Sofradex ®	drops (for ear or eye), dexamethasone 0.05%, framycetin sulphate 0.5%, gramicidin 0.005%.	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.1 Drugs acting on the ear	12.1.1 Otitis externa	Dexamethasone 0.1%, neomycin sulphate 3250 units/ml, glacial acetic acid 2%	ear spray, dexamethasone 0.1%, neomycin sulphate 3250 units/ml, glacial acetic acid 2%.	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.1 Drugs acting on the ear	12.1.1 Otitis externa	Flumetasone 0.02%/ Cloquinol 1%	Ear Drops solution (Branded Locorten-Vioform ® now discontinued)	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.1 Drugs acting on the ear	12.1.1 Otitis externa	Clotrimazole	solution 1%	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.1 Drugs acting on the ear	12.1.1 Otitis externa	Bethnesol-N ®	drops (for ear, eye or nose), betamethasone sodium phosphate 0.1%, neomycin sulphate 0.5%	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.1 Drugs acting on the ear	12.1.1 Otitis externa	Betamethasone sodium phosphate	drops (for ear, eye or nose) 0.1%.	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.1 Drugs acting on the ear	12.1.1 Otitis media	Phenazone/ Lidocaine	Phenazone 40mg/g / Lidocaine 10mg/g ear drops	Unspecified	All prescribers	For use in line with NICE guideline Otitis media (acute): antimicrobial prescribing (NG91) or the Pharmacy First relevant clinical pathway
12. Ear, nose, and oropharynx	12.1 Drugs acting on the ear	12.1.3 Removal of ear wax	Sodium bicarbonate	ear drops 5%	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.1 Drugs acting on the ear	12.1.3 Removal of ear wax	Olive oil	ear drops	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.1 Drugs used in nasal allergy	Beclometasone dipropionate	aqueous nasal spray 50micrograms per metered spray, nose drops	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.1 Drugs used in nasal allergy	Budesonide	aqueous nasal spray 64micrograms/metered spray	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.1 Drugs used in nasal allergy	Fluticasone furoate	nasal spray 27.5 micrograms per metered spray	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.1 Drugs used in nasal allergy	Fluticasone propionate	nasal spray 50 microgram - metered spray	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.1 Drugs used in nasal allergy	Azelastine	nasal spray 140micrograms/metered spray	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.1 Drugs used in nasal allergy	Mometasone furoate	nasal spray 50 micrograms - metered spray	Unspecified	All prescribers	First choice corticosteroid nasal spray.
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.2 Topical nasal decongestants	Sodium chloride	nasal drops, 0.9%.	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.2 Topical nasal decongestants	Ipratropium bromide	nasal spray 0.03% (21micrograms per metered spray)	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.3 Nasal preparations for infection	Naseptin ®	nasal cream, chlorhexidine hydrochloride 0.1%, neomycin sulphate 0.5%	Unspecified	All prescribers	Use in line with Infection Control policies
12. Ear, nose, and oropharynx	12.2 Drugs acting on the nose	12.2.3 Nasal preparations for infection	Mupirocin	nasal ointment 2%.	Unspecified	All prescribers	Use in line with Infection Control policies
12. Ear, nose, and oropharynx	12.3 Drugs acting on the oropharynx	12.3.1 Drugs for oral ulceration and inflammation	Benzydamine hydrochloride	oral rinse 0.15%; spray 0.15%	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.3 Drugs acting on the oropharynx	12.3.2 Oropharyngeal anti-infective drugs	Nystatin	oral suspension 100,000units in 1ml	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.3 Drugs acting on the oropharynx	12.3.2 Oropharyngeal anti-infective drugs	Miconazole	20mg/g oromucosal gel sugar free	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.3 Drugs acting on the oropharynx	12.3.4 Mouthwashes, gargles, and dentifrices	Chlorhexidine gluconate	mouthwash 0.2%; dental gel 1%	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.3 Drugs acting on the oropharynx	12.3.5 Treatment of dry mouth	Salivix ®	pastilles	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.3 Drugs acting on the oropharynx	12.3.5 Treatment of dry mouth	Pilocarpine	tablets 5mg	Unspecified	All prescribers	
12. Ear, nose, and oropharynx	12.3 Drugs acting on the oropharynx	12.3.5 Treatment of dry mouth	Artificial Saliva - Biotene Oralbalance ®	saliva replacement gel	Unspecified	All prescribers	
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Emulsifying Ointment BP	ointment	Unspecified	Specialist INITIATION	Primary care initiation is permissible following specialist advice For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Cetomacrogol Formula A	cream	Unspecified	Specialist INITIATION	Primary care initiation is permissible following specialist advice For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Zerocream®	cream	Second line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children.
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	ZeroAQS®	emollient cream	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children.
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Zeroderm®	ointment	Second line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children.

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Aquamax®	cream	Unspecified	Specialist INITIATION	Primary care initiation is permissible following specialist advice For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Epimax Original®	cream	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Epimax®	ointment	First line	All prescribers	Drug Safety Update July 2024: Epimax Ointment and Epimax Paraffin-Free Ointment: reports of ocular surface toxicity and ocular chemical injury For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
		13.2.1 Emollients	Epimax® paraffin - free	paraffin-free ointment	Unspecified	All prescribers	Drug Safety Update July 2024: Epimax Ointment and Epimax Paraffin-Free Ointment: reports of ocular surface toxicity and ocular chemical injury For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Epimax® Oatmeal	cream	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Cetaben®	lotion	Unspecified	Specialist INITIATION	Primary care initiation is permissible following specialist advice For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Epimax ExCetra ®	cream	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Unguentum M®	cream 100g	Unspecified	Specialist INITIATION	Primary care initiation is permissible following specialist advice For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	AproDerm®	emollient cream, gel	Second line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	AproDerm®	ointment	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Epimax Isomol ®	gel	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Fifty:50	ointment	Unspecified	Specialist INITIATION	Primary care initiation is permissible following specialist advice For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Hydromol®	ointment	Unspecified	Specialist INITIATION	Primary care initiation is permissible following specialist advice For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Urea	10% cream	Unspecified	Specialist INITIATION	Primary care initiation is permissible following specialist advice For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Balneum Plus®	cream urea 5%, lauromacrogols 3%	Unspecified	Specialist INITIATION	Primary care initiation is permissible following specialist advice For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	ImuDerm®	emollient urea 5%, glycerol 5%	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Dermatonics Once®	Heel balm urea 25%	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Dermol®	cream benzalkonium chloride 0.1%, chlorhexidine 0.1%	Unspecified	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children.
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Dermol® 500	lotion benzalkonium chloride 0.1%, chlorhexidine 0.1%	Unspecified	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children.
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Miclaro® Oat	cream	Second line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children.
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Hydromol Intensive®	cream	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children.
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Flexitol®	10% urea cream, 25% urea heel balm	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children.
13. Skin	13.2 Emollients & barrier preparations	13.2.1 Emollients	Aproderm® colloidal oat	cream	First line	All prescribers	For use in line with NW London Prescribing Guideline for Emollients (Including Bath and Shower Preparations) for Adults and Children.
13. Skin	13.2 Emollients & barrier preparations	13.2.2 Barrier preparations	Metanium ®	ointment	Unspecified	All prescribers	
13. Skin	13.2 Emollients & barrier preparations	13.2.2 Barrier preparations	Conotrane ®	cream	Unspecified	All prescribers	
13. Skin	13.3 Topical local anaesthetics and antipruritics	13.3	Crotamiton	cream 10%	Unspecified	All prescribers	
13. Skin	13.3 Topical local anaesthetics and antipruritics	13.3	Doxepin	cream 5%	Unspecified	All prescribers	
13. Skin	13.3 Topical local anaesthetics and antipruritics	13.3	Menthol	Menthol 2% in Aqueous cream	Unspecified	All prescribers	
13. Skin	13.4 Topical corticosteroids	13.4	Hydrocortisone	cream 1%, ointment 1%	First line	All prescribers	1st line mild corticosteroid
13. Skin	13.4 Topical corticosteroids	13.4	Clobetasone butyrate	cream 0.05%; ointment 0.05% (Eumovate ®)	First line	All prescribers	1st line moderate potency corticosteroid
13. Skin	13.4 Topical corticosteroids	13.4	Mometasone furoate	cream 0.1%; ointment 0.1%;	First line	All prescribers	1st line potent corticosteroid
13. Skin	13.4 Topical corticosteroids	13.4	Hydrocortisone butyrate	cream 0.1% scalp lotion 0.1%	Unspecified	All prescribers	Potent
13. Skin	13.4 Topical corticosteroids	13.4	Fluticasone propionate	cream 0.05% ointment 0.005%	Unspecified	All prescribers	Potent
13. Skin	13.4 Topical corticosteroids	13.4	Betamethasone (as valerate)	cream 0.1%; ointment 0.1%; lotion 0.1%; scalp application 0.1% (Betacap ®); foam scalp application 0.1% (Bettamousse®); cream 0.025%; ointment 0.025%; medicated plasters 2.25mg	Unspecified	All prescribers	Potent Betamethasone scalp application 0.1% (Betacap ®) is a 1st line potent corticosteroid scalp application. Betamethasone medicated plasters should only be used in place of cream or ointment when they are required because of the site of application or when there is another clear clinical benefit.
13. Skin	13.4 Topical corticosteroids	13.4	Fluocinolone acetonide	Fluocinolone acetonide 0.00625% cream (Synalar 1 in 4 Dilution) Fluocinolone acetonide 0.00625% ointment (Synalar 1 in 4 Dilution) Fluocinolone acetonide 0.025% cream (Synalar) Fluocinolone acetonide 0.025% gel (Synalar) Fluocinolone acetonide 0.025% ointment (Synalar)	Unspecified	All prescribers	Synalar 1 in 4 Dilution - moderate Synalar - potent
13. Skin	13.4 Topical corticosteroids	13.4	Clobetasol propionate	cream 0.05%; ointment 0.05%; scalp application 0.05% (Dermovate ®); shampoo 0.05% (Etrivex ®)	Unspecified	All prescribers	Very potent
13. Skin	13.4 Topical corticosteroids	13.4	Diffucortolone valerate	oil cream 0.3%; ointment 0.3% (Nerisone Forte ®)	Unspecified	All prescribers	Very potent
13. Skin	13.4 Topical corticosteroids	13.4	Diprosalic ®	ointment, betamethasone 0.05%, salicylic acid 3%; scalp application, betamethasone 0.05%, salicylic acid 2%	Unspecified	All prescribers	Potent; with salicylic acid
13. Skin	13.4 Topical corticosteroids	13.4	Clotrimazole 1%, hydrocortisone 1% (Canestan HC ®)	cream hydrocortisone 1%, clotrimazole 1%	Unspecified	All prescribers	Mild potency; with antifungal
13. Skin	13.4 Topical corticosteroids	13.4	Fusidic acid 2%, hydrocortisone 1% (Fucidin H ®)	cream, hydrocortisone acetate 1%, fusidic acid 2%	Unspecified	All prescribers	Mild potency; with antibiotic
13. Skin	13.4 Topical corticosteroids	13.4	Timodine®	cream, hydrocortisone 0.5%, nystatin 100000 units/g, benzalkonium chloride solution 0.2%, dimeticon 350' 10%	Unspecified	All prescribers	Mild potency; with antimicrobials
13. Skin	13.4 Topical corticosteroids	13.4	Trimovate®	cream, clobetasone butyrate 0.05%, oxytetracycline 3%, nystatin 100 000 units/g	Unspecified	All prescribers	Moderate potency; with antimicrobials
13. Skin	13.4 Topical corticosteroids	13.4	Fucibet ®	cream, betamethasone 0.1%, fusidic acid 2%; Lipid cream betamethasone 0.1%, fusidic acid 2%	Unspecified	All prescribers	Potent corticosteroid; with antibiotic
13. Skin	13.4 Topical corticosteroids	13.4	Lotriderm ®	cream, betamethasone dipropionate 0.064%, clotrimazole 1%	Unspecified	Specialist INITIATION	Potent corticosteroid; with antifungal
13. Skin	13.4 Topical corticosteroids	13.4	Betamethasone 0.1%, neomycin sulphate 0.5%	cream & ointment betamethasone 0.1%, neomycin sulphate 0.5%	Unspecified	Specialist INITIATION	Potent corticosteroid; with antibiotic
13. Skin	13.4 Topical corticosteroids	13.4	Betamethasone 0.1%, clioquinol 3%	cream & ointment betamethasone 0.1%, clioquinol 3%	Unspecified	All prescribers	Potent corticosteroid; with antimicrobial
13. Skin	13.5 Preparations for eczema and psoriasis	13.5.2 Preparations for psoriasis	Dovobet ®	ointment & gel, betamethasone 0.05%, calcipotriol 50micrograms/g	First line	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
13. Skin	13.5 Preparations for eczema and psoriasis	13.5.2 Preparations for psoriasis	Enstilar®	cutaneous foam spray, betamethasone (as dipropionate) 0.5mg/g, calcipotriol (as monohydrate) 50micrograms/g	Second line	All prescribers	For use in line with NICE's guidance
13. Skin	13.5 Preparations for eczema and psoriasis	13.5.2 Preparations for psoriasis	Coal tar, salicylic acid and sulphur (Sebco ®)	scalp ointment, coal tar solution 12%, salicylic acid 2%, precipitated sulphur 4% in a coconut oil emollient basis (Sebco®).	Unspecified	All prescribers	
13. Skin	13.5 Preparations for eczema and psoriasis	13.5.2 Preparations for psoriasis	Psoriderm	Scalp lotion (Shampoo) Coal tar 2.5%, Lecithin 0.3%	Unspecified	All prescribers	
13. Skin	13.5 Preparations for eczema and psoriasis	13.5.2 Preparations for psoriasis	Calcitriol (1,25-dihydroxycolecalciferol)	ointment 3micrograms/g	Unspecified	All prescribers	
13. Skin	13.5 Preparations for eczema and psoriasis	13.5.2 Preparations for psoriasis	Calcipotriol	ointment 50 micrograms/g;	Unspecified	All prescribers	
13. Skin	13.5 Preparations for eczema and psoriasis	13.5.3 Drugs affecting the immune response	Tacrolimus	ointment 0.03%, 0.1%	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
13. Skin	13.5 Preparations for eczema and psoriasis	13.5.3 Drugs affecting the immune response	Pimecrolimus	cream 1%	Unspecified	Specialist INITIATION	For use in line with NICE's guidance
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Benzoyl peroxide	5% gel	Unspecified	All prescribers	For mild acne
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Azelaic acid	cream 20%; 15% gel	Unspecified	All prescribers	For mild acne
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Adapalene	cream 0.1%; gel 0.1%	Unspecified	All prescribers	For mild to moderate acne
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Epiduo®	Adapalene 0.1% / Benzoyl peroxide 2.5% gel; Adapalene 0.3% / Benzoyl peroxide 2.5% gel	Unspecified	All prescribers	For mild to moderate acne
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Benzoyl peroxide 5%/ clindamycin 1%	gel benzoyl peroxide 5%, clindamycin 1%	Unspecified	All prescribers	When possible use non-antibiotic antimicrobials rather than antibiotics. For mild to moderate acne.
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Trifarotene	50micrograms/g cream	Third line	All prescribers	For use in line with NICE guidelines NG198, when the specified treatment options are not tolerated or contraindicated: • For the initial 12 weeks' treatment course, after the combination therapies or topical benzoyl peroxide, trifarotene to be used as a third line option • For maintenance treatment, after fixed combination of topical adapalene and topical benzoyl peroxide or topical monotherapy with adapalene, azelaic acid, or benzoyl peroxide, trifarotene to be used as a third line option
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Clindamycin	topical solution 1%; lotion 1%	Unspecified	All prescribers	When possible use non-antibiotic antimicrobials rather than antibiotics.
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Erythromycin	topical solution, erythromycin 40mg, zinc acetate 12mg/ml (Zineryl ®)	Unspecified	All prescribers	When possible use non-antibiotic antimicrobials rather than antibiotics.
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Aknemycin® Plus	solution tretinoin 0.025%, erythromycin 4%	Unspecified	All prescribers	When possible use non-antibiotic antimicrobials rather than antibiotics.
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Treclin®	gel tretinoin 0.025%, clindamycin (as phosphate) 1%	Unspecified	All prescribers	When possible use non-antibiotic antimicrobials rather than antibiotics.
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Nicotinamide	gel 4%	Unspecified	All prescribers	
13. Skin	13.6 Acne and rosacea	13.6.1 Topical preparations for acne	Clascoterone	cream 10mg/g	Unspecified	Specialist INITIATION	To be used as monotherapy for moderate-to-severe facial acne vulgaris as a second or third line treatment option for the following cohorts: - Where patients do not wish to take oral antibiotics - As an alternative to oral contraceptives - Where spironolactone may be indicated Specialist to prescribe until confirmation of response at 12 weeks then continuation in primary care
13. Skin	13.6 Acne and rosacea	13.6.2 Oral preparations for acne	Co-cyprindiol	tablets, cyproterone acetate 2mg, ethinylestradiol 35micrograms	Unspecified	All prescribers	See BNF. Prescribing this by the approved name co-cyprindiol rather than by the brand name Dianette® reduces the cost.
13. Skin	13.6 Acne and rosacea	13.6.3 Topical preparations for rosacea	Brimonidine tartrate	gel 3.3mg/g (0.33%) brimonide (equivalent to 5mg/g of brimonide tartrate)	Unspecified	All prescribers	
13. Skin	13.6 Acne and rosacea	13.6.3 Topical preparations for rosacea	Ivermectin	cream 10mg/g (1%)	Unspecified	All prescribers	
13. Skin	13.7 Preparations for warts and callouses	13.7 Preparations for warts and calluses	Silver nitrate	95% caustic pencils	Unspecified	All prescribers	
13. Skin	13.7 Preparations for warts and callouses	13.7 Preparations for warts and calluses	Salactol ®	paint, salicylic acid 16.7%, lactic acid 16.7% in flexible collodion	First line	All prescribers	
13. Skin	13.7 Preparations for warts and callouses	13.7 Preparations for warts and calluses	Occlusal®	solution salicylic acid 26%	Third line	All prescribers	
13. Skin	13.7 Preparations for warts and callouses	13.7 Preparations for warts and calluses	Imiquimod	cream 5%	Unspecified	All prescribers	
13. Skin	13.8 Sunscreens and camouflagers	13.8.1 Sunscreen preparations	Anthelios sunscreen lotion SPF 50+	sunscreen lotion SPF 50	Unspecified	All prescribers	To be used in line with ACBS indication
13. Skin	13.8 Sunscreens and camouflagers	13.8.1 Sunscreen preparations	Diclofenac sodium	gel 3% in a sodium hyaluronate basis (Solaraze ®)	Unspecified	All prescribers	
13. Skin	13.8 Sunscreens and camouflagers	13.8.1 Sunscreen preparations	Fluorouracil	cream 5%	Unspecified	Specialist INITIATION	Can be initiated in primary care following recommendation of a specialist for actinic keratosis only.
13. Skin	13.8 Sunscreens and camouflagers	13.8.1 Sunscreen preparations	Tirbanibulin	10mg/g ointment 250mg sachets	Second line	All prescribers	For use, as second line option after fluorouracil 5% cream for treating non-hyperkeratotic, non-hypertrophic actinic keratosis (Olsen grade 1) of the face or scalp in adults. The third line option is diclofenac 3% gel or Fluorouracil 0.5% / Salicylic acid 10% cutaneous solution (Aktierall 5mg/g / 100mg/g cutaneous solution)

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
13. Skin	13.8 Sunscreens and camouflagers	13.8.1 Sunscreen preparations	Actikerall®	solution, fluorouracil 0.5%, salicylic acid 10%	Unspecified	Specialist INITIATION	
13. Skin	13.9 Shampoos and other preparations for scalp and hair conditions	13.9	Capasal ®	shampoo	Unspecified	All prescribers	
13. Skin	13.9 Shampoos and other preparations for scalp and hair conditions	13.9	Ketoconazole	shampoo 2%	Second line	All prescribers	
13. Skin	13.9 Shampoos and other preparations for scalp and hair conditions	13.9	Eflornithine	cream 11.5%	Unspecified	All prescribers	
13. Skin	13.10 Anti-infective skin preparations	13.10.1	Mupirocin	cream 2%; ointment 2%	Unspecified	All prescribers	
13. Skin	13.10 Anti-infective skin preparations	13.10.1	Metronidazole	cream, gel 0.75%	Unspecified	All prescribers	Prescribe by brand name. The different brands have different licensed indications.
13. Skin	13.10 Anti-infective skin preparations	13.10.1	Fusidic acid	cream 2%; ointment 2%	Unspecified	All prescribers	Prefer an antibacterial that is not used systemically where possible, to minimise the development of resistant organisms.
13. Skin	13.10 Anti-infective skin preparations	13.10.2	Miconazole nitrate	cream 2%	Unspecified	All prescribers	
13. Skin	13.10 Anti-infective skin preparations	13.10.2	Terbinafine	cream 1%	Unspecified	All prescribers	
13. Skin	13.10 Anti-infective skin preparations	13.10.2	Ketoconazole	cream 2%	Unspecified	All prescribers	
13. Skin	13.10 Anti-infective skin preparations	13.10.2	Clotrimazole	cream 1%; spray 1%; solution 1%	Unspecified	All prescribers	
13. Skin	13.10 Anti-infective skin preparations	13.10.2	Amorolfine	nail lacquer 5%	Unspecified	All prescribers	
13. Skin	13.10 Anti-infective skin preparations	13.10.3	Aciclovir	cream 5%	Unspecified	All prescribers	
13. Skin	13.10 Anti-infective skin preparations	13.10.4	Permethrin	cream rinse 1%; cream 5%	Unspecified	All prescribers	
13. Skin	13.10 Anti-infective skin preparations	13.10.4	Malathion	liquid 0.5%	Unspecified	All prescribers	
13. Skin	13.11 Skin cleansers, antiseptics, and desloughing agents.	13.11.6	Hydrogen peroxide	cream 1%	Unspecified	All prescribers	
13. Skin	13.12 Antiperspirants.	13.12	Aluminium chloride hexahydrate	20% roll-on applicator	Unspecified	All prescribers	Guidance on usage is at www.sweathelp.com
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Typhoid vaccine	injection, Typhim Vi®	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Tuberculin Purified Protein Derivative (Tuberculin PPD)	20 units per ml (2 units in 0.1ml); 100 units per ml (10 units in 0.1ml)	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Respiratory syncytial virus (RSV) vaccine (Abrysvo®)	powder and solvent for solution for injection 0.5ml vials	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Shingles (Herpes Zoster) vaccine (Shingrix®)	powder and suspension for suspension for injection 0.5ml vials	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Pneumococcal polysaccharide vaccine	injection, Pneumovax 23	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Pneumococcal polysaccharide conjugate vaccine (adsorbed)	injection Prevenar 13 ®	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Meningococcal polysaccharide A, C, W135 and Y vaccine	Injection,	Unspecified	All prescribers	NHS England advise that this vaccine should not be prescribed on the NHS exclusively for the purpose of travel.
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Meningococcal group B vaccine	injection	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Measles, mumps and rubella (MMR) vaccine	injection	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Influenza vaccine, seasonal	LAIV or inactivated	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Human papilloma virus vaccine	Gardasil 9 vaccine suspension for injection 0.5ml pre-filled syringes	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Hepatitis B vaccine		Unspecified	All prescribers	NHS England advise that this vaccine should not be prescribed on the NHS exclusively for the purpose of travel.
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Hepatitis A with hepatitis B vaccine	injection 20mg in 1ml, 1ml pre-filled syringe (Twinrix ® Adult); 0.5ml pre-filled syringe (Twinrix ® Paediatric)	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Hepatitis A vaccine	injection (Avaxim®, Vaqta® Paediatric)	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Diphtheria / Tetanus / Pertussis (acellular component) / Hepatitis B (surface antigen) / Poliomyelitis (inactivated) / Haemophilus type b conjugate vaccine	injection Vaxelis; Infanrix Hexa	Unspecified	All prescribers	

BNF chapter	BNF section	BNF sub-section	Drug name (shading indicates that the drug appears in more than one section)	Formulations	Order of preference	Initiate by specialist only?	Comments
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Diphtheria, tetanus, pertussis and inactivated poliomyelitis (absorbed) - DTap/IPV	injection	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Diphtheria (low dose), tetanus, pertussis and inactivated poliomyelitis (absorbed) vaccine	injection	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	Diphtheria (low dose), tetanus and inactivated poliomyelitis vaccine	injection	Unspecified	All prescribers	
14. Immunological products and vaccines	14.4 Vaccines and antisera	14.4	BCG intradermal	Bacillus Calmette-Guerin Vaccine (Live) Intradermal injection	Unspecified	All prescribers	NHS England advise that this vaccine should not be prescribed on the NHS exclusively for the purpose of travel.
14. Immunological products and vaccines	14.5 Immunoglobulins	14.5	Tetanus immunoglobulin	injection	Unspecified	All prescribers	
15. Anaesthesia	15.1 General anaesthesia	15.1.3 Antimuscarinic drugs	Hyoscine hydrobromide	injection 400micrograms in 1ml, 600micrograms in 1ml	Unspecified	All prescribers	
15. Anaesthesia	15.1 General anaesthesia	15.1.3 Antimuscarinic drugs	Glycopyrronium bromide (Glycopyrrolate)	injection 200micrograms in 1ml, 600micrograms in 3ml	Unspecified	All prescribers	
15. Anaesthesia	15.1 General anaesthesia	15.1.4.1 Benzodiazepines	Midazolam	injection 1mg/ml 5ml ampoules	Unspecified	For use in palliative care	For use in palliative care. See palliative care section in the BNF and the NPSA Rapid Response Report on reducing risk of overdose with midazolam injection in adults. Ensure that flumazenil is available where midazolam is used.
15. Anaesthesia	15.1 General anaesthesia	15.1.4.2 Non-opioid analgesics	Ketorolac trometamol	injection 30mg/ml	Unspecified	Specialist INITIATION	
	15.1 General anaesthesia	15.1.4.3 Opioid analgesics	Alfentanil	injection 500micrograms/ml (2ml ampoules)	For use only in palliative care	Specialist INITIATION	For use only in palliative care.
15. Anaesthesia	15.1 General anaesthesia	15.1.7 Antagonists for central and respiratory depression	Naloxone hydrochloride	injection 400micrograms in 1ml	Unspecified	All prescribers	
	15.1 General anaesthesia	15.1.7 Antagonists for central and respiratory depression	Flumazenil	injection 100micrograms/ml 5ml ampoules	Unspecified	All prescribers	Ensure that flumazenil is available where midazolam is used.
15. Anaesthesia	15.2 Local anaesthesia	15.2	Lidocaine	plasters 5% (700mg/plaster)	Unspecified	See comments column	High cost. Not recommended in NICE's guideline on neuropathic pain (Neuropathic pain - pharmacological management CG173). NHS England guidance https://www.england.nhs.uk/publication/items-which-should-not-be-routinely-prescribed-in-primary-care-guidance-for-cogs/ Should not be initiated in primary care and should be deprescribed in all patients (apart from exception below*). If, in exceptional circumstances, there is a clinical need for lidocaine plasters to be prescribed in primary care, this should be undertaken in a cooperation arrangement with a multi-disciplinary team and/or other healthcare professional. *Exception: The recommendations above do not apply to patients who have been treated in line with NICE CG173 on neuropathic pain but are still experiencing neuropathic pain associated with previous herpes zoster infection (post-herpetic neuralgia).
15. Anaesthesia	15.2 Local anaesthesia	15.2	Lidocaine with chlorhexidine	Gel containing lidocaine hydrochloride 2% and chlorhexidine gluconate 0.25%, 6ml, 11ml syringe (Instillagel ®).	Unspecified	All prescribers	
15. Anaesthesia	15.2 Local anaesthesia	15.2	Lidocaine hydrochloride	ointment 5%; injection 5mg/ml (0.5%)	Unspecified	All prescribers	
15. Anaesthesia	15.2 Local anaesthesia	15.2	EMLA®	cream, lidocaine 2.5%, prilocaine 2.5%,	Unspecified	All prescribers	
Appendix 2 Borderline substances	A2.4 Feed supplements	A2.4.2 Fibre, vitamin and mineral supplements	Renavit®	450mg tablets	Unspecified	All prescribers	ACBS Indications Dietary management of water-soluble vitamin deficiency in adults with renal failure on dialysis