

Reasoning analysis — x-ai/grok-4.6 (50 runs, `real_grok46_50`)

Each model rationale is split into sentences; sentences making the same point across runs are grouped. **Shared** = the point appears in at least 60% of the runs in that group; **varying** = it appears in some runs but not most. Counts are runs, not sentences. 'Evidence cited' lists the numbers and times the model quoted, with the number of runs quoting each.

decision: Profile / basal 00:00

keep — 50/50 runs

Shared reasoning

- (40/50) Evidence: local 00:00-03:00 TBR 10-18% with means near 90-105 mg/dL.

Varying reasoning

- (12/50) Dependencies: do not cut this basal in the same step as an overnight target review.
- (11/50) Dependencies: target 90, always-on SMB, DIA 10 h, and afternoon UAM stacks.
- (10/50) Change unsupported.
- (8/50) Dependencies: overnight target 90 mg/dL and Dynamic ISF or SMB stacking.
- (8/50) Uncertainty: residual evening IOB and DIA 10 h are not separated from this rate.
- (6/50) Dependencies: do not change with target, DIA, or SMB delivery.
- ... 17 more varying points

Points made by a single run only: 32.

Phrases used by at least half the runs: “target 90” (33), “dia 10” (26), “90 mg/dl” (25).

Evidence cited: 03:00 (35), 00:00 (31), 10h (27), 90mg/dl (25), 0.6u/h (10), 18.5% (10), 23:00 (9), 18% (9), 106mg/dl (6), 04:00 (6), 105mg/dl (5), 08:00 (5).

decision: Profile / basal 05:00

keep — 50/50 runs

Varying reasoning

- (25/50) Uncertainty: one 42 mg/dL near 07:19 local on 21 Aug.
- (20/50) Evidence: local 05:00-06:00 average about 104 mg/dL with high time in range and low time below range.
- (15/50) Evidence: local 05:00 TIR 95.2% and mean 104.2 mg/dL.
- (13/50) Evidence: local 04:00-06:00 TIR 95-97.5% with low TAR.
- (10/50) No isolated fasting defect.
- (9/50) Dependencies: 06:00 basal and overnight IOB.
- ... 16 more varying points

Points made by a single run only: 29.

Phrases used by at least half the runs: “42 mg/dl” (28), “local 05 00” (25).

Evidence cited: 05:00 (36), 06:00 (32), 42mg/dl (28), 104.2mg/dl (15), 95.2% (13), 07:19 (9), 4.8% (9), 07:00 (9), 06:19 (8), 00:00 (7), 104mg/dl (7), 04:00 (7).

decision: Profile / basal 06:00

keep — 50/50 runs

Shared reasoning

- (34/50) Evidence: local 06:00-08:00 remains in range with TIR 97.5-98.7% and means near 100-110 mg/dL.

Varying reasoning

- (12/50) Uncertainty: one 42 mg/dL point on 21 Aug 06:19 is not a repeated dawn defect.
- (7/50) Dependencies: later 08:00-10:00 rise is meal-timed.
- (5/50) Uncertainty: later 09:00-10:00 rise can be breakfast UAM.
- (4/50) Evidence: hours 6-8 local TIR 98-99% and almost no TAR, so dawn basal is not insufficient.
- (4/50) Dependencies: 05:00 step and morning meals after 08:00.
- (4/50) Dependencies: 08:00 target 100 mg/dL and unlogged intake.
- ... 21 more varying points

Points made by a single run only: 47.

Evidence cited: 08:00 (41), 06:00 (40), 99% (16), 110mg/dl (15), 42mg/dl (12), 09:00 (11), 10:00 (11), 07:00 (9), 98.7% (9), 05:00 (5), 06:19 (5), 07:19 (4).

decision: Profile / basal 10:00

keep — 50/50 runs

Varying reasoning

- (13/50) Uncertainty: meals versus basal cannot be split.
- (9/50) Dependencies: unused CR 13.2 g/U and always-on SMB.
- (8/50) Dependencies: CR unused;
- (6/50) Dependencies: meal logging before any daytime basal change.
- (6/50) Not labeled a basal defect.
- (6/50) Uncertainty: unlogged intake and GLP-1.
- ... 28 more varying points

Points made by a single run only: 38.

Phrases used by at least half the runs: “late morning” (38).

Evidence cited: 10:00 (15), 09:00 (7), 20.2% (7), 13.2g/u (6), 08:00 (5), 19:00 (4), 18:00 (3), 13:00 (2), 20% (2), 15:00 (2), 148mg/dl (2), 12:00 (2).

decision: Profile / basal 16:00

keep — 50/50 runs

Varying reasoning

- (19/50) Evidence: 16:00 local has the highest TAR then 18:00-19:00 TBR.
- (7/50) Evidence: local 16:00 is the peak time-above-range hour, but 18:00-19:00 time below range follows the same afternoons.
- (6/50) Uncertainty: meal versus under-basal.
- (5/50) Evidence is 14:00-17:00 TAR then 19:00 TBR.
- (5/50) Dependencies: UAM, Dynamic ISF, CR 10.5 g/U, and evening corrections; do not raise this with any of those.
- (4/50) Dependencies: evening hypo risk.
- ... 29 more varying points

Points made by a single run only: 46.

Evidence cited: 19:00 (36), 16:00 (30), 18:00 (21), 32.1% (14), 0.5u/h (13), 15.3% (7), 14:00 (6), 42mg/dl (5), 17:00 (4), 162.8mg/dl (3), 10.5g/u (2), 10h (2).

decision: Profile / basal 18:00

keep — 50/50 runs

Varying reasoning

- (15/50) Evidence: local 18:00 average 124.2 mg/dL with mixed TAR and TBR, then 19:00 TBR 15.3 percent and 41 mg/dL extremes.
- (11/50) Evidence: glucose is already falling from the afternoon peak through 18:00, with 7.7% below range.
- (11/50) Dependencies: 16:00 basal, 19:00 basal, and DIA.
- (9/50) Uncertainty: residual IOB versus basal.
- (7/50) Evidence: local hour 18 already shows TBR 7.7% after the afternoon peak.
- (7/50) Uncertainty: activity and GLP-1 are unmeasured.
- ... 20 more varying points

Points made by a single run only: 51.

Evidence cited: 18:00 (40), 19:00 (30), 7.7% (22), 16:00 (16), 42mg/dl (8), 10h (8), 124.2mg/dl (7), 15.3% (4), 0.6u/h (3), 0.5u/h (2), 41mg/dl (1), 0.1u/h (1).

decision: Profile / basal 19:00

keep — 50/50 runs

Varying reasoning

- (24/50) Evidence: local 19:00 TBR 15.3% and minima 41 mg/dL.
- (11/50) Evidence: hour 19:00 local has 15.3 percent time below range.
- (7/50) Dependencies: DIA 10 h and UAM IOB.
- (7/50) Dependencies: overnight target 90 and DIA 10 h.
- (5/50) Dependencies: do not lower this in the same step as SMB, DynISF, or 16:00 basal.
- (4/50) Uncertainty: delayed afternoon SMB versus this 0.7 U/h step.
- ... 17 more varying points

Points made by a single run only: 71.

Phrases used by at least half the runs: “41 mg/dl” (26), “tbr 15.3” (26).

Evidence cited: 15.3% (42), 19:00 (37), 41mg/dl (26), 0.7u/h (20), 10h (11), 16:00 (7), 18:00 (3), 23:00 (2), 105.4mg/dl (2), 00:00 (1), 14:00 (1), 17:00 (1).

decision: Profile / CR 00:00

keep — 50/50 runs

Varying reasoning

- (27/50) Evidence: zero carb entries and cob 0, so this CR is unused.
- (17/50) Dependencies: meal logging must exist before any CR review.
- (9/50) No meal-linked evidence supports a change.
- (8/50) Dependencies: UAM meal path.
- (8/50) Uncertainty: no overnight meal tests.
- (6/50) Evidence: zero carb treatments, so this CR was not used.
- ... 19 more varying points

Points made by a single run only: 28.

Phrases used by at least half the runs: “carb entries” (33), “zero carb” (25).

Evidence cited: 13g/u (1).

decision: Profile / CR 04:00

keep — 50/50 runs

Varying reasoning

- (17/50) Evidence: no carb treatments in this or any segment.
- (8/50) Uncertainty: unused setting cannot be validated.
- (8/50) Dependencies: empty declared meal strategy.
- (8/50) Dependencies: meal logging.
- (7/50) 04:00-08:00 glucose is already stable.
- (7/50) Evidence: no carb treatments in the early-morning CR window.
- ... 24 more varying points

Points made by a single run only: 29.

Evidence cited: 08:00 (5), 04:00 (5), 11.1g/u (4), 06:00 (1), 05:00 (1), 01:00 (1).

decision: Profile / CR 08:00

keep — 50/50 runs

Varying reasoning

- (8/50) Evidence: local 08:00-10:00 rise occurs with COB 0, so UAM, not this CR, dosed any meal.
- (8/50) Uncertainty: meal size unknown.
- (8/50) Uncertainty: breakfast composition unknown.
- (7/50) Evidence: late-morning rise occurs with COB 0, so UAM, not this CR, delivered insulin.
- (7/50) Dependencies: do not tighten CR while always-on UAM remains the meal path.
- (5/50) Morning rises occurred without announced carbs, so this CR was not the dosing path.
- ... 27 more varying points

Points made by a single run only: 46.

Evidence cited: 08:00 (8), 10:00 (8), 13.2g/u (6), 09:00 (4), 12:00 (1), 11:00 (1).

decision: Profile / CR 16:00

keep — 50/50 runs

Varying reasoning

- (9/50) Evidence: strongest CR sits on the same clock as peak TAR and the 18:00-19:00 crash, but it was unused.
- (9/50) Dependencies: afternoon UAM and 16:00 basal.
- (7/50) Uncertainty: high.
- (6/50) Evidence: tightest CR sits under the worst TAR hour, but no carbs were entered so it did not dose meals.
- (5/50) Dependencies: do not tighten CR while TBR is binding.
- (4/50) Dependencies: keep until carbs exist.
- ... 24 more varying points

Points made by a single run only: 65.

Evidence cited: 16:00 (14), 10.5g/u (12), 19:00 (10), 18:00 (4), 0.5u/h (2), 10h (1).

decision: Profile / CR 20:00

keep — 50/50 runs

Varying reasoning

- (10/50) Evidence: no evening carb entries.
- (10/50) Dependencies: DIA, IOB, overnight target.
- (10/50) Uncertainty: no dinner carb log.
- (7/50) Evidence: evening TBR with cob 0 is not a CR test.
- (7/50) Dependencies: leftover afternoon IOB dominates this window more than CR.
- (5/50) Uncertainty: leftover afternoon IOB versus any late snack cannot be split.
- ... 28 more varying points

Points made by a single run only: 27.

Evidence cited: 10h (5), 19:00 (3), 12.6g/u (2), 90mg/dl (2), 20:00 (2), 22:00 (1), 23:00 (1).

decision: Profile / ISF 00:00

keep — 50/50 runs

Varying reasoning

- (12/50) Dependencies: do not change profile ISF in the same step as DynISF 70%.
- (11/50) Evidence: Dynamic ISF is on and sampled variable_sens ranged about 28-124 mg/dL/U, so 56 is not the delivered ISF.
- (8/50) Enacted sensitivity is Dynamic ISF, with variable_sens far from 56.
- (8/50) Dependencies: Dynamic ISF 70 percent; do not change ISF and Dynamic ISF together.
- (7/50) Evidence: enacted variable_sens ranged about 28-124 mg/dL/U under DynISF 70%, so 56 was not the delivered ISF.
- (7/50) Dependencies: DynISF 70%, SMB, DIA.
- ... 24 more varying points

Points made by a single run only: 54.

Phrases used by at least half the runs: “dynamic isf” (29).

Evidence cited: 70% (31), 124mg/dl (6), 10h (4), 56mg/dl (4), 5.1% (2), 00:00 (1), 0.2u (1), 178mg/dl (1), 20u (1), 43mg/dl (1), 180mg/dl (1).

decision: Profile / target 00:00

keep — 40/50 runs

Varying reasoning

- (16/40) Dependencies: 23:00 target, DIA, basal 00:00.
- (10/40) Evidence: overnight mean is near 90 mg/dL while time below range is high, which can reflect leftover IOB rather than a wrong target.
- (8/40) Evidence: 00:00-03:00 TBR is high under this target.
- (6/40) Dependencies: this is the first profile lever to re-review after low-IOB nights, but not a supported change until that check, and not in the same step as basal.
- (6/40) Evidence: overnight target 90 mg/dL sits in a high-TBR band, but lows may be residual IOB rather than the target number.
- (6/40) Uncertainty: residual IOB may be the driver rather than the target number.
- ... 14 more varying points

Points made by a single run only: 47.

Evidence cited: 00:00 (21), 90mg/dl (18), 03:00 (8), 23:00 (7), 01:00 (6), 0.6u/h (3), 18% (3), 18.5% (3), 32.6% (2), 10h (2), 90.6mg/dl (2), 101mg/dl (1).

change — 10/50 runs

Varying reasoning

- (5/10) Local 00:00-03:00 TBR is 10-18% while means are already about 90-106 mg/dL.
- (5/10) Dependencies: change together with 23:00 as one overnight alignment and do not change basal, ISF, SMB, or Dynamic ISF simultaneously.
- (4/10) Evidence: local 00:00 average 93.3 mg/dL with 11.4% below range and 0% above; local 01:00 average 90.6 mg/dL with 18.5% below range and 0% above, all under this 90 mg/dL segment.
- (4/10) Uncertainty: residual afternoon IOB may still cause lows at 100 mg/dL.
- (3/10) Candidate 100 mg/dL already exists at 08:00.
- (3/10) Dependency: do not also change 00:00 basal, 23:00 target, SMB, or DynISF in the same step.

- ... 1 more varying points

Points made by a single run only: 12.

Phrases used by at least half the runs: “100 mg/dl” (7), “local 00 00” (5), “00 target” (5), “uncertainty residual” (5).

Evidence cited: 00:00 (9), 100mg/dl (7), 18.5% (7), 23:00 (7), 01:00 (6), 08:00 (6), 03:00 (6), 11.4% (3), 90.6mg/dl (3), 90mg/dl (3), 18% (3), 0% (1).

decision: Profile / target 08:00

keep — 50/50 runs

Varying reasoning

- (10/50) Uncertainty: afternoon TAR is meal/UAM, not this target being too high.
- (9/50) Uncertainty: temporary targets of 135 mg/dL also appear.
- (9/50) Evidence: daytime target 100 mg/dL already sits under afternoon TAR that is better explained by unannounced intake.
- (7/50) Evidence: daytime target 100 is already higher than overnight 90 and aligns with better late-morning TIR than afternoon TAR.
- (6/50) Evidence is 08:00 target segment and 10:00-17:00 TAR with COB 0.
- (4/50) Evidence: daytime target is already higher than night while afternoon TAR is meal-and-UAM timed.
- ... 24 more varying points

Points made by a single run only: 53.

Phrases used by at least half the runs: “daytime target” (37), “target 100” (27), “100 mg/dl” (26), “daytime target 100” (26).

Evidence cited: 100mg/dl (26), 135mg/dl (14), 08:00 (5), 14:00 (4), 17:00 (4), 90mg/dl (2), 00:00 (2), 10:00 (2), 163mg/dl (2), 06:00 (1), 32% (1), 16:00 (1).

decision: Profile / target 23:00

keep — 44/50 runs

Shared reasoning

- (33/44) Dependencies: profile.target.00_00.

Varying reasoning

- (15/44) Uncertainty: same IOB-versus-target confound as 00:00.
- (10/44) Uncertainty: same leftover-IOB confound as the 00:00 target.
- (9/44) Evidence: 23:00 local already has 10.1% below range heading into the overnight 90 mg/dL target.
- (7/44) Evidence: 23:00 local TBR about 10% begins the early-night low cluster on the same 90 mg/dL target.
- (6/44) Evidence: local 23:00 TBR 10.1% begins the overnight low cluster.
- (6/44) Keep both overnight target rows together rather than editing one.
- ... 12 more varying points

Points made by a single run only: 33.

Phrases used by at least half the runs: “target 00 00” (24).

Evidence cited: 00:00 (42), 23:00 (34), 10.1% (22), 90mg/dl (17), 01:00 (6), 10% (5), 105.7mg/dl (4), 10h (2), 03:00 (2), 18.5% (1), 19:00 (1).

change — 6/50 runs

Shared reasoning

- (4/6) Dependencies: pair only with 00:00 target 100 mg/dL; hold basal and automation.
- (4/6) This segment starts the overnight 90 mg/dL window where TBR rises from 23:00.

Varying reasoning

- (3/6) Changing it together with 00:00 is one overnight-target review, not a second experiment.
- (2/6) Uncertainty: evening IOB from afternoon UAM.

Points made by a single run only: 8.

Phrases used by at least half the runs: “90 mg/dl” (5), “overnight 90 mg/dl” (4), “segment starts” (3), “one overnight” (3).

Evidence cited: 00:00 (6), 23:00 (5), 90mg/dl (5), 10.1% (4), 100mg/dl (2), 03:00 (1), 19:00 (1).

decision: Profile / DIA

keep — 50/50 runs

Varying reasoning

- (8/50) Evidence: DIA 10 h can keep afternoon and evening insulin active into 00:00 to 03:00, matching the late low pattern.
- (8/50) Evidence: 10 h can leave afternoon insulin active at 19:00-01:00.
- (7/50) Uncertainty: shortening DIA could hide IOB and worsen lows.
- (7/50) Dependencies: ISF and SMB.
- (6/50) Dependencies: changing DIA with target or SMB would confound IOB accounting.
- (5/50) Uncertainty: free_peak_oref peak is not in the package;
- ... 24 more varying points

Points made by a single run only: 66.

Phrases used by at least half the runs: “shortening dia” (25).

Evidence cited: 10h (23), 19:00 (8), 03:00 (8), 01:00 (7), 6h (7), 00:00 (5), 4h (3), 6u (3), 23:00 (2), 5.1% (2), 25u (1), 5.9u (1).

decision: AAPS / SMB and UAM

keep — 50/50 runs

Varying reasoning

- (17/50) No source component is null, so verify is not used.
- (16/50) Evidence: UAM with SMB matches zero carbs and COB 0 and is the observed meal engine.
- (7/50) Uncertainty: disabling UAM without announced meals would leave afternoon TAR uncovered.

- (7/50) Uncertainty: it also contributes to evening overshoot.
- (6/50) Dependencies: turning UAM off without a carb workflow would leave afternoon rises uncovered; do not pair with the target change.
- (5/50) Evidence: with 0 g carbs, UAM is the only meal path.
- ... 32 more varying points

Points made by a single run only: 58.

Evidence cited: 00:00 (2), 17:00 (2), 16:00 (1), 70% (1), 08:00 (1), 14:00 (1).

decision: AAPS / SMB activation conditions

keep — 50/50 runs

Shared reasoning

- (36/50) High-TT SMB off matches 135 mg/dL temporary targets.

Varying reasoning

- (13/50) Evidence: always-on SMB makes COB and after-carb flags idle, which matches COB 0.
- (7/50) Dependencies: do not change activation with interval or DynISF in one step.
- (5/50) Uncertainty: always-SMB may over-deliver on unannounced rises, but disabling it is not isolated from DynISF.
- (4/50) No component is null, so verify is not used.
- (4/50) Uncertainty: always-on SMB can still dose toward overnight target 90.
- (4/50) Dependencies: overnight TBR and meal strategy.
- ... 28 more varying points

Points made by a single run only: 53.

Phrases used by at least half the runs: “always on smb” (35), “high tt” (35).

Evidence cited: 135mg/dl (23), 90mg/dl (3), 19:00 (1), 110mg/dl (1), 5.1% (1).

decision: AAPS / SMB strength and frequency

keep — 50/50 runs

Varying reasoning

- (6/50) Evidence: configured 3, 15, and 20 minutes look modest versus 0.5 to 0.8 U/h basal, but reason strings show larger custom UAM boosts and SMBs.
- (6/50) Dependencies: do not change delivery in the same step as overnight target.
- (5/50) Dependencies: do not retune minutes in the same step as Dynamic ISF or max basal.
- (5/50) Dependencies: do not change interval and UAM minutes together or with DynISF.
- (4/50) Evidence: 3-minute cadence and UAM boosts can stack on afternoon rises; sampled SMB sometimes exceeded the 15/20 minute inventory story.
- (4/50) Evidence: 3-minute interval and UAM 20 minutes can restack, and delivered SMB sometimes exceeded 15-minute profile-basal math.
- ... 33 more varying points

Points made by a single run only: 70.

Evidence cited: 20min (30), 3min (14), 1.6u (10), 0.8u/h (7), 19:00 (5), 70% (2), 4u/h (2), 16:00 (2), 15min (2), 00:00 (1), 0.3u (1), 10h (1).

decision: AAPS / max basal and max IOB limits

keep — 50/50 runs

Shared reasoning

- (31/50) Evidence: observed IOB peaked near 5.9 U and enacted temps near 3-4 U/h, so 25 U and 12 U/h did not bind.

Varying reasoning

- (4/50) Evidence: 25 U and 12 U/h are loose versus TDD about 20 U and were not binding; hypos occurred at IOB well below 25 U.
- (4/50) Uncertainty: they are loose versus TDD about 20 U.
- (4/50) No null component.
- (3/50) Dependencies: default multipliers are not inventory items and are not routine optimization targets.
- (3/50) Uncertainty: lowering unused ceilings may do nothing until actuators change.
- (3/50) Keep; do not invent a new cap.
- ... 18 more varying points

Points made by a single run only: 89.

Phrases used by at least half the runs: “12 u/h” (32), “near u/h” (30).

Evidence cited: 4u/h (46), 25u (37), 12u/h (32), 6u (23), 5.9u (20), 20u (11), 19.7u (9), 5.935u (3), 16.2u (2), 0.8u/h (1), 19:00 (1), 10:00 (1).

decision: AAPS / Autosens and Dynamic ISF

keep — 50/50 runs

Varying reasoning

- (15/50) Dependencies: do not change DynISF with profile ISF or SMB.
- (14/50) Dependencies: do not enable Autosens on top of Dynamic ISF; do not change the 70 percent factor with profile ISF or SMB delivery.
- (12/50) A sensitivity_ratio of 1 is a neutral current adjustment, not disabled Autosens.
- (9/50) Uncertainty: 70 percent versus SMB interval 3 min.
- (7/50) Autosens off is consistent with Autosens target flags.
- (7/50) Evidence: Autosens is off, so a sensitivity_ratio of 1 is not an Autosens diagnosis.
- ... 35 more varying points

Points made by a single run only: 62.

Phrases used by at least half the runs: “profile isf” (37).

Evidence cited: 70% (46), 56mg/dl (2), 19:00 (2), 00:00 (1), 124mg/dl (1), 5.1% (1), 10h (1), 3min (1), 14:00 (1), 42mg/dl (1).

decision: AAPS / carbohydrate absorption model

keep — 50/50 runs

Varying reasoning

- (11/50) UAM is not extra COB counting.
- (8/50) Evidence: both values are idle with 0 g carbs and COB 0.
- (7/50) Dependencies: become relevant only after carb entries exist.
- (7/50) Dependencies: empty meal strategy.
- (6/50) Uncertainty: no meal-linked absorption curve.
- (5/50) Dependencies: empty meal_strategies and unused CR; no e-carb duration exists to infer.
- ... 31 more varying points

Points made by a single run only: 34.

Evidence cited: 7h (8), 10h (4), 5min (1), 8mg/dl (1).

profile recommendation

target — 33/50 runs

Shared reasoning

- (28/33) Keep every profile segment, including overnight target 90 mg/dL, daytime target 100 mg/dL, and all basal, CR, ISF, and DIA values.

Varying reasoning

- (9/33) Local 01:00 time below range is high near target 90 mg/dL, but leftover UAM and Dynamic ISF IOB, DIA 10 h, unlogged meals, and overnight basal are not separated.
- (8/33) Local 00:00-03:00 TBR is real, but the same lows are compatible with residual afternoon UAM/SMB, DIA 10 h, and SMB-always at target 90.
- (5/33) Night basal is not changed because 04:00-08:00 is already stable.
- (5/33) Package evidence does not isolate one profile segment enough to support an old-to-new value change.
- (4/33) The 08:00 target is already 100 mg/dL, so 100 mg/dL is an existing user setpoint rather than a new invented number.
- (4/33) Afternoon TAR is not a clean 16:00 basal defect while COB is 0.
- ... 11 more varying points

Points made by a single run only: 58.

Phrases used by at least half the runs: “90 mg/dl” (32), “target 90” (20), “overnight target” (20), “100 mg/dl” (18), “00 basal” (18), “target 90 mg/dl” (17).

Evidence cited: 90mg/dl (32), 00:00 (24), 100mg/dl (18), 0.6u/h (16), 10h (15), 08:00 (13), 23:00 (12), 03:00 (10), 16:00 (9), 01:00 (8), 04:00 (8), 18.5% (5).

basal — 17/50 runs

Shared reasoning

- (15/17) Keep every current basal, target, CR, ISF, and DIA value.

Varying reasoning

- (7/17) The only clean fasting-like window is local 04:00-08:00 and it is already stable.
- (4/17) Keep every current profile segment, including overnight basal 0.6 U/h from 00:00 and 0.7 U/h from 19:00, and verify whether the 23:00-03:00 lows persist when IOB is already near zero.

- (4/17) Afternoon highs occur while 16:00 basal is already 0.5 U/h and no carbs were logged, so a basal increase is unsupported.
- (3/17) Overnight means are near the 90 mg/dL target, but 00:00-03:00 and 19:00 lows can be leftover insulin.
- (2/17) The only clean fasting-like window (04:00-08:00 local) is stable on 0.7-0.8 U/h, while early-night TBR and afternoon TAR have stronger competing explanations in unannounced meals, UAM/SMB, Dynamic ISF, DIA 10 h, and target 90.
- (2/17) Overnight TBR with near-zero TAR can be residual UAM and Dynamic ISF IOB plus DIA 10 h, a tight 90 mg/dL target, or slightly strong 00:00 basal.
- ... 6 more varying points

Points made by a single run only: 31.

Phrases used by at least half the runs: “cr isf” (14), “keep every” (14), “isf and dia” (12), “0.6 u/h” (11), “target cr isf” (11), “basal target cr” (11), “0.5 u/h” (9), “dia 10” (9).

Evidence cited: 16:00 (11), 0.6u/h (11), 10h (10), 00:00 (10), 08:00 (10), 0.5u/h (9), 04:00 (7), 90mg/dl (7), 03:00 (7), 19:00 (7), 0.7u/h (4), 05:00 (4).

primary recommendation

target — 33/50 runs

Varying reasoning

- (8/33) Review raising only the 00:00 target from 90 to 100 mg/dL.
- (8/33) Local 00:00-03:00 sits on a 90 mg/dL target with TBR 10-18.5% and a 01:00 mean of 90.6 mg/dL.
- (8/33) Keep the profile and first separate overnight lows from leftover IOB.
- (8/33) Keep the profile and verify overnight lows before any edit.
- (6/33) That does not isolate target 90 mg/dL from 00:00 basal 0.6 U/h.
- (5/33) Overnight lows sit at a 90 mg/dL target with almost no overnight highs, but leftover afternoon IOB and 00:00 basal 0.6 U/h can produce the same pattern.
- ... 15 more varying points

Points made by a single run only: 39.

Phrases used by at least half the runs: “90 mg/dl” (29), “overnight target” (17), “target 90” (17).

Evidence cited: 90mg/dl (29), 00:00 (21), 03:00 (17), 0.6u/h (14), 10h (10), 100mg/dl (9), 23:00 (9), 08:00 (7), 19:00 (4), 01:00 (3), 90.6mg/dl (2), 18.5% (2).

basal — 16/50 runs

Varying reasoning

- (7/16) Keep the current profile and separate leftover IOB from overnight basal and target.
- (4/16) Keep the current basal profile and verify overnight lows against evening insulin.
- (3/16) Evening and 00:00-03:00 lows can come from afternoon UAM or correction stacks and 10 h DIA rather than from 0.6 U/h overnight basal or the 90 mg/dL overnight target.
- (3/16) Evening and overnight lows can still be residual UAM or DynISF IOB, a tight 90 mg/dL target, or overnight basal 0.6 U/h.
- (3/16) Early-night TBR is real, but 04:00-08:00 local control is strong and afternoon highs are not a clean under-basal signal.
- (3/16) Evidence does not isolate one setting, so no basal, target, CR, ISF, or DIA value is changed.
- ... 7 more varying points

Points made by a single run only: 22.

Phrases used by at least half the runs: “target 90” (9), “overnight basal” (9), “90 mg/dl” (8).

Evidence cited: 90mg/dl (8), 0.6u/h (7), 10h (6), 03:00 (5), 04:00 (4), 00:00 (4), 16:00 (4), 08:00 (3), 19:00 (2), 0.7u/h (2), 42mg/dl (1), 0.5u/h (1).

profile — 1/50 runs

Points made by a single run only: 3.

Phrases used by at least half the runs: “catch up dynamic” (1), “meal timed observations” (1), “profile value change” (1), “target 90 mg/dl” (1), “profile and collect” (1), “explained by unannounced” (1), “mg/dl those causes” (1), “timed observations first” (1), “collect meal timed” (1), “intake uam catch” (1), “evening or overnight” (1), “observations first afternoon” (1).

Evidence cited: 10h (1), 90mg/dl (1).

summary

50 runs

Shared reasoning

- (47/50) Over 14 days with 98.7% CGM coverage, mean glucose was 121.6 mg/dL, time in range 86.0%, time above range 8.9%, and time below range 5.1% with CV 32.6%.

Varying reasoning

- (19/50) Afternoon highs (about 14:00-17:00 local) were followed by evening and overnight lows, especially 19:00 and 00:00-03:00.
- (16/50) Competing explanations are not separated, so every profile and AAPS inventory value is kept and the next step is observation rather than a value change.
- (13/50) The selected AAPS profile is complete and declared applicable.
- (11/50) No carbs were logged, so meals were handled by always-on SMB/UAM and Dynamic ISF rather than CR or e-carbs.
- (9/50) No carbs or e-carbs were logged; declared meal strategy fields are empty.
- (7/50) No carb or e-carb entries were recorded, so meals were covered by always-on SMB and UAM with Dynamic ISF, not by CR.
- ... 18 more varying points

Points made by a single run only: 28.

Phrases used by at least half the runs: “time in range” (37), “121.6 mg/dl” (36), “below range” (36), “dynamic isf” (34), “mean glucose” (32), “time below range” (28), “cgm coverage” (28), “afternoon highs” (27).

Evidence cited: 5.1% (44), 121.6mg/dl (36), 86.0% (34), 19:00 (28), 98.7% (25), 8.9% (22), 32.6% (17), 00:00 (15), 03:00 (14), 90mg/dl (13), 14:00 (10), 17:00 (10).

issues

50 runs

Shared reasoning

- (49/50) Time below range 5.1% with clustered local lows at 00:00-03:00 and 19:00, including 41-42 mg/dL.

Varying reasoning

- (28/50) Mid-period profile switch, GLP-1 notes, site changes, and a 19 Aug occlusion reduce period stationarity.
- (20/50) Max IOB 25 U and max basal 12 U/h are far above observed delivery and did not bind.
- (17/50) Afternoon TAR (local 14:00-17:00, 16:00 TAR 32.1%) then evening crash, consistent with late UAM/DynISF stacking.
- (16/50) Afternoon time above range peaks near 14:00-17:00 local and is not separable from unlogged intake.

- (14/50) No carb entries and an empty declared meal-strategy list, so CR, prebolus, immediate bolus percent, and e-carbs cannot be tested.
- (12/50) Glucose variability is high (CV 32.6 percent);
- ... 20 more varying points

Points made by a single run only: 57.

Phrases used by at least half the runs: “42 mg/dl” (40), “41 42 mg/dl” (36), “glp notes” (35), “time below range” (32), “dynamic isf” (31), “profile switch” (30), “max iob 25” (27), “12 u/h” (25).

Evidence cited: 19:00 (45), 5.1% (45), 42mg/dl (40), 25u (30), 01:00 (29), 12u/h (25), 32.6% (25), 17:00 (24), 14:00 (22), 03:00 (21), 16:00 (20), 90mg/dl (18).

step findings: data_quality

50 runs

Shared reasoning

- (46/50) CGM quality is good: 3995 valid entries, 98.7% coverage, about 5-minute cadence, 10 gaps, largest gap 65 minutes, no invalid or duplicate points.

Varying reasoning

- (26/50) Mid-period events include a 13 Aug profile-switch note, GLP-1 notes, site and sensor changes, and an 19 Aug occlusion.
- (20/50) The selected AAPS profile is complete and historically applicable.
- (14/50) Insulin treatments are abundant (655 insulin entries, 275.95 U) but carb entries are zero and every sampled loop status shows COB 0.
- (11/50) The selected aaps_profile is complete and profile_context states historical applicability is yes.
- (11/50) The authoritative aaps_profile is complete.
- (9/50) First and last calendar days are partial.
- ... 13 more varying points

Points made by a single run only: 29.

Phrases used by at least half the runs: “10 gaps” (46), “65 minutes” (45), “largest gap” (43), “minute cadence” (33), “3995 valid” (32), “10 gaps largest” (32), “aaps_profile is complete” (31), “profile switch” (30), “gaps largest gap” (29), “gap 65 minutes” (29), “largest gap 65” (29), “cadence 10 gaps” (28).

Evidence cited: 65min (46), 98.7% (43), 275.95u (9), 10h (6), 5min (5), 19.7u (2), 16.2u (2), 2u (1).

step findings: safety_overview

50 runs

Shared reasoning

- (36/50) Local TBR is highest at 01:00 (18.5 percent) and 19:00 (15.3 percent).

Varying reasoning

- (24/50) Extremes include 41-42 mg/dL and highs 286-293 mg/dL.
- (17/50) Time below range 5.1% is the binding safety constraint.
- (17/50) Mean glucose 121.6 mg/dL, TIR 86.0 percent, TAR 8.9 percent, TBR 5.1 percent, CV 32.6 percent.
- (17/50) Afternoon TAR peaks at 16:00 (32.1%).

- (15/50) Sampled IOB reached about 4.7-5.9 U.
- (12/50) Afternoon time above range peaks at 14:00-17:00.
- ... 26 more varying points

Points made by a single run only: 32.

Phrases used by at least half the runs: “42 mg/dl” (43), “41 42 mg/dl” (38), “tbr 5.1” (30).

Evidence cited: 19:00 (44), 5.1% (43), 42mg/dl (43), 18.5% (36), 15.3% (34), 01:00 (30), 32.6% (28), 03:00 (22), 16:00 (20), 5.9u (17), 23:00 (16), 32.1% (16).

step findings: basal_targets

50 runs

Shared reasoning

- (37/50) 04:00-08:00 is the most stable fasting-like window.
- (30/50) 00:00-03:00 sits near target 90 with high TBR.

Varying reasoning

- (19/50) The 16:00 basal cut to 0.5 U/h and the 19:00 rise to 0.7 U/h sit beside the 18:00-19:00 crash, so raising evening basal is unsupported.
- (17/50) The 09:00-16:00 rise and 16:00 TAR are more consistent with unannounced intake than with basal 0.7 then 0.5 U/h.
- (10/50) Scheduled basal is about 16.2 U/day and matches the profile name.
- (8/50) Overnight means sit near target with excess time below range from 00:00-03:00, then 05:00-08:00 time in range is 97-99%.
- (8/50) The 09:00 to 10:00 rise is meal-shaped after a stable dawn.
- (5/50) Afternoon highs occur while basal is already lowest at 16:00 (0.5 U/h) and coincide with likely unlogged intake and UAM, so they are not a basal hole.
- ... 23 more varying points

Points made by a single run only: 48.

Phrases used by at least half the runs: “0.5 u/h” (40), “00 basal” (39), “90 mg/dl” (38), “16 00 basal” (32), “target 90” (28).

Evidence cited: 16:00 (46), 08:00 (45), 0.5u/h (40), 90mg/dl (38), 19:00 (30), 04:00 (30), 03:00 (23), 05:00 (22), 00:00 (21), 0.8u/h (18), 99% (17), 100mg/dl (17).

step findings: profile_cr_isf_dia

50 runs

Shared reasoning

- (40/50) Flat ISF 56 mg/dL/U is not the dosed sensitivity while Dynamic ISF is on; loop variable_sens ranged about 28-124 mg/dL/U.

Varying reasoning

- (24/50) CR segments exist but were unused because carb entries and COB are zero.
- (13/50) CR segments cannot be tested because no carbs were logged.
- (11/50) DIA 10 h can leave afternoon insulin active at 19:00 and overnight.
- (10/50) CR segments exist but are unused because no carbs were logged.

- (9/50) CR, ISF, and DIA stay withheld.
- (7/50) Correction windows are not clean ISF tests.
- ... 18 more varying points

Points made by a single run only: 49.

Phrases used by at least half the runs: “dia 10” (48), “isf 56” (44), “cr segments” (44), “56 mg/dl/u” (38), “flat isf 56” (37), “dynamic isf” (35), “isf 56 mg/dl/u” (31), “correction windows” (27).

Evidence cited: 10h (50), 56mg/dl (38), 124mg/dl (14), 19:00 (7), 6h (6), 03:00 (5), 4h (4), 01:00 (4), 00:00 (3), 70% (2), 12.6g/u (2), 23:00 (2).

step findings: automation_smb_uam

50 runs

Varying reasoning

- (27/50) SMB and UAM are on, SMB is always enabled, and COB-based SMB is off, which matches zero announced carbs.
- (23/50) High temporary targets at 135 mg/dL should suppress SMB.
- (16/50) UAM is the meal engine, not duplicate COB counting.
- (9/50) Delivery is frequent at a 3-minute interval, and loop reasons show UAM boosts and SMB sizes larger than a naive 15-20 minutes of 0.5-0.8 U/h basal, including a 1.6 U UAM high boost.
- (7/50) SMB and UAM always-on with 3-minute interval, 15-minute SMB and 20-minute UAM basal minutes is internally consistent with a no-carb workflow.
- (5/50) Nominal 15/20 minute SMB caps look small on 0.5-0.8 U/h basal, but sampled reasons show UAM boosts, microboluses above the stated standard max, and at least one 1.6 U SMB with IOB already 4.73 U.
- ... 34 more varying points

Points made by a single run only: 72.

Phrases used by at least half the runs: “smb and uam” (45).

Evidence cited: 20min (31), 1.6u (23), 135mg/dl (15), 3min (14), 0.4u (11), 19:00 (7), 0.8u/h (7), 15min (6), 4u/h (6), 4.73u (5), 18:00 (4), 10h (4).

step findings: automation_sensitivity_limits

50 runs

Shared reasoning

- (36/50) Autosens is off; a sensitivity_ratio of 1 is a neutral adjustment, not proof of Autosens.
- (36/50) Max basal 12 U/h and max IOB 25 U are not binding.
- (32/50) Carb absorption settings are idle without COB.

Varying reasoning

- (27/50) Dynamic ISF is on at 70%.
- (10/50) Carb-absorption settings had no COB to act on.
- (7/50) Autosens is off and target raise or lower is off.
- (5/50) DynISF 70% is the active high-glucose aggressor. max_basal 12 U/h and max_iob 25 U do not bind observed delivery; enacted temps meet a lower maxSafeBasal near 4 U/h and peak sampled IOB is about 5.9 U.

- (5/50) Automation is compensating for missing meal announcements more than for a proven basal or CR error.
- (5/50) Automation is compensating for unlogged meals rather than proving a weak profile.
- ... 12 more varying points

Points made by a single run only: 55.

Phrases used by at least half the runs: “12 u/h” (50), “carb absorption” (50), “max basal 12” (36), “basal 12 u/h” (36), “max iob 25” (36), “dynamic isf” (35), “u/h and max” (26).

Evidence cited: 25u (50), 12u/h (50), 70% (48), 4u/h (37), 6u (19), 5.9u (11), 20u (8), 19.7u (4), 0.8u/h (3), 5.935u (2), 5min (2), 7h (2).

step findings: meal_bolus_strategy

50 runs

Shared reasoning

- (48/50) User-declared meal_strategies is an empty list and strategy_stable_for_period is null, so there is no declared prebolus or immediate-bolus percent.

Varying reasoning

- (14/50) Observed execution is correction boluses plus UAM/SMB only.
- (13/50) Recurring 08:00-10:00 and 14:00-17:00 rises with later 18:00-19:00 TBR match unannounced intake plus UAM/SMB, not a CR-based bolus plan.
- (12/50) Observed execution is unannounced intake handled by UAM/SMB, tiny corrections, and occasional high temporary targets of 135.
- (8/50) Adding a meal bolus on top of current UAM/SMB would raise evening and overnight hypo risk.
- (8/50) Observed execution matches that declaration: only correction boluses and UAM/SMB.
- (5/50) Temporary targets of 135 mg/dL are present but are not a numeric bolus percentage.
- ... 16 more varying points

Points made by a single run only: 51.

Phrases used by at least half the runs: “observed execution” (44), “user declared” (38), “declared meal_strategies” (38), “immediate bolus” (37), “empty list” (36), “immediate bolus percent” (34), “strategy_stable_for_period is null” (32), “user declared meal_strategies” (31), “list and strategy_stable_for_period” (29).

Evidence cited: 135mg/dl (8), 19:00 (7), 17:00 (7), 14:00 (6), 4h (6), 18:00 (5), 10:00 (4), 09:00 (4), 6h (3), 08:00 (3), 10h (2), 16:00 (2).

step findings: meal_ecarbs_absorption

50 runs

Varying reasoning

- (26/50) E-carbs are not a pump extended bolus.
- (16/50) No e-carb amount, start offset, or duration is declared or logged.
- (14/50) Extended carb entries are 0.
- (12/50) Carb-absorption model settings are idle.
- (11/50) No e-carb amount, start offset, or duration exists in the declaration or in telemetry.
- (9/50) Idle absorption settings cannot be judged.
- ... 23 more varying points

Points made by a single run only: 31.

Phrases used by at least half the runs: “extended bolus” (34), “pump extended” (29), “pump extended bolus” (28).

Evidence cited: 4h (4), 19:00 (3), 6h (3), 3h (2), 18:00 (2), 10h (2), 5h (1), 12:00 (1), 17:00 (1), 180min (1), 03:00 (1), 7h (1).

step findings: synthesis_plan

50 runs

Shared reasoning

- (37/50) The ranked explanation is afternoon unannounced absorption handled late by UAM, SMB, and Dynamic ISF, leaving IOB into 18:00-19:00 and some 00:00-03:00 lows.

Varying reasoning

- (8/50) Ranked causes: unannounced intake plus UAM lag for 10:00 and 14:00-17:00 highs; leftover IOB plus always-on SMB plus night target 90, with overnight basal still competing, for 23:00-03:00 and 19:00 lows;
- (8/50) Safe order is keep the profile, observe IOB-clear nights and log meals, then consider only one later lever if the pattern remains separated.
- (8/50) The conservative sequence is keep the current profile, observe meals and overnight IOB versus lows, then consider one later setting only after those are separated.
- (8/50) One conservative profile review is 00:00 target 90 to 100 mg/dL.
- (7/50) 16:00 basal 0.5 U/h is not a supported hole;
- (5/50) Max basal 12 and max IOB 25 were not the binding brakes.
- ... 31 more varying points

Points made by a single run only: 102.

Phrases used by at least half the runs: “target 90” (44), “00 basal” (39), “overnight target” (32), “16 00 basal” (31), “dynamic isf” (26), “overnight target 90” (26), “dia 10” (26).

Evidence cited: 16:00 (35), 19:00 (31), 10h (29), 00:00 (29), 90mg/dl (22), 03:00 (21), 0.6u/h (15), 0.5u/h (13), 18:00 (12), 100mg/dl (9), 23:00 (9), 17:00 (6).

meal strategy

50 runs

Varying reasoning

- (17/50) Declared meal_strategies is empty and Nightscout shows zero carbs, zero e-carbs, and correction-only response windows.
- (17/50) There is no declared prebolus, no declared immediate bolus percent, and no e-carbs.
- (11/50) There is no declared prebolus time, no declared immediate-bolus percent, and no e-carb plan.
- (11/50) User-declared meal_strategies is empty and strategy_stable_for_period is null.
- (8/50) The recurring 14:00-17:00 rise then 18:00-19:00 drop is therefore UAM/DynISF execution, not a CR or e-carb setting that can be tuned from this file.
- (6/50) Do not add e-carbs or invent a prebolus or immediate-bolus percent.
- ... 26 more varying points

Points made by a single run only: 82.

Phrases used by at least half the runs: “immediate bolus” (48), “bolus percent” (45), “immediate bolus percent” (40), “declared prebolus” (37), “meal_strategies is empty” (37), “declared meal_strategies” (26).

Evidence cited: 10h (24), 19:00 (16), 18:00 (10), 135mg/dl (10), 03:00 (5), 14:00 (4), 17:00 (4), 01:00 (3), 90mg/dl (3), 09:00 (2), 4h (2), 16:00 (2).