

Reasoning analysis — meta-llama/llama-4-maverick (49 runs, `real_llama4maverick_50`)

Each model rationale is split into sentences; sentences making the same point across runs are grouped. **Shared** = the point appears in at least 60% of the runs in that group; **varying** = it appears in some runs but not most. Counts are runs, not sentences. 'Evidence cited' lists the numbers and times the model quoted, with the number of runs quoting each.

decision: Profile / basal 00:00

keep — 30/49 runs

Varying reasoning

- (10/30) No strong evidence to change basal rate at 00:00, evidence from basal_targets, uncertainty due to meal and correction effects, dependencies on data_quality and safety_overview.
- (9/30) Current basal rate is generally appropriate.
- (5/30) Dependencies: basal_targets.
- (4/30) Evidence: Basal rate settings and glucose patterns.
- (3/30) Basal rate seems reasonable.
- (3/30) No strong evidence for change at this time.
- ... 5 more varying points

Points made by a single run only: 7.

Phrases used by at least half the runs: "basal rate" (23).

Evidence cited: 00:00 (4), 0.6u/h (1).

change — 19/49 runs

Varying reasoning

- (5/19) Evidence from glucose patterns, uncertainty due to limited meal data, dependencies on basal_targets and safety_overview.
- (5/19) Glucose variability, evidence from basal_targets stage.
- (4/19) Basal rate analysis.
- (3/19) Evidence from glucose patterns
- (2/19) Uncertainty: Medium.

Points made by a single run only: 6.

Phrases used by at least half the runs: "glucose patterns" (11).

decision: Profile / basal 05:00

keep — 48/49 runs

Varying reasoning

- (24/48) No strong evidence to change.
- (9/48) No strong evidence to change the current basal rate.

- (8/48) Current basal rate is generally appropriate.
- (6/48) Evidence from glucose data, uncertainty due to meal effects, dependencies on data_quality and safety_overview.
- (5/48) The uncertainty is low.
- (4/48) Evidence: Basal rate settings and glucose patterns.
- ... 4 more varying points

Points made by a single run only: 8.

Phrases used by at least half the runs: “basal rate” (25).

Evidence cited: 05:00 (4), 0.7u/h (1).

decision: Profile / basal 06:00

keep — 47/49 runs

Varying reasoning

- (20/47) No strong evidence to change.
- (9/47) No strong evidence to change the current basal rate.
- (8/47) Current basal rate is generally appropriate.
- (6/47) Evidence from glucose data, uncertainty due to meal effects, dependencies on data_quality and safety_overview.
- (6/47) Dependencies: basal_targets.
- (4/47) Evidence: Basal rate settings and glucose patterns.
- ... 5 more varying points

Points made by a single run only: 6.

Phrases used by at least half the runs: “basal rate” (25).

Evidence cited: 06:00 (4), 0.8u/h (1).

change — 1/49 runs

Points made by a single run only: 1.

Phrases used by at least half the runs: “glucose patterns” (1).

decision: Profile / basal 10:00

keep — 48/49 runs

Varying reasoning

- (21/48) No strong evidence to change.
- (9/48) No strong evidence to change the current basal rate.
- (8/48) Current basal rate is generally appropriate.
- (6/48) Evidence from glucose data, uncertainty due to meal effects, dependencies on data_quality and safety_overview.

- (6/48) Dependencies: basal_targets.
- (4/48) Evidence: Basal rate settings and glucose patterns.
- ... 5 more varying points

Points made by a single run only: 6.

Phrases used by at least half the runs: “basal rate” (25).

Evidence cited: 10:00 (4), 0.7u/h (1).

decision: Profile / basal 16:00

keep — 46/49 runs

Varying reasoning

- (20/46) No strong evidence to change.
- (9/46) No strong evidence to change the current basal rate.
- (8/46) Current basal rate is generally appropriate.
- (6/46) Dependencies: basal_targets.
- (5/46) Evidence from glucose data, uncertainty due to meal effects, dependencies on data_quality and safety_overview.
- (4/46) Evidence: Basal rate settings and glucose patterns.
- ... 4 more varying points

Points made by a single run only: 8.

Phrases used by at least half the runs: “basal rate” (25).

Evidence cited: 16:00 (4), 0.5u/h (1).

change — 2/49 runs

Points made by a single run only: 2.

Phrases used by at least half the runs: “glucose patterns” (1), “frequent correction boluses” (1).

decision: Profile / basal 18:00

keep — 48/49 runs

Varying reasoning

- (21/48) No strong evidence to change.
- (9/48) No strong evidence to change the current basal rate.
- (8/48) Current basal rate is generally appropriate.
- (6/48) Evidence from glucose data, uncertainty due to meal effects, dependencies on data_quality and safety_overview.
- (6/48) Dependencies: basal_targets.
- (4/48) Evidence: Basal rate settings and glucose patterns.

- ... 5 more varying points

Points made by a single run only: 6.

Phrases used by at least half the runs: “basal rate” (25).

Evidence cited: 18:00 (4), 0.6u/h (1).

decision: Profile / basal 19:00

keep — 48/49 runs

Varying reasoning

- (21/48) No strong evidence to change.
- (9/48) No strong evidence to change the current basal rate.
- (8/48) Current basal rate is generally appropriate.
- (6/48) Dependencies: basal_targets.
- (5/48) Evidence from glucose data, uncertainty due to meal effects, dependencies on data_quality and safety_overview.
- (4/48) Evidence: Basal rate settings and glucose patterns.
- ... 5 more varying points

Points made by a single run only: 7.

Phrases used by at least half the runs: “basal rate” (25).

Evidence cited: 19:00 (4), 0.7u/h (1).

decision: Profile / CR 00:00

keep — 48/49 runs

Shared reasoning

- (31/48) No strong evidence to change.

Varying reasoning

- (10/48) Current CR is generally appropriate.
- (3/48) CR seems appropriate.
- (3/48) Uncertainty: Low.
- (3/48) Uncertainty: Medium.
- (2/48) Uncertainty due to lack of meal data.
- (2/48) No strong evidence for change, depends on meal_bolus_strategy.
- ... 4 more varying points

Points made by a single run only: 12.

Evidence cited: 00:00 (4), 13g/u (1).

change — 1/49 runs

Points made by a single run only: 1.

Phrases used by at least half the runs: “effects suggest potential” (1), “meal effects” (1), “adjustment in cr” (1), “potential adjustment” (1).

decision: Profile / CR 04:00

keep — 49/49 runs

Shared reasoning

- (32/49) No strong evidence to change.

Varying reasoning

- (10/49) Current CR is generally appropriate.
- (3/49) CR seems appropriate.
- (3/49) Uncertainty: Low.
- (3/49) Uncertainty: Medium.
- (2/49) Uncertainty due to lack of meal data.
- (2/49) No strong evidence for change, depends on meal_bolus_strategy.
- ... 4 more varying points

Points made by a single run only: 12.

Evidence cited: 04:00 (4), 11.1g/u (1).

decision: Profile / CR 08:00

keep — 49/49 runs

Shared reasoning

- (32/49) No strong evidence to change.

Varying reasoning

- (10/49) Current CR is generally appropriate.
- (3/49) CR seems appropriate.
- (3/49) Uncertainty: Low.
- (3/49) Uncertainty: Medium.
- (2/49) Uncertainty due to lack of meal data.
- (2/49) No strong evidence for change, depends on meal_bolus_strategy.
- ... 4 more varying points

Points made by a single run only: 12.

Evidence cited: 08:00 (4), 13.2g/u (1).

decision: Profile / CR 16:00

keep — 49/49 runs

Shared reasoning

- (32/49) No strong evidence to change.

Varying reasoning

- (10/49) Current CR is generally appropriate.
- (3/49) CR seems appropriate.
- (3/49) Uncertainty: Low.
- (3/49) Uncertainty: Medium.
- (2/49) Uncertainty due to lack of meal data.
- (2/49) No strong evidence for change, depends on meal_bolus_strategy.
- ... 4 more varying points

Points made by a single run only: 12.

Evidence cited: 16:00 (4), 10.5g/u (1).

decision: Profile / CR 20:00

keep — 49/49 runs

Shared reasoning

- (32/49) No strong evidence to change.

Varying reasoning

- (10/49) Current CR is generally appropriate.
- (3/49) CR seems appropriate.
- (3/49) Uncertainty: Low.
- (3/49) Uncertainty: Medium.
- (2/49) Uncertainty due to lack of meal data.
- (2/49) No strong evidence for change, depends on meal_bolus_strategy.
- ... 4 more varying points

Points made by a single run only: 12.

Evidence cited: 20:00 (4), 12.6g/u (1).

decision: Profile / ISF 00:00

keep — 45/49 runs

Varying reasoning

- (20/45) No strong evidence to change.
- (10/45) Dependencies on profile_cr_isf_dia.
- (8/45) Current ISF is generally appropriate.
- (5/45) Evidence from glucose data, uncertainty due to correction bolus effects, dependencies on profile_cr_isf_dia.
- (4/45) Uncertainty: Medium.
- (3/45) Evidence: ISF settings and glucose patterns.
- ... 5 more varying points

Points made by a single run only: 8.

Evidence cited: 00:00 (2), 56mg/dl (1).

change — 4/49 runs

Shared reasoning

- (3/4) Evidence from glucose variability and correction bolus analysis.

Points made by a single run only: 4.

Phrases used by at least half the runs: “glucose variability” (3), “correction bolus” (2), “variability and correction” (2).

decision: Profile / target 00:00

keep — 49/49 runs

Varying reasoning

- (20/49) No strong evidence to change.
- (8/49) Current target is generally appropriate.
- (8/49) No strong evidence to change the current target.
- (8/49) Dependencies: basal_targets.
- (5/49) Uncertainty due to glucose variability.
- (5/49) Uncertainty: Low.
- ... 7 more varying points

Points made by a single run only: 6.

Evidence cited: 00:00 (4), 90mg/dl (1).

decision: Profile / target 08:00

keep — 49/49 runs

Varying reasoning

- (20/49) No strong evidence to change.

- (8/49) Current target is generally appropriate.
- (8/49) No strong evidence to change the current target.
- (8/49) Dependencies: basal_targets.
- (5/49) Uncertainty due to glucose variability.
- (5/49) Uncertainty: Low.
- ... 7 more varying points

Points made by a single run only: 6.

Evidence cited: 08:00 (4), 100mg/dl (1).

decision: Profile / target 23:00

keep — 49/49 runs

Varying reasoning

- (20/49) No strong evidence to change.
- (8/49) Current target is generally appropriate.
- (8/49) No strong evidence to change the current target.
- (8/49) Dependencies: basal_targets.
- (5/49) Uncertainty due to glucose variability.
- (5/49) Uncertainty: Low.
- ... 7 more varying points

Points made by a single run only: 6.

Evidence cited: 23:00 (4), 90mg/dl (1).

decision: Profile / DIA

keep — 49/49 runs

Shared reasoning

- (30/49) No strong evidence to change.

Varying reasoning

- (10/49) Current DIA is generally appropriate.
- (5/49) Evidence from insulin action profiles, dependencies on profile_cr_isf_dia.
- (4/49) Uncertainty: Low.
- (4/49) Uncertainty: Medium.
- (3/49) DIA seems appropriate.
- (2/49) No strong evidence for change, depends on profile_cr_isf_dia.
- ... 4 more varying points

Points made by a single run only: 8.

Evidence cited: 10h (1).

decision: AAPS / SMB and UAM

keep — 49/49 runs

Varying reasoning

- (19/49) SMB/UAM settings are generally appropriate.
- (14/49) No strong evidence for change, dependencies on automation_smb_uam.
- (8/49) SMB and UAM are appropriately enabled.
- (5/49) Uncertainty: Low.
- (4/49) Dependencies on safety_overview, basal_targets, and profile_cr_isf_dia.
- (4/49) SMB and UAM are effectively managing glucose variability.
- ... 5 more varying points

Points made by a single run only: 8.

decision: AAPS / SMB activation conditions

keep — 49/49 runs

Varying reasoning

- (25/49) SMB activation conditions are generally appropriate.
- (15/49) Dependencies: automation_smb_uam.
- (6/49) Current SMB activation conditions are reasonable.
- (6/49) Uncertainty: Low.
- (4/49) SMB activation conditions are appropriately set.
- (3/49) Dependencies on safety_overview, basal_targets, and profile_cr_isf_dia.
- ... 2 more varying points

Points made by a single run only: 10.

Phrases used by at least half the runs: “activation conditions” (34), “smb activation” (33), “smb activation conditions” (32).

decision: AAPS / SMB strength and frequency

keep — 49/49 runs

Varying reasoning

- (26/49) SMB delivery settings are generally appropriate.
- (15/49) Dependencies: automation_smb_uam.
- (9/49) SMB delivery settings are reasonable.
- (6/49) Uncertainty: Low.
- (4/49) SMB delivery settings are appropriately configured, evidence from automation_smb_uam, uncertainty due to glucose variability, dependencies on safety_overview, basal_targets, and profile_cr_isf_dia.

- (2/49) SMB delivery settings are appropriately configured.
- ... 1 more varying points

Points made by a single run only: 11.

Phrases used by at least half the runs: “smb delivery” (26).

Evidence cited: 3min (1).

decision: AAPS / max basal and max IOB limits

keep — 26/49 runs

Varying reasoning

- (7/26) Current max basal and max IOB limits are generally appropriate.
- (6/26) Safety limits are set high but not changed.
- (4/26) No strong evidence to change.
- (3/26) No strong evidence to change, but high max IOB may pose risk
- (3/26) Safety limits are high but reasonable.
- (3/26) Dependencies: automation_sensitivity_limits.
- ... 1 more varying points

Points made by a single run only: 4.

change — 23/49 runs

Shared reasoning

- (16/23) High max IOB poses a risk of excessive insulin dosing.

Varying reasoning

- (6/23) Uncertainty: Medium.
- (5/23) Dependencies: automation_sensitivity_limits.
- (2/23) High max IOB poses a risk, evidence from automation_sensitivity_limits analysis, uncertainty due to limited data, dependencies on safety_overview, basal_targets, and profile_cr_isf_dia.
- (2/23) High limits pose a risk; reduce to safer values.

Points made by a single run only: 12.

Phrases used by at least half the runs: “max iob” (22), “high max iob” (20).

Evidence cited: 15u (2), 25u (2), 20u (1).

decision: AAPS / Autosens and Dynamic ISF

keep — 48/49 runs

Varying reasoning

- (16/48) Autosens and Dynamic ISF settings are generally appropriate.
- (15/48) Evidence: Dynamic ISF is enabled.
- (15/48) Dependencies: automation_sensitivity_limits.
- (6/48) Uncertainty: Medium.
- (4/48) Autosens is disabled and Dynamic ISF is enabled with appropriate settings, evidence from automation_sensitivity_limits, uncertainty due to glucose variability, dependencies on safety_overview, basal_targets, profile_cr_isf_dia, and automation_smb_uam.
- (4/48) Current settings are appropriate.
- ... 3 more varying points

Points made by a single run only: 7.

Phrases used by at least half the runs: “dynamic isf” (33).

Evidence cited: 70% (1).

verify — 1/49 runs

Points made by a single run only: 1.

Phrases used by at least half the runs: “autosens is disabled” (1), “disabled its impact” (1).

decision: AAPS / carbohydrate absorption model

keep — 49/49 runs

Varying reasoning

- (26/49) Carb absorption model settings are generally appropriate.
- (16/49) No strong evidence for change, dependencies on meal_ecarbs_absorption.
- (5/49) Uncertainty: Low.
- (4/49) Carb absorption settings are appropriately configured, evidence from meal_ecarbs_absorption, uncertainty due to lack of meal data, dependencies on meal_bolus_strategy.
- (4/49) No meal carb entries to assess carbohydrate absorption.
- (4/49) Carb absorption settings are reasonable.
- ... 1 more varying points

Points made by a single run only: 12.

Phrases used by at least half the runs: “carb absorption” (28).

Evidence cited: 5min (1), 8mg/dl (1).

profile recommendation

basal — 43/49 runs

Shared reasoning

- (26/43) Adjust basal rate at 00:00 to 0.65 U/h.

Varying reasoning

- (11/43) Keep basal rates with close monitoring.
- (5/43) Glucose variability suggests potential basal rate adjustments.
- (5/43) Basal rates seem reasonable but need further analysis with meal data.
- (3/43) Keep basal rates as is and monitor glucose control No strong evidence to change basal rates.
- (3/43) Keep basal rates with next check on overnight glucose patterns.
- (3/43) To improve glucose control based on observed variability.

Points made by a single run only: 12.

Phrases used by at least half the runs: “basal rate” (27), “adjust basal” (26), “rate at 00” (23), “adjust basal rate” (23).

Evidence cited: 00:00 (24), 0.65u/h (16), 0.55u/h (6), 0.6u/h (4), 16:00 (1).

isf — 4/49 runs

Shared reasoning

- (3/4) Adjust ISF to 50 mg/dL/U.

Points made by a single run only: 4.

Phrases used by at least half the runs: “adjust isf” (3), “isf to 50” (3), “50 mg/dl/u” (3), “glucose variability” (3).

Evidence cited: 50mg/dl (3), 56mg/dl (1).

cr — 1/49 runs

Points made by a single run only: 2.

Phrases used by at least half the runs: “keep cr” (1), “need meal” (1), “verify with meal” (1), “cr settings seem” (1), “appropriate but need” (1), “seem generally appropriate” (1).

aaps — 1/49 runs

Points made by a single run only: 2.

Phrases used by at least half the runs: “max iob profile” (1), “appropriate but max” (1), “max iob needs” (1), “iob needs adjustment” (1), “adjust max iob” (1), “generally appropriate” (1).

primary recommendation

basal — 26/49 runs

Shared reasoning

- (16/26) Review basal rates.

Varying reasoning

- (12/26) Adjust basal rates.
- (7/26) Adjust basal rates to improve glucose control.

- (2/26) Basal rates need adjustment based on observed glucose patterns.
- (2/26) Basal rates vary throughout the day, and adjustments may be needed based on glucose patterns.
- (2/26) Adjust basal rate profile for better overnight control.
- (2/26) Consider adjusting basal rates to reduce glucose variability.
- ... 2 more varying points

Points made by a single run only: 4.

Phrases used by at least half the runs: “basal rates” (26), “adjust basal” (19), “adjust basal rates” (16), “glucose patterns” (13).

AAPS safety limits — 7/49 runs

Shared reasoning

- (5/7) Adjust max IOB limit.

Varying reasoning

- (4/7) Reduce max IOB from 25 U to 15 U to prevent excessive insulin dosing.
- (2/7) Current max IOB is set to 25 U, which is high and poses a risk.

Points made by a single run only: 2.

Phrases used by at least half the runs: “max iob” (7), “adjust max iob” (6), “reduce max iob” (5), “excessive insulin” (4).

Evidence cited: 25u (4), 15u (2).

profile — 5/49 runs

Shared reasoning

- (3/5) Review and adjust ISF.

Points made by a single run only: 6.

Evidence cited: 56mg/dl (1).

AAPS — 4/49 runs

Shared reasoning

- (4/4) Reduce max IOB to prevent excessive insulin dosing.

Phrases used by at least half the runs: “excessive insulin” (4), “reduce max iob” (4), “excessive insulin dosing” (3), “max iob reduce” (3), “adjust max iob” (3), “prevent excessive insulin” (3), “iob reduce max” (3), “iob to prevent” (2).

meal strategy — 3/49 runs

Shared reasoning

- (3/3) Start logging meal data..

Points made by a single run only: 3.

Phrases used by at least half the runs: “start logging meal” (3), “meal data logging” (2).

Meal Logging — 1/49 runs

Points made by a single run only: 1.

Phrases used by at least half the runs: “meals will improve” (1), “improve understanding” (1), “meals logging meals” (1), “understanding of glucose” (1), “start logging meals” (1), “logging meals logging” (1), “glucose patterns” (1).

meal logging — 1/49 runs

Points made by a single run only: 1.

Phrases used by at least half the runs: “improving meal bolus” (1), “meal bolus strategies” (1), “meal data logging” (1), “start logging meal” (1), “crucial for assessing” (1), “assessing and improving” (1).

Meal Strategy — 1/49 runs

Points made by a single run only: 1.

Phrases used by at least half the runs: “strategy could improve” (1), “meal bolus strategy” (1), “bolus strategy recording” (1), “strategy recording carb” (1), “recording carb intake” (1), “consistent meal bolus” (1), “implement meal bolus” (1), “improve glucose control” (1).

summary

49 runs

Varying reasoning

- (17/49) The current AAPS profile and settings are generally appropriate but require adjustments for better glucose control.
- (16/49) The user’s glucose management relies heavily on correction boluses and SMB/UAM.
- (9/49) The glucose data shows a generally good time in range (TIR) but with instances of hyperglycemia and hypoglycemia.
- (9/49) The glucose data shows variability with instances of hypo and hyperglycemia.
- (7/49) The analysis indicates generally good glucose control with some variability.
- (6/49) The user’s glucose data shows variability with both hypo and hyperglycemic events.
- ... 15 more varying points

Points made by a single run only: 10.

Phrases used by at least half the runs: “glucose management” (25).

Evidence cited: 86% (1), 86.0% (1).

issues

49 runs

Varying reasoning

- (29/49) High max IOB setting.
- (18/49) Lack of meal data.
- (6/49) Variability in glucose levels.
- (5/49) Glucose variability.
- (4/49) Glucose variability with hyperglycemia and hypoglycemia events.
- (3/49) No explicit meal bolus strategy.
- ... 2 more varying points

Points made by a single run only: 4.

Phrases used by at least half the runs: “max iob” (29), “high max” (29), “high max iob” (27).

Evidence cited: 25u (3), 12u/h (1).

step findings: data_quality

49 runs

Varying reasoning

- (24/49) Good CGM coverage with minor gaps.
- (18/49) Good CGM data quality with 98.7% coverage.
- (7/49) CGM data coverage is 98.7% with 10 gaps, the largest being 65 minutes.

Points made by a single run only: 1.

Phrases used by at least half the runs: “cgm coverage” (31), “good cgm” (31).

Evidence cited: 98.7% (27), 65min (7).

step findings: safety_overview

49 runs

Varying reasoning

- (23/49) TIR is 86.0%, TAR is 8.9%, and TBR is 5.1%.
- (10/49) Generally safe with some hyperglycemia and hypoglycemia events.
- (9/49) Generally safe with instances of hypo and hyperglycemia.
- (8/49) Significant time in range but with hypo and hyperglycemic events.
- (5/49) Average glucose is 121.6 mg/dL.
- (2/49) Glucose is generally within range but with variability.

Evidence cited: 8.9% (22), 5.1% (22), 86.0% (15), 86% (14), 121.6mg/dl (8).

step findings: basal_targets

49 runs

Varying reasoning

- (18/49) Basal rates and targets seem generally appropriate.
- (14/49) Basal rates and targets vary throughout the day.
- (9/49) Basal rates and target glucose levels vary throughout the day.
- (6/49) Basal rates and targets are generally stable.

Points made by a single run only: 4.

Phrases used by at least half the runs: “basal rates” (49), “rates and targets” (40).

step findings: profile_cr_isf_dia

49 runs

Varying reasoning

- (16/49) CR, ISF, and DIA settings are generally appropriate.
- (9/49) DIA is set to 10 hours.
- (6/49) CR and ISF settings need review in context of meal and correction effects.
- (6/49) CR and ISF settings need evaluation against meal and correction bolus responses.
- (3/49) CR, ISF, and DIA settings are in place.
- (3/49) CR, ISF, and DIA settings are within reasonable ranges.
- ... 4 more varying points

Points made by a single run only: 7.

Phrases used by at least half the runs: “cr and isf” (26).

Evidence cited: 10h (9), 56mg/dl (1), 13.2g/u (1).

step findings: automation_smb_uam

49 runs

Varying reasoning

- (27/49) SMB and UAM are enabled.
- (7/49) SMB and UAM are enabled with aggressive settings.
- (6/49) SMB/UAM settings are generally appropriate.
- (3/49) SMB/UAM is enabled and active.
- (2/49) SMB/UAM are enabled with high safety limits and frequent SMB delivery.

Points made by a single run only: 5.

Phrases used by at least half the runs: “smb and uam” (33), “uam are enabled” (30).

step findings: automation_sensitivity_limits

49 runs

Varying reasoning

- (28/49) Autosens is disabled, but Dynamic ISF is enabled.
- (16/49) Dynamic ISF is enabled, and safety limits are set high.
- (15/49) Max basal and max IOB limits are high.
- (2/49) Dynamic ISF is enabled, which may be compensating for some variability.

Points made by a single run only: 5.

Phrases used by at least half the runs: “dynamic isf” (46), “isf is enabled” (37).

Evidence cited: 70% (7), 25u (6), 12u/h (2).

step findings: meal_bolus_strategy

49 runs

Varying reasoning

- (26/49) No explicit meal bolus strategy is declared.
- (12/49) Meal strategy relies on correction boluses and SMB/UAM.
- (5/49) Lack of carb entries and explicit meal strategy declarations.
- (2/49) The user relies on SMB/UAM without explicit carb counting.
- (2/49) Lack of meal data makes it hard to assess meal bolus strategy.
- (2/49) No explicit meal logging or bolusing is observed.

Points made by a single run only: 5.

Phrases used by at least half the runs: “meal bolus” (25).

step findings: meal_ecarbs_absorption

49 runs

Varying reasoning

- (14/49) No e-carb data available.
- (11/49) No e-carb entries are used.
- (4/49) No carb entries or e-carbs were recorded.
- (3/49) No carb entries are logged.
- (3/49) No e-carb usage is declared or observed.
- (3/49) No e-carb or extended bolus data is available.
- ... 3 more varying points

Points made by a single run only: 4.

Phrases used by at least half the runs: “carb entries” (20).

step findings: synthesis_plan

49 runs

Varying reasoning

- (18/49) Primary recommendation is to adjust basal rates based on observed glucose patterns.
- (8/49) Primary recommendation is to adjust max IOB to a safer limit.
- (5/49) The primary recommendation is to adjust the basal rates for better glucose control.
- (4/49) The primary recommendation is to review basal rates.
- (3/49) Primary recommendation is to start logging meal data.
- (2/49) The primary recommendation is to adjust the basal rate profile for better overnight control.
- ... 3 more varying points

Points made by a single run only: 3.

Phrases used by at least half the runs: “primary recommendation” (40).

meal strategy

49 runs

Varying reasoning

- (18/49) Rely on SMB/UAM for glucose management.
- (11/49) No explicit meal bolus strategy;
- (7/49) Rely on correction boluses and SMB/UAM.
- (7/49) Lack of explicit meal strategy and carb entries.
- (6/49) Lack of meal data.
- (4/49) Continue using correction boluses and temp targets as needed.
- ... 16 more varying points

Points made by a single run only: 14.

Phrases used by at least half the runs: “rely on smb/uam” (29).