Chapter 6. The Gamma and Zeta Functions

Analytic continuation and functional equations of Γ and ζ

Contents

1 The gamma function **2**

1.1 Analytic continuation 2

1.2 Further properties of Γ 3

2 The zeta function **5**

2.1 Functional equation and analytic continuation 5

3 Selected exercises **8**

Overview

$1/\Gamma(s)$ is essentially the simplest entire function with zeros exactly at $s = 0, -1, -2, \ldots$. The zeta function enters number theory through Euler's identity

$$\prod_p \frac{1}{1-p^{-s}} = \sum_{n=1}^{\infty} \frac{1}{n^s} = \zeta(s)$$

over all primes. ζ is analytic for $\mathrm{Re}(s) > 1$; Riemann tied the primes to its meromorphic continuation to $\mathbb{C}$. We prove this and the functional equation (symmetry about $\mathrm{Re}(s) = 1/2$, via the theta function), and estimate ζ near $\mathrm{Re}(s) = 1$ for the prime number theorem (Chapter 7). The variable is s.

1 The gamma function

Motivation. Continuation, product formula and $\Gamma(s)\Gamma(1-s) = \pi/\sin \pi s$ are the tools needed for ζ.

For $s > 0$ the **gamma function** is

$$\Gamma(s) = \int_0^{\infty} e^{-t} t^{s-1}\, dt. \tag{1}$$

The integral converges: t^{s-1} is integrable near 0, and e^{-t} decays at infinity.

Proposition 1.1

The gamma function extends to an analytic function in the half-plane $\mathrm{Re}(s) > 0$, and is still given there by the integral formula (1).

Proof. It suffices to work in strips $S_{\delta,M} = \{\delta < \mathrm{Re}(s) < M\}$. With $\sigma = \mathrm{Re}(s)$, $|e^{-t}t^{s-1}| = e^{-t}t^{\sigma-1}$, so $\Gamma(s) = \lim_{\epsilon\to 0} F_\epsilon(s)$, $F_\epsilon(s) = \int_\epsilon^{1/\epsilon} e^{-t}t^{s-1}\, dt$, holomorphic by Theorem 5.4. By Theorem 5.2 it suffices that $F_\epsilon \to \Gamma$ uniformly on the strip. Now

$$|\Gamma(s) - F_\epsilon(s)| \le \int_0^{\epsilon} e^{-t}t^{\sigma-1}\, dt + \int_{1/\epsilon}^{\infty} e^{-t}t^{\sigma-1}\, dt.$$

The first integral is $\le \epsilon^\delta/\delta$ for $0 < \epsilon < 1$, and the second is $\le \int_{1/\epsilon}^{\infty} e^{-t}t^{M-1}\, dt \le C\int_{1/\epsilon}^{\infty} e^{-t/2}\, dt \to 0$, uniformly in the strip. ∎

1.1 Analytic continuation

Though the integral diverges elsewhere, a meromorphic function on $\mathbb{C}$ equals Γ for $\mathrm{Re}(s) > 0$: its **analytic continuation**, still Γ (unique, as the complement of the poles is connected).

Lemma 1.2

If $\mathrm{Re}(s) > 0$, then

$$\Gamma(s+1) = s\Gamma(s). \tag{2}$$

As a consequence $\Gamma(n+1) = n!$ for $n = 0, 1, 2, \ldots$.

Proof. Integrating by parts,

$$\int_\epsilon^{1/\epsilon} \frac{d}{dt}(e^{-t}t^s)\,dt = -\int_\epsilon^{1/\epsilon} e^{-t}t^s\,dt + s\int_\epsilon^{1/\epsilon} e^{-t}t^{s-1}\,dt,$$

and letting $\epsilon \to 0$ the left side vanishes, since $e^{-t}t^s \to 0$ as $t \to 0$ or $t \to \infty$. Also $\Gamma(1) = \int_0^\infty e^{-t}\,dt = 1$; apply (2) successively. ∎

Theorem 1.3

Γ continues to a meromorphic function on $\mathbb{C}$ whose only singularities are simple poles at $s = 0, -1, \ldots$, with $\operatorname{res}_{s=-n}\Gamma = (-1)^n/n!$.

Proof. It suffices to extend Γ to each half-plane $\operatorname{Re}(s) > -m$, $m \geq 1$. For $\operatorname{Re}(s) > -m$ define

$$F_m(s) = \frac{\Gamma(s+m)}{(s+m-1)(s+m-2)\cdots s}.$$

F_m is meromorphic in $\operatorname{Re}(s) > -m$ with simple poles at $s = 0, -1, \ldots, -m+1$ and residues

$$\operatorname{res}_{s=-n} F_m(s) = \frac{\Gamma(-n+m)}{(m-1-n)!(-1)(-2)\cdots(-n)} = \frac{(m-n-1)!}{(m-1-n)!(-1)(-2)\cdots(-n)} = \frac{(-1)^n}{n!}.$$

By Lemma 1.2, $F_m = \Gamma$ for $\operatorname{Re}(s) > 0$, so by uniqueness $F_m = F_k$ $(1 \leq k \leq m)$ on the domain of F_k; this gives the continuation. ∎

Remark

By analytic continuation, $\Gamma(s+1) = s\Gamma(s)$ holds whenever $s \neq 0, -1, -2, \ldots$. At $s = -n$, $n \geq 1$, both sides are infinite and $\operatorname{res}_{s=-n}\Gamma(s+1) = -n\operatorname{res}_{s=-n}\Gamma(s)$. At $s = 0$, $\Gamma(1) = \lim_{s\to 0} s\Gamma(s)$.

An alternate proof. For $\operatorname{Re}(s) > 0$ split $\Gamma(s) = \int_0^1 e^{-t}t^{s-1}\,dt + \int_1^\infty e^{-t}t^{s-1}\,dt$. The second integral is entire; expanding e^{-t} and integrating termwise, $\int_0^1 e^{-t}t^{s-1}\,dt = \sum_{n=0}^\infty \frac{(-1)^n}{n!(n+s)}$, so

$$\Gamma(s) = \sum_{n=0}^\infty \frac{(-1)^n}{n!(n+s)} + \int_1^\infty e^{-t}t^{s-1}\,dt \qquad \text{for } \operatorname{Re}(s) > 0. \tag{3}$$

The series is meromorphic on $\mathbb{C}$ with poles at $-n$, residues $(-1)^n/n!$: for $|s| < R$ split it at $N > 2R$; the finite part has the right poles, and for $n > N$, $\left|\frac{(-1)^n}{n!(n+s)}\right| \leq \frac{1}{n!R}$, so the tail is holomorphic. Hence (3) holds on $\mathbb{C}$.

1.2 Further properties of Γ

Theorem 1.4

For all $s \in \mathbb{C}$,

$$\Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin \pi s}. \tag{4}$$

$\Gamma(s)\Gamma(1-s)$ and $\pi/\sin \pi s$ are meromorphic with simple poles at the integers, so by analytic continuation it suffices to prove (4) for $0 < s < 1$.

Lemma 1.5

For $0 < a < 1$, $\displaystyle\int_0^\infty \frac{v^{a-1}}{1+v}\,dv = \frac{\pi}{\sin \pi a}$.

Proof. With $v = e^x$, $\int_0^\infty \frac{v^{a-1}}{1+v}\,dv = \int_{-\infty}^\infty \frac{e^{ax}}{1+e^x}\,dx$, which equals $\pi/\sin \pi a$ by Example 2. ∎

Proof of Theorem 1.4. For $0 < s < 1$ and $t > 0$, the change of variables $u = vt$ gives $\Gamma(1-s) = \int_0^\infty e^{-u}u^{-s}\,du = t\int_0^\infty e^{-vt}(vt)^{-s}\,dv$. Hence

$$\begin{aligned}\Gamma(1-s)\Gamma(s) &= \int_0^\infty e^{-t}t^{s-1}\Gamma(1-s)\,dt = \int_0^\infty e^{-t}t^{s-1}\Big(t\int_0^\infty e^{-vt}(vt)^{-s}\,dv\Big)dt \\ &= \int_0^\infty\int_0^\infty e^{-t[1+v]}v^{-s}\,dt\,dv = \int_0^\infty \frac{v^{-s}}{1+v}\,dv = \frac{\pi}{\sin\pi(1-s)} = \frac{\pi}{\sin \pi s}.\end{aligned}$$

∎

With $s = 1/2$, since $\Gamma(s) > 0$ for $s > 0$, $\Gamma(1/2) = \sqrt{\pi}$.

Theorem 1.6

The function Γ has the following properties:
(i) $1/\Gamma(s)$ is an entire function of s with simple zeros at $s = 0, -1, -2, \ldots$ and it vanishes nowhere else.
(ii) $1/\Gamma(s)$ has growth $\left|\frac{1}{\Gamma(s)}\right| \le c_1 e^{c_2|s|\log|s|}$. Therefore, $1/\Gamma$ is of order 1 in the sense that for every $\epsilon > 0$, there exists a bound $c(\epsilon)$ so that $\left|\frac{1}{\Gamma(s)}\right| \le c(\epsilon)e^{c_2|s|^{1+\epsilon}}$.

Proof. By Theorem 1.4,

$$\frac{1}{\Gamma(s)} = \Gamma(1-s)\frac{\sin \pi s}{\pi}, \tag{5}$$

so the simple poles of $\Gamma(1-s)$ at $s = 1, 2, \ldots$ are cancelled by zeros of $\sin \pi s$, and $1/\Gamma$ is entire with simple zeros at $0, -1, -2, \ldots$.

For the estimate, first $\int_1^\infty e^{-t}t^\sigma\,dt \le e^{(\sigma+1)\log(\sigma+1)}$ for $\sigma > 0$: choose n with $\sigma \le n \le \sigma + 1$; then $\int_1^\infty e^{-t}t^\sigma\,dt \le \int_0^\infty e^{-t}t^n\,dt = n! \le n^n = e^{n\log n} \le e^{(\sigma+1)\log(\sigma+1)}$. Since (3) holds on $\mathbb{C}$, by (5)

$$\frac{1}{\Gamma(s)} = \Big(\sum_{n=0}^\infty \frac{(-1)^n}{n!(n+1-s)}\Big)\frac{\sin \pi s}{\pi} + \Big(\int_1^\infty e^{-t}t^{-s}\,dt\Big)\frac{\sin \pi s}{\pi}.$$

$\left|\int_1^\infty e^{-t}t^{-s}\,dt\right| \le \int_1^\infty e^{-t}t^{|\sigma|}\,dt \le e^{(|s|+1)\log(|s|+1)}$ and $|\sin \pi s| \le e^{\pi|s|}$, so the second term is $\le c_1 e^{c_2|s|\log|s|}$. For the first term: if $|\mathrm{Im}(s)| > 1$ it is $\le ce^{\pi|s|}$. If $|\mathrm{Im}(s)| \le 1$, let k be the integer with $k - 1/2 \le \mathrm{Re}(s) < k + 1/2$. If $k \ge 1$,

$$\sum_{n=0}^\infty \frac{(-1)^n}{n!(n+1-s)}\frac{\sin \pi s}{\pi} = (-1)^{k-1}\frac{\sin \pi s}{\pi(k-1)!(k-s)} + \sum_{n\ne k-1}\frac{(-1)^n}{n!(n+1-s)}\frac{\sin \pi s}{\pi};$$

the first term is bounded since $\sin \pi s$ vanishes at $s = k$, and the second is majorized by $c\sum 1/n!$. If $k \le 0$ then $\mathrm{Re}(s) < 1/2$ and the series is again bounded by $c\sum 1/n!$. ∎

Theorem 1.7

For all $s \in \mathbb{C}$,

$$\frac{1}{\Gamma(s)} = e^{\gamma s} s \prod_{n=1}^{\infty} \left(1 + \frac{s}{n}\right) e^{-s/n}.$$

Here γ is **Euler's constant**, $\gamma = \lim_{N\to\infty} \left(\sum_{n=1}^{N} \frac{1}{n} - \log N\right)$. The limit exists: $\sum_{n=1}^{N} \frac{1}{n} - \log N = \sum_{n=1}^{N-1} \int_n^{n+1} \left[\frac{1}{n} - \frac{1}{x}\right] dx + \frac{1}{N}$, and by the mean value theorem $0 \le \frac{1}{n} - \frac{1}{x} \le \frac{1}{n^2}$ for $n \le x \le n+1$; so the sum is $\sum_{n=1}^{N-1} a_n + \frac{1}{N}$ with $0 \le a_n \le 1/n^2$, and $\sum a_n$ converges.

Proof. By Hadamard's theorem, since $1/\Gamma$ is entire of growth order 1 with simple zeros at $0, -1, -2, \ldots$,

$$\frac{1}{\Gamma(s)} = e^{As+B} s \prod_{n=1}^{\infty} \left(1 + \frac{s}{n}\right) e^{-s/n}.$$

Since $s\Gamma(s) \to 1$ as $s \to 0$, $B = 0$ (or an integer multiple of $2\pi i$, which gives the same result). Putting $s = 1$ and using $\Gamma(1) = 1$,

$$e^{-A} = \prod_{n=1}^{\infty} \left(1 + \frac{1}{n}\right) e^{-1/n} = \lim_{N\to\infty} e^{\sum_{n=1}^{N} [\log(1+1/n) - 1/n]} = \lim_{N\to\infty} e^{-(\sum_{n=1}^{N} 1/n) + \log N + \log(1+1/N)} = e^{-\gamma}.$$

So $A = \gamma + 2\pi i k$; since $\Gamma(s)$ is real for real s, $k = 0$. ∎

So $1/\Gamma$ is characterized (up to two constants) as entire with (i) simple zeros exactly at $s = 0, -1, -2, \ldots$ and (ii) order 1. $\sin \pi s$ (zeros at all integers) satisfies the stricter $O(e^{c|s|})$, which fails for $1/\Gamma$.

2 The zeta function

Motivation. To use complex analysis on ζ, continue it to $\mathbb{C} - \{1\}$ and relate $\zeta(s)$ to $\zeta(1-s)$.

The **Riemann zeta function** is initially defined for real $s > 1$ by $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$.

2.1 Functional equation and analytic continuation

Proposition 2.1

The series defining $\zeta(s)$ converges for $\mathrm{Re}(s) > 1$, and the function ζ is holomorphic in this half-plane.

Proof. For $s = \sigma + it$, $|n^{-s}| = |e^{-s \log n}| = e^{-\sigma \log n} = n^{-\sigma}$. If $\sigma \ge 1 + \delta > 1$ the series is uniformly bounded by $\sum 1/n^{1+\delta} < \infty$; so it converges uniformly on each half-plane $\mathrm{Re}(s) \ge 1 + \delta$ and defines a holomorphic function in $\mathrm{Re}(s) > 1$. ∎

The continuation is subtler than for Γ; it relates ζ to the **theta function** $\vartheta(t) = \sum_{n=-\infty}^{\infty} e^{-\pi n^2 t}$, $t > 0$, which satisfies (by Poisson summation, (4) of Chapter 4) $\vartheta(t) = t^{-1/2}\vartheta(1/t)$. We need

$$\vartheta(t) \le C t^{-1/2} \quad \text{as } t \to 0, \qquad |\vartheta(t) - 1| \le C e^{-\pi t} \quad \text{for } t \ge 1:$$

the first follows from the functional equation, the second from $\sum_{n\ge1} e^{-\pi n^2 t} \le \sum_{n\ge1} e^{-\pi n t} \le C e^{-\pi t}$ for $t \ge 1$.

Theorem 2.2

If $\mathrm{Re}(s) > 1$, then

$$\pi^{-s/2}\Gamma(s/2)\zeta(s) = \frac{1}{2}\int_0^\infty u^{(s/2)-1}[\vartheta(u) - 1]\,du.$$

Proof. The key is

$$\int_0^\infty e^{-\pi n^2 u} u^{(s/2)-1}\,du = \pi^{-s/2}\Gamma(s/2)n^{-s}, \qquad n \geq 1 : \tag{6}$$

with $u = t/\pi n^2$ the left side becomes $\left(\int_0^\infty e^{-t}t^{(s/2)-1}\,dt\right)(\pi n^2)^{-s/2}$. Since $\frac{\vartheta(u)-1}{2} = \sum_{n=1}^\infty e^{-\pi n^2 u}$, and the estimates on ϑ justify interchanging sum and integral,

$$\frac{1}{2}\int_0^\infty u^{(s/2)-1}[\vartheta(u) - 1]\,du = \sum_{n=1}^\infty \int_0^\infty u^{(s/2)-1}e^{-\pi n^2 u}\,du = \pi^{-s/2}\Gamma(s/2)\sum_{n=1}^\infty n^{-s} = \pi^{-s/2}\Gamma(s/2)\zeta(s).$$

■

The **xi function**, a more symmetric modification of ζ, is defined for $\mathrm{Re}(s) > 1$ by

$$\xi(s) = \pi^{-s/2}\Gamma(s/2)\zeta(s). \tag{7}$$

Theorem 2.3

ξ is holomorphic for $\mathrm{Re}(s) > 1$ and continues meromorphically to $\mathbb{C}$ with simple poles at $s = 0, 1$, and $\xi(s) = \xi(1 - s)$.

Proof. Heuristically: multiply $\sum_n e^{-\pi n^2 u} = u^{-1/2}\sum_n e^{-\pi n^2/u}$ by $u^{(s/2)-1}$ and integrate, using (6) and the change $u \mapsto 1/u$ (discarding the $n = 0$ terms). Precisely: let $\psi(u) = [\vartheta(u) - 1]/2$. The functional equation of ϑ gives

$$\psi(u) = u^{-1/2}\psi(1/u) + \frac{1}{2u^{1/2}} - \frac{1}{2}.$$

By Theorem 2.2, for $\mathrm{Re}(s) > 1$,

$$\begin{aligned}
\pi^{-s/2}\Gamma(s/2)\zeta(s) &= \int_0^1 u^{(s/2)-1}\psi(u)\,du + \int_1^\infty u^{(s/2)-1}\psi(u)\,du \\
&= \int_0^1 u^{(s/2)-1}\Big[u^{-1/2}\psi(1/u) + \frac{1}{2u^{1/2}} - \frac{1}{2}\Big]du + \int_1^\infty u^{(s/2)-1}\psi(u)\,du \\
&= \frac{1}{s-1} - \frac{1}{s} + \int_1^\infty (u^{(-s/2)-1/2} + u^{(s/2)-1})\psi(u)\,du
\end{aligned}$$

(substituting $u \mapsto 1/u$ in the first integral). Therefore

$$\xi(s) = \frac{1}{s-1} - \frac{1}{s} + \int_1^\infty (u^{(-s/2)-1/2} + u^{(s/2)-1})\psi(u)\,du.$$

As ψ decays exponentially, the integral is entire, giving the continuation with simple poles at $0, 1$; the integral and $\frac{1}{s-1} - \frac{1}{s}$ are invariant under $s \mapsto 1 - s$. ■

Theorem 2.4

The zeta function has a meromorphic continuation into the entire complex plane, whose only singularity is a simple pole at $s = 1$.

Proof. By (7), $\zeta(s) = \pi^{s/2}\xi(s)/\Gamma(s/2)$; the simple zero of $1/\Gamma(s/2)$ at 0 cancels the pole of ξ, so ζ has only a simple pole at $s = 1$. ■

A more elementary approach gives the continuation to $\mathrm{Re}(s) > 0$ and is useful for the growth of ζ near $\mathrm{Re}(s) = 1$: compare $\sum n^{-s}$ with $\int_1^\infty x^{-s}\,dx$.

Proposition 2.5

There is a sequence of entire functions $\{\delta_n(s)\}_{n=1}^{\infty}$ that satisfy the estimate $|\delta_n(s)| \le |s|/n^{\sigma+1}$, where $s = \sigma + it$, and such that

$$\sum_{1\le n<N} \frac{1}{n^s} - \int_1^N \frac{dx}{x^s} = \sum_{1\le n<N} \delta_n(s), \tag{8}$$

whenever N is an integer > 1.

Proof. Set

$$\delta_n(s) = \int_n^{n+1} \Big[\frac{1}{n^s} - \frac{1}{x^s}\Big] dx. \tag{9}$$

By the mean-value theorem applied to x^{-s}, $\big|\frac{1}{n^s} - \frac{1}{x^s}\big| \le \frac{|s|}{n^{\sigma+1}}$ for $n \le x \le n+1$, so $|\delta_n(s)| \le |s|/n^{\sigma+1}$; and $\int_1^N \frac{dx}{x^s} = \sum_{1\le n<N} \int_n^{n+1} \frac{dx}{x^s}$. ∎

Corollary 2.6

For $\mathrm{Re}(s) > 0$ we have

$$\zeta(s) - \frac{1}{s-1} = H(s),$$

where $H(s) = \sum_{n=1}^{\infty} \delta_n(s)$ is holomorphic in the half-plane $\mathrm{Re}(s) > 0$.

Proof. For $\mathrm{Re}(s) > 1$ let $N \to \infty$ in (8): $\int_1^N x^{-s} dx \to \frac{1}{s-1}$, and $|\delta_n(s)| \le |s|/n^{\sigma+1}$ makes $\sum \delta_n(s)$ converge uniformly on compacta of $\mathrm{Re}(s) \ge \delta > 0$. So the identity holds for $\mathrm{Re}(s) > 1$, and H is holomorphic in $\mathrm{Re}(s) > 0$, extending ζ there. ∎

For $\mathrm{Re}(s) > 1$, $|\zeta(s)| \le \sum n^{-\sigma}$, so ζ is bounded in $\mathrm{Re}(s) \ge 1 + \delta$. The growth near $\mathrm{Re}(s) = 1$ is mild (the following estimates are crude but suffice for Chapter 7).

Proposition 2.7

Suppose $s = \sigma + it$ with $\sigma, t \in \mathbb{R}$. Then for each σ_0, $0 \le \sigma_0 \le 1$, and every $\epsilon > 0$, there exists a constant c_ϵ so that (i) $|\zeta(s)| \le c_\epsilon |t|^{1-\sigma_0+\epsilon}$, if $\sigma_0 \le \sigma$ and $|t| \ge 1$.
(ii) $|\zeta'(s)| \le c_\epsilon |t|^{\epsilon}$, if $1 \le \sigma$ and $|t| \ge 1$.

In particular $\zeta(1 + it) = O(|t|^{\epsilon})$ as $|t| \to \infty$, and likewise for ζ'.

Proof. Besides $|\delta_n(s)| \le |s|/n^{\sigma+1}$ we have $|\delta_n(s)| \le 2/n^{\sigma}$, from (9) and $|n^{-s}| = n^{-\sigma}$, $|x^{-s}| \le n^{-\sigma}$ for $x \ge n$. Combining via $A = A^{\delta} A^{1-\delta}$ $(0 \le \delta \le 1)$,

$$|\delta_n(s)| \le \Big(\frac{|s|}{n^{\sigma_0+1}}\Big)^{\delta} \Big(\frac{2}{n^{\sigma_0}}\Big)^{1-\delta} \le \frac{2|s|^{\delta}}{n^{\sigma_0+\delta}} \qquad (\sigma \ge \sigma_0).$$

Take $\delta = 1 - \sigma_0 + \epsilon$ and use Corollary 2.6: for $\sigma \ge \sigma_0$,

$$|\zeta(s)| \le \Big|\frac{1}{s-1}\Big| + 2|s|^{1-\sigma_0+\epsilon} \sum_{n=1}^{\infty} \frac{1}{n^{1+\epsilon}},$$

which gives (i). For (ii), by the Cauchy integral formula on the circle of radius r about s,

$$\zeta'(s) = \frac{1}{2\pi r} \int_0^{2\pi} \zeta(s + re^{i\theta}) e^{-i\theta}\, d\theta.$$

Take $r = \epsilon$; this circle lies in $\mathrm{Re} \ge 1 - \epsilon$, and (ii) follows from (i) (with $\sigma_0 = 1 - \epsilon$ and 2ϵ in place of ϵ). ∎

3 Selected exercises

Motivation. Four exercises from Stein–Shakarchi, Chapter 6 (book numbers kept).

Exercise 6.1 (Gauss's limit formula)

Prove that

$$\Gamma(s) = \lim_{n\to\infty} \frac{n^s\, n!}{s(s+1)\cdots(s+n)} \qquad \text{whenever } s \neq 0, -1, -2, \dots.$$

Solution.

Step 1. Let $H_n = \sum_{k=1}^{n} 1/k$. The partial products of Theorem 1.7 are

$$P_n(s) = e^{\gamma s} s \prod_{k=1}^{n} \Big(1 + \frac{s}{k}\Big) e^{-s/k},$$

and $P_n(s) \to 1/\Gamma(s)$ for every s.

Step 2. Since $s\prod_{k\le n}(1+s/k) = s(s+1)\cdots(s+n)/n!$ and $n^{-s} = e^{-s\log n}$,

$$P_n(s) = \frac{s(s+1)\cdots(s+n)}{n!\, n^s}\, e^{-s(H_n - \log n - \gamma)}.$$

Step 3. $H_n - \log n - \gamma \to 0$, so $s(s+1)\cdots(s+n)/(n!\,n^s) \to 1/\Gamma(s)$, non-zero for $s \neq 0, -1, \dots$ (Theorem 1.6); take reciprocals. ∎

Exercise 6.3 (Duplication formula)

Using Exercise 5.6(b), prove that $\Gamma(s)\Gamma(s+1/2) = 2^{1-2s}\sqrt{\pi}\,\Gamma(2s)$ for all s with $2s \neq 0, -1, -2, \dots$.

Solution. Fix s with $2s \notin \{0, -1, -2, \dots\}$, so none of s, $s+1/2$, $2s$ is a pole of Γ.

Step 1. By Exercise 6.1 applied to s and $s+\frac{1}{2}$ (with the same n),

$$\Gamma(s)\Gamma(s+\tfrac{1}{2}) = \lim_{n\to\infty} \frac{n^{2s+1/2}(n!)^2}{\prod_{k=0}^{n}(s+k)(s+k+\frac{1}{2})}.$$

Step 2. Since $(s+k)(s+k+\frac{1}{2}) = \frac{1}{4}(2s+2k)(2s+2k+1)$,

$$\prod_{k=0}^{n}(s+k)(s+k+\tfrac{1}{2}) = 2^{-2n-2}\prod_{j=0}^{2n+1}(2s+j).$$

Step 3. Exercise 6.1 applied to $2s$, along the subsequence $2n+1$, gives

$$\Gamma(2s) = \lim_{n\to\infty} \frac{(2n+1)^{2s}(2n+1)!}{\prod_{j=0}^{2n+1}(2s+j)}.$$

Step 4. Dividing the limits in Steps 1 and 3 (the second is non-zero), and using Step 2,

$$\frac{\Gamma(s)\Gamma(s+\frac{1}{2})}{\Gamma(2s)} = \lim_{n\to\infty} \Big(\frac{n}{2n+1}\Big)^{2s} \cdot \frac{2^{2n+2}\, n^{1/2}(n!)^2}{(2n+1)!}.$$

Step 5. We evaluate the two factors. First, $(n/(2n+1))^{2s} = e^{2s\log(n/(2n+1))} \to e^{-2s\log 2} = 2^{-2s}$. Second, since $(2n+1)! = (2n)!\,(2n+1)$,

$$\frac{2^{2n+2}n^{1/2}(n!)^2}{(2n+1)!} = 4\cdot\frac{2^{2n}(n!)^2}{(2n)!\,(2n+1)^{1/2}}\cdot\Big(\frac{n}{2n+1}\Big)^{1/2} \to 4\sqrt{\frac{\pi}{2}}\cdot\frac{1}{\sqrt{2}} = 2\sqrt{\pi},$$

by Exercise 5.6(b).

Step 6. Hence $\Gamma(s)\Gamma(s+\frac12) = 2^{-2s}\cdot 2\sqrt{\pi}\,\Gamma(2s) = 2^{1-2s}\sqrt{\pi}\,\Gamma(2s)$. ∎

Exercise 6.5 (Γ on the line $\mathrm{Re}(s) = 1/2$)

Show that $|\Gamma(\frac12 + it)|^2 = \dfrac{2\pi}{e^{\pi t} + e^{-\pi t}}$ for $t \in \mathbb{R}$.

Solution.

Step 1. For $\mathrm{Re}(s) > 0$, conjugating the integral (1) gives $\overline{\Gamma(s)} = \int_0^\infty e^{-u}u^{\bar s - 1}\,du = \Gamma(\bar s)$. Both $s \mapsto \overline{\Gamma(\bar s)}$ and Γ are meromorphic on $\mathbb{C}$ and agree on $\mathrm{Re}(s) > 0$, so $\overline{\Gamma(s)} = \Gamma(\bar s)$ for all s off the poles.

Step 2. For $s = \frac12 + it$ we have $\bar s = \frac12 - it = 1 - s$. By Theorem 1.4,

$$|\Gamma(s)|^2 = \Gamma(s)\Gamma(\bar s) = \Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin(\frac{\pi}{2} + i\pi t)} = \frac{\pi}{\cos(i\pi t)} = \frac{\pi}{\cosh \pi t} = \frac{2\pi}{e^{\pi t} + e^{-\pi t}}.$$

∎

Exercise 6.7 (The Beta function)

For $\mathrm{Re}(\alpha) > 0$ and $\mathrm{Re}(\beta) > 0$ let $B(\alpha,\beta) = \int_0^1 (1-t)^{\alpha-1}t^{\beta-1}\,dt$.

(a) Prove that $B(\alpha,\beta) = \dfrac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$.

(b) Show that $B(\alpha,\beta) = \displaystyle\int_0^\infty \frac{u^{\alpha-1}}{(1+u)^{\alpha+\beta}}\,du$.

Solution. All powers of positive reals are principal, so $(xy)^w = x^w y^w$ and $|x^w| = x^{\mathrm{Re}\,w}$ for $x, y > 0$. In particular $|(1-t)^{\alpha-1}t^{\beta-1}| = (1-t)^{\mathrm{Re}\,\alpha-1}t^{\mathrm{Re}\,\beta-1}$ is integrable on $(0,1)$, so B is well defined.

(a)

Step 1. The double integral

$$\Gamma(\alpha)\Gamma(\beta) = \int_0^\infty\int_0^\infty t^{\alpha-1}s^{\beta-1}e^{-t-s}\,dt\,ds$$

converges absolutely, since the modulus of the integrand is $t^{\mathrm{Re}\,\alpha-1}s^{\mathrm{Re}\,\beta-1}e^{-t-s}$. Hence Fubini's theorem and changes of variables are allowed.

Step 2. The map $(u,r) \mapsto (t,s) = (u(1-r), ur)$ is a C^1 bijection from $(0,\infty)\times(0,1)$ onto $(0,\infty)^2$, with inverse $u = t+s$, $r = s/(t+s)$. Its Jacobian determinant is $(1-r)u + ru = u$.

Step 3. Therefore

$$\Gamma(\alpha)\Gamma(\beta) = \int_0^\infty\int_0^1 u^{\alpha+\beta-1}(1-r)^{\alpha-1}r^{\beta-1}e^{-u}\,dr\,du = \Gamma(\alpha+\beta)\,B(\alpha,\beta).$$

Since Γ never vanishes (Theorem 1.6(i)), we may divide by $\Gamma(\alpha+\beta)$.

(b) Substitute $t = 1/(1+u)$, $u \in (0,\infty)$. Then $1-t = u/(1+u)$ and $dt = -du/(1+u)^2$, and t decreases from 1 to 0. Hence

$$B(\alpha,\beta) = \int_0^\infty \frac{u^{\alpha-1}}{(1+u)^{\alpha-1}} \frac{1}{(1+u)^{\beta-1}} \frac{du}{(1+u)^2} = \int_0^\infty \frac{u^{\alpha-1}}{(1+u)^{\alpha+\beta}}\, du.$$

■