Lecture 3. Basic Parametric Distributions and Models

Contents

1 Outline 1

2 Exponential Distribution 1

3 Weibull Distribution 2

4 Weibull and the Extreme Value Distribution 2

5 Log-normal Distribution 2

6 Log-logistic Distribution 3

7 Gamma Distribution 3

8 Regression Models for Survival Data 4

8.1 Approach 1: Log-linear / accelerated failure time (AFT) model 4

8.2 Approach 2: Multiplicative (proportional hazards) model 5

8.3 Approach 3: Additive hazards model 6

1 Outline

- Parametric families: exponential, Weibull, extreme value, log-normal, log-logistic, gamma.
- Regression: accelerated failure time (log-linear), multiplicative (PH), additive hazards.

2 Exponential Distribution

Definition 3.1: Exponential(λ)

$f(x) = \lambda e^{-\lambda x}$, $S(x) = e^{-\lambda x}$, $h(x) = \lambda$ (constant), $H(x) = \lambda x$.

- Memoryless: the failure rate given survival to t does not depend on t.
- **Empirical check**: $\hat{H}(x)$ against x should be a line through the origin.
- Simple but too restrictive in practice.
- Weibull with $\alpha = 1$; gamma with $\beta = 1$.

3 Weibull Distribution

Definition 3.2: Weibull(λ, α), $\lambda, \alpha > 0$

$$h(x) = \lambda\alpha x^{\alpha-1}, \qquad H(x) = \lambda x^{\alpha}, \qquad S(x) = \exp(-\lambda x^{\alpha}).$$

- Simple yet flexible: hazard increasing ($\alpha > 1$), decreasing ($\alpha < 1$), constant ($\alpha = 1$).
- **Empirical check**: $\ln H(x) = \ln\lambda + \alpha \ln x$, so $\ln \hat{H}(x)$ against $\ln x$ should be linear (slope α).

4 Weibull and the Extreme Value Distribution

Motivation. On the log scale the Weibull is a location–scale model, the basis of AFT regression.

Definition 3.3: Standard extreme value distribution

$$f_W(w) = \exp\{w - e^w\}, \qquad S_W(w) = \exp\{-e^w\}, \qquad -\infty < w < \infty.$$

A special case of the Gompertz family.

Theorem 3.4: Log-linear representation of the Weibull

If $Y \equiv \ln X = \mu + \sigma W$ with W standard extreme value, then $X \sim \text{Weibull}(\lambda, \alpha)$ with

$$\alpha = \frac{1}{\sigma}, \qquad \lambda = e^{-\mu/\sigma} \quad \Big(\text{equivalently } \mu = \frac{-\ln\lambda}{\alpha},\ \sigma = \frac{1}{\alpha}\Big).$$

Proof [Supplement]. $S_W(w) = \int_w^\infty e^{v-e^v}\,dv = \int_{e^w}^\infty e^{-u}\,du = \exp(-e^w)$ $(u = e^v)$. Then

$$S_X(x) = P(\ln X > \ln x) = P\Big(W > \frac{\ln x - \mu}{\sigma}\Big) = \exp\Big\{-e^{(\ln x-\mu)/\sigma}\Big\} = \exp\Big\{-e^{-\mu/\sigma}x^{1/\sigma}\Big\},$$

i.e. $\exp(-\lambda x^{\alpha})$. ■

So a log-linear model with extreme value error gives a Weibull X.

5 Log-normal Distribution

Definition 3.5

If $Y = \ln X \sim N(\mu, \sigma^2)$ then $X \sim$ log-normal(μ, σ^2): $S(x) = 1 - \Phi\Big(\frac{\ln x - \mu}{\sigma}\Big)$.

- Close tie to the normal.
- Hump-shaped hazards: from 0 up to a maximum, then decreasing.
- Criticized because the hazard *always decreases to 0 in the tail.*
- Adequate when large x are not of interest.

6 Log-logistic Distribution

Motivation. The log-logistic allows rising-then-falling hazards in closed form.

Definition 3.6: Log-logistic

If $Y = \ln X = \mu + \sigma W$ with W standard logistic, LG$(0,1)$:

$$f_W(w) = \frac{e^w}{(1+e^w)^2}, \qquad S_W(w) = \frac{1}{1+e^w},$$

then $Y \sim \text{LG}(\mu, \sigma^2)$ with $f_Y(y) = \dfrac{e^{(y-\mu)/\sigma}}{\sigma\left(1+e^{(y-\mu)/\sigma}\right)^2}$ and $X \sim$ log-logistic.

Derivation: survival function of X

$$S_X(x) = S_W\Big(\frac{\ln x - \mu}{\sigma}\Big) = \frac{1}{1+\exp\left(\frac{\ln x-\mu}{\sigma}\right)} = \frac{1}{1+e^{-\mu/\sigma}x^{1/\sigma}} = \frac{1}{1+\lambda x^\alpha},$$

with $\alpha = 1/\sigma$ and $\lambda = e^{-\mu/\sigma}$.

Theorem 3.7: Log-logistic hazard

$$h_X(x) = \frac{\lambda\alpha x^{\alpha-1}}{1+\lambda x^\alpha}.$$

It is monotone decreasing when $\alpha \le 1$. When $\alpha > 1$ it increases to a maximum at $x^* = \{(\alpha-1)/\lambda\}^{1/\alpha}$ and then decreases to 0 as $x \to \infty$.

Proof [Supplement]. $H(x) = -\ln S(x) = \ln(1+\lambda x^\alpha)$, so $h(x) = H'(x) = \lambda\alpha x^{\alpha-1}/(1+\lambda x^\alpha)$. Differentiating,

$$h'(x) = \lambda\alpha\,\frac{(\alpha-1)x^{\alpha-2}(1+\lambda x^\alpha) - x^{\alpha-1}\lambda\alpha x^{\alpha-1}}{(1+\lambda x^\alpha)^2} = \lambda\alpha\,\frac{x^{\alpha-2}\left[(\alpha-1)-\lambda x^\alpha\right]}{(1+\lambda x^\alpha)^2}.$$

For $\alpha \le 1$ the bracket is negative; for $\alpha > 1$ it changes sign at x^*; and $h(x) \sim \alpha/x \to 0$. ■

Like the log-normal hazard, but in closed form.

7 Gamma Distribution

Definition 3.8: Gamma(λ, β), $\lambda, \beta > 0$

$$f(x) = \frac{\lambda^\beta x^{\beta-1}e^{-\lambda x}}{\Gamma(\beta)}, \qquad S(x) = \int_x^\infty \frac{\lambda^\beta s^{\beta-1}e^{-\lambda s}}{\Gamma(\beta)}\,\mathrm{d}s, \qquad h(x) = \frac{x^{\beta-1}e^{-\lambda x}}{\int_x^\infty s^{\beta-1}e^{-\lambda s}\,\mathrm{d}s}.$$

No closed form for S, h in general.

- $\beta = 1$: exponential(λ). $\beta \to \infty$: approximately normal. $\lambda = 1/2$, integer β: $\chi^2_{2\beta}$.
- $\beta > 1$: h increases from $h(0) = 0$ to λ.
- $\beta < 1$: h decreases from $h(0) = \infty$ to λ.

- Less tractable; the Weibull is more popular.

Derivation: why the gamma hazard tends to λ monotonically

Substitute $s = x + u$ in the reciprocal hazard:

$$\frac{1}{h(x)} = \int_x^\infty \left(\frac{s}{x}\right)^{\beta-1} e^{-\lambda(s-x)}\,\mathrm{d}s = \int_0^\infty \left(1+\frac{u}{x}\right)^{\beta-1} e^{-\lambda u}\,\mathrm{d}u.$$

As $x \to \infty$, $1/h(x) \to \int_0^\infty e^{-\lambda u}\,\mathrm{d}u = 1/\lambda$. For $\beta > 1$, $(1+u/x)^{\beta-1}$ decreases in x, so h increases; for $\beta < 1$ the reverse.

8 Regression Models for Survival Data

Heterogeneity requires covariate adjustment, e.g. for groups imbalanced in age, tumour size, positive nodes or hormonal status in non-randomized studies.

8.1 Approach 1: Log-linear / accelerated failure time (AFT) model

Definition 3.9: Log-linear model

$$Y \equiv \ln X = \mu + \boldsymbol{\gamma}'\mathbf{z} + \sigma W,$$

with $W \sim F$ known and $\mu, \sigma, \boldsymbol{\gamma}$ unknown.

Theorem 3.10: Log-linear $\Rightarrow$ AFT

Let $X_0 = \exp(\mu + \sigma W)$ with survival S_0. Then

$$S(x \mid \mathbf{z}) = S_0(x\,e^{-\boldsymbol{\gamma}'\mathbf{z}}), \qquad h(x \mid \mathbf{z}) = \eta\, h_0(\eta x), \quad \eta = e^{-\boldsymbol{\gamma}'\mathbf{z}}.$$

Proof [Supplement].

$$\begin{aligned} P(X > x \mid \mathbf{z}) &= P(Y > \ln x \mid \mathbf{z}) = P(\mu + \sigma W > \ln x - \boldsymbol{\gamma}'\mathbf{z}) \\ &= P\big(\exp(\mu + \sigma W) > x \exp(-\boldsymbol{\gamma}'\mathbf{z})\big) \\ &= S_0(xe^{-\boldsymbol{\gamma}'\mathbf{z}}). \end{aligned}$$

$S(x \mid \mathbf{z}) = S_0(\eta x)$ gives $f(x \mid \mathbf{z}) = \eta f_0(\eta x)$, so

$$h(x \mid \mathbf{z}) = \eta f_0(\eta x)/S_0(\eta x) = \eta h_0(\eta x).$$

■

- Covariates rescale time: **accelerated** if $\eta = e^{-\boldsymbol{\gamma}'\mathbf{z}} > 1$, **decelerated** if $\eta < 1$.
- Signs are **opposite**: positive γ lengthens survival ($\eta < 1$).

Derivation: interpretation of γ

Since $X = e^{\mu+\boldsymbol{\gamma}'\mathbf{z}}e^{\sigma W}$,

$$\frac{\mathrm{E}(X \mid \mathbf{z}_1)}{\mathrm{E}(X \mid \mathbf{z}_2)} = \frac{e^{\mu+\boldsymbol{\gamma}'\mathbf{z}_1}\mathrm{E}(e^{\sigma W})}{e^{\mu+\boldsymbol{\gamma}'\mathbf{z}_2}\mathrm{E}(e^{\sigma W})} = e^{\boldsymbol{\gamma}'(\mathbf{z}_1-\mathbf{z}_2)} \implies \ln\frac{\mathrm{E}(X \mid \mathbf{z}_1)}{\mathrm{E}(X \mid \mathbf{z}_2)} = \boldsymbol{\gamma}'(\mathbf{z}_1-\mathbf{z}_2).$$

γ is the log ratio of mean (or any quantile of) survival times per unit increase in z.

Definition 3.11: Three choices of F

1. $W \sim N(0,1)$: $\ln X \sim N(\mu+\boldsymbol{\gamma}'\mathbf{z}, \sigma^2)$, i.e. X is **log-normal**.
2. W standard extreme value, $f_W(w) = \exp(w - e^w)$: $X \sim \text{Weibull}(\lambda, \alpha)$ with $\alpha = 1/\sigma$ and $\lambda = \exp\{-(\frac{\mu}{\sigma} + \frac{\boldsymbol{\gamma}'\mathbf{z}}{\sigma})\}$.
3. W standard logistic: the **log-logistic** regression model.

8.2 Approach 2: Multiplicative (proportional hazards) model

Definition 3.12: Multiplicative hazards model

$$h(x \mid \mathbf{z}) = h_0(x)\, C(\boldsymbol{\beta}'\mathbf{z}),$$

with baseline $h_0(x) \geq 0$ parametric or unspecified; $C(\boldsymbol{\beta}'\mathbf{z}) = e^{\boldsymbol{\beta}'\mathbf{z}}$ gives the **Cox proportional hazards model**.

- β is the **log hazard ratio**, constant over time: $h(x \mid \mathbf{z}_1)/h(x \mid \mathbf{z}_2) = e^{\boldsymbol{\beta}'(\mathbf{z}_1-\mathbf{z}_2)}$.
- Integrating, $H(x \mid \mathbf{z}) = e^{\boldsymbol{\beta}'\mathbf{z}}H_0(x)$ and $S(x \mid \mathbf{z}) = \exp\{-e^{\boldsymbol{\beta}'\mathbf{z}}H_0(x)\} = \{S_0(x)\}^{e^{\boldsymbol{\beta}'\mathbf{z}}}$.
- **Graphical check**: $\ln[-\ln S(x \mid \mathbf{z})] = \boldsymbol{\beta}'\mathbf{z} + \ln[-\ln S_0(x)]$, so log-cumulative-hazard curves should be *parallel*.

Example 3.13: Weibull baseline under PH

If $h_0(x) = \alpha\lambda x^{\alpha-1}$ then $h(x \mid \mathbf{z}) = \alpha\lambda x^{\alpha-1}\exp(\boldsymbol{\beta}'\mathbf{z})$ and

$$S(x \mid \mathbf{z}) = \exp\left[-\lambda\exp(\boldsymbol{\beta}'\mathbf{z})x^{\alpha}\right], \quad \text{i.e.} \quad X \mid \mathbf{z} \sim \text{Weibull}(\lambda e^{\boldsymbol{\beta}'\mathbf{z}}, \alpha).$$

Theorem 3.14: Weibull is closed under both AFT and PH

The Weibull is the only continuous distribution closed under both AFT and PH.

Derivation: matching the two parametrizations

Under AFT with Weibull baseline, $h(x \mid \mathbf{z}) = \eta\,\alpha\lambda(\eta x)^{\alpha-1} = \eta^{\alpha}\cdot\alpha\lambda x^{\alpha-1}$: PH with hazard ratio $\eta^{\alpha} = e^{-\alpha\boldsymbol{\gamma}'\mathbf{z}}$, so

$$\boldsymbol{\beta} = -\alpha\boldsymbol{\gamma} = -\boldsymbol{\gamma}/\sigma.$$

8.3 Approach 3: Additive hazards model

Definition 3.15: Additive hazards model

$$h(x \mid \mathbf{z}) = h_0(x) + \sum_{j=1}^{p} z_j(x)\beta_j(x).$$

- Covariate effects may vary over time.
- $\beta_j(x)$ must be constrained so that $h(x \mid \mathbf{z}) \geq 0$.