

Two Field Spread: numerical algorithms

v0.4.1 implementation and next experiment specification

24 September 2026

Status. The Markov pilot and execution tape are implemented and tested. Long-run LMF diagnostics, convergence acceptance, fractional transport and calendar subordination are not completed. This draft describes the actual software; it does not amend the supplied theory paper or letter.

1 Model, agents and the inherited numerical construction

The state is two nonnegative densities, $\rho_B(x_j, u_n)$ and $\rho_A(x_j, u_n)$, in one market. A cell carries volume $\rho_i \Delta x$; x is log price. Fixed endpoint reservoirs are excluded from executable volume. Lit and latent sources aggregate supply; diffusion aggregates repricing; cancellation removes standing volume; symmetric reaction removes overlap. These are an aggregate closure with an order-level random-walk interpretation. The current solver does not simulate individual traders, wealth, beliefs or strategy choice.

The market-making sector is represented by pending completion stocks and source placement. All explicitly executed child volume in this pilot is attributed to that sector. Its opposite-side source is *completion-equivalent*: survival and execution probabilities are assumed inside the completion/placement specification. Source delivery is neither a new trade print nor independently observed settlement. Reaction loss is accounted for as a field sink, without inventing aggressor signs or prices for it. Thus the trade tape records the specified aggressor programme, not every possible microscopic match.

Let Q_B denote bid replenishment owed after sales and Q_A ask replenishment owed after purchases. The quote-setting sample obeys

$$Q_\Sigma = Q_A + Q_B, \quad Q_\Delta = Q_A - Q_B, \quad (1)$$

$$\Sigma = \Sigma_0 + \chi_s Q_\Sigma, \quad \mu = \mu_0 - \chi_m Q_\Delta, \quad q_b = \mu - \Sigma/2, \quad q_a = \mu + \Sigma/2. \quad (2)$$

Observed quotes p_b, p_a are side-density threshold crossings, not q_b, q_a . Zero signed inventory does not imply zero total pending exposure. Also, $\Sigma_0 = 12$ is imposed in this pilot: the existence of its baseline spread is not evidence of spontaneous spread formation from zero standoff.

Diana–Gebbie explicitly identify the need for two densities to represent a nonzero spread [1]. The correlation-emergence software supplies the operational DTRW, execution-record and observation conventions [2, 3]. Its two coupled books are not identified with the two sides of this market. The new density core implements the supplied spread paper’s Markov branch. For the common reaction-free interior operator, the prototype has

$$\rho^{n+1} = e^{-\nu \Delta u} \rho^n + \frac{r}{2} \delta_x^2 \rho^n + \Delta u S^n, \quad r = 2D\Delta u / \Delta x^2,$$

where $\delta_x^2 \rho_j = \rho_{j-1} - 2\rho_j + \rho_{j+1}$. Here cancellation is $(1 - \nu \Delta u) \rho^n$. Their difference is $(e^{-\nu \Delta u} - 1 + \nu \Delta u) \rho^n$, of one-step order Δu^2 . Matched transport agrees to roundoff at $\nu = \kappa = 0$; complete trajectories are not asserted identical. The prototype consumes before its field update; the present paper convention consumes after it. Phase labels preserve this distinction. See `compatibility-v0.4.1.json` for the executed comparison.

2 Algorithm 1: implemented Markov update

Inputs: previous post-event densities, post-event pending stocks P_B, P_A , fixed $\Delta x, \Delta u$, parameters and event-integrated requests. Both sides read one frozen field; new executions cannot complete immediately.

1. Deliver old pending cohorts: $C_i = (1 - e^{-\Delta u/\tau_h})P_i$ and $Q_i = P_i - C_i$. For the next-update control set $C_i = P_i$. This is the only “immediate” convention used. Form q_b, q_a from Q above.
2. Evaluate the paper’s lit and latent sources S_i^{ext} on the outward side of q_i . Normalize uniform nodal weights g_B on $[q_b - w, q_b]$ and g_A on $[q_a, q_a + w]$ so that $\Delta x \sum_j g_{i,j} = 1$. Reject empty or reservoir-truncated supports. No supply is placed between the placement edges.
3. For interior nodes, compute the pre-event field

$$\tilde{\rho}_{i,j} = (1 - 2a - \nu\Delta u - \kappa\Delta u\rho_{i,j})\rho_{i,j} + a(\rho_{i,j-1} + \rho_{i,j+1}) + \Delta u S_{i,j}^{\text{ext}} + C_i g_{i,j}, \quad a = D\Delta u/\Delta x^2.$$

Require $2a + \nu\Delta u + \kappa\Delta u\rho_{i,j} \leq 1$ everywhere. Keep endpoint reservoirs fixed; reject inadmissible states without clipping.

4. Read pre-event quotes by linear interpolation at density threshold ϵ : the rightmost downward bid crossing and leftmost upward ask crossing. Record crossing counts; reject a selected plateau, insufficient selected slope, reservoir failure or contact.
5. Execute each request in outward price priority, starting at its pre-event quote and ending at the configured execution depth. Restrict to interior node centres. Remove at most $\tilde{\rho}_{i,j}\Delta x$ per node; record actual fills, requests and unfilled demand. No reservoir is consumed. The current tape requires one aggressor side per event.
6. Let v_A be actual buy volume and v_B actual sell volume. Set $P_B^{\text{new}} = Q_B + v_A$, $P_A^{\text{new}} = Q_A + v_B$. Read post-event quotes and save both inventory samples, actual removals and separate source increments. An event at u is applied at the end of the interval ending at u ; pre/post fields share u with different phase labels.

Checks on every update. For each side, the interior volume change is boundary flux minus cancellation, minus symmetric reaction, plus external supply, plus completion, minus execution. The boundary increment is $D\Delta u(\rho_0 - \rho_1 + \rho_J - \rho_{J-1})/\Delta x$. Reaction loss is $\kappa\Delta u\Delta x \sum_j \rho_{A,j}\rho_{B,j}$ on each side. It cancels from imbalance and occurs twice in total density. Cumulative initiation must equal cumulative completion plus post-event pending stock, separately by side.

Initialization and recovery. Relax the same positive-reaction equations with empty pending stock and no executions to the configured residual. Recompute geometric completion weights when Δu changes, retaining physical τ_h . Preserve pending tails beyond a finite plotting horizon. Volume-balanced round trips need not restore liquidity at the quote at the same time: delivery occurs elsewhere and transport takes time. Neither the ledger nor a stationary field establishes dealer profitability or a strategic equilibrium. Hard source supports require grid-phase refinement; the current late spread offset has not been accepted as physical.

3 Algorithm 2: implemented execution tape

1. For event n retain nonzero executed node volumes $v_{n,j}$, parent identifier, aggressor sign ε_n (+1 buy, -1 sell), operational index and time. No fill means no trade observation. Preserve each fill row and the requested/unfilled amounts in the event record.
2. Following the prior executable convention, compute the primary execution log price

$$z_n = \frac{\sum_j v_{n,j} x_j}{\sum_j v_{n,j}}.$$

Store e^{z_n} as the geometric execution price and $\sum_j v_{n,j} e^{x_j} / \sum_j v_{n,j}$ as the separate arithmetic price VWAP. Do not call these identical. Store pre/post bid, ask, log midpoint and spread, and the placement and pending state that generated the event.

3. Preserve the true aggressor sign separately from the quote sign $\text{sign}(z_n - m_n^{\text{pre}})$ and tick sign $\text{sign}(z_n - z_{n-1})$. The first tick label is zero; a zero tick subsequently carries the last nonzero tick label, as in v2.0.0.
4. Using consecutive executed events, calculate $r_n^m = m_n^{\text{post}} - m_{n-1}^{\text{post}}$, $r_n^T = z_n - z_{n-1}$ and $\Delta s_n = s_n^{\text{post}} - s_{n-1}^{\text{post}}$. Retain absolute increments. First increments are missing, not zero. An instantaneous pre/post midpoint jump is stored separately from r_n^m , which also includes inter-event field evolution.

Prices are measured from actual density consumption; there is no independent synthetic price process. The current six same-sign children demonstrate the measurement path only. Their sign variance is zero, so no sign ACF or LMF claim is available from this run.

4 Algorithm 3: lagged correlation

The estimator is implemented and tested; the long-run protocol below has not yet been executed. For aligned finite series X, Y , define

$$\hat{C}_{XY}(k) = \frac{\sum_{n \in I_k} (X_n - \bar{X}_k)(Y_{n+k} - \bar{Y}_k)}{\sqrt{\sum_{n \in I_k} (X_n - \bar{X}_k)^2 \sum_{n \in I_k} (Y_{n+k} - \bar{Y}_k)^2}}.$$

Each mean is taken over its own overlapping slice. Positive k means X leads Y . Save pair counts, lag units and phase alignment. Zero variance is undefined, not zero correlation. Do not silently compress missing observations. This reproduces the v2.0.0 executable Pearson ACF. Its supplementary text instead prints a global-mean covariance normalization; that documentation mismatch is not carried into this supplement.

For future runs, retain raw price paths but use $r^m, r^T, |r^m|, |r^T|$ for price ACFs. For a positive price, $|P| = P$; an absolute log-price level depends on the arbitrary origin. Spread-level ACFs require a checked stationary window. Price-level ACFs, if retained, must be labelled descriptive rather than evidence of long memory. A fixed valuation centre and fixed reservoirs can cause mean reversion; diffusive prices are not guaranteed.

5 Next numerical experiments: isolate the market maker

No chartist or fundamentalist population is required for the current spread question. Aggregate source functions can represent their net activity only after a specified behavioural closure. Existing stationary sources do not already encode trend following, valuation switching or persistent order signs. The fixed μ_0 is a valuation reference, not a simulated fundamentalist.

1. First resolve grid, time-step, domain, threshold and support-phase sensitivity for the saved pilot. Retain the supplied equations and physical event volumes; do not smooth the source to force attractive plots.
2. Reuse the same input tape in delayed and next-update completion controls. Separate width feedback ($\chi_s = 0$) from centre feedback ($\chi_m = 0$); the existing jointly frozen control cannot identify them individually. Retain equal actual filled volumes or report that a comparison failed.
3. Add a matched balanced buy/sell programme. Show both Q_Σ and Q_Δ , so small signed inventory with material pending exposure can be distinguished from an empty book. Show q_b, q_a separately from p_b, p_a . During sustained flow, test stable pending exposure and flow balance; after stopping, show remaining completion and density recovery. Do not describe source delivery as a second executed leg.
4. Introduce a minimal aggregate order-splitting generator, with stored parent IDs, independent centred parent signs and an explicitly specified length law. For the LMF benchmark use a parent survival tail $\Pr(L \geq \ell) \sim \ell^{-\beta}$, $1 < \beta < 2$. Under its order-splitting assumptions the sign ACF exponent is $\gamma = \beta - 1$ [4]. Consecutive same-sign parents remain distinct. Register initialization, finite-size truncation, run length, lag window, seeds and independent replicates before inspecting correlation slopes. Match an independent-sign null at the same child volume and rate. A run generator is an aggregate order-flow closure, not evidence of endogenous LMF generation by RD.
5. Apply that tape to the density solver. Measure ACFs of $\varepsilon, r^m, r^T, |r^m|, |r^T|, s$. Measure cross-correlations of (ε, r^m) , $(\varepsilon, \Delta s)$, (r^m, r^T) , $(r^m, \Delta s)$, $(|r^m|, s)$ and $(|r^T|, s)$. Use independent path groups for uncertainty; reflected paths are paired, not additional independent replicates. Check stationarity and censoring; retain all inadmissible exits rather than selecting only successful paths.

Signed midpoint–spread correlation may cancel by buy/sell symmetry; magnitude correlations can still show coupling. Bid/ask bounce may affect short-lag trade returns. Neither is a result established by the current pilot.

Focused figure target. Keep F2 density snapshots, F3 prices and market-maker state (now including actual execution prices), F4 matched mechanism controls and V1 profile video. Add F5: a three-by-three layout with three aligned path panels and the six ACFs. Add F6: the six cross-correlations in square panels. These are numerical outputs from retained trajectories, not a separate theoretical illustration set. Exact panels can share axes and controls without adding a gallery of unrelated stylised facts.

Square-root impact. A single small-volume programme establishes no impact exponent. Any subsequent scaling test must register a matched no-programme baseline, actual executed volume, duration and participation controls, a valid scale window and convergence checks. A pre-contact, finite-threshold spread model does not inherit the single-front square-root limit automatically. The minimal reaction–diffusion impact literature provides a comparison target [5], not a universal acceptance criterion. Broad impact scaling remains a later extension.

6 Future memory and clock compatibility

Keep four mechanisms distinct: event-time order splitting (β), operational transport memory (α_u), round-trip completion (h, τ_h) and calendar observation/subordination (α_t). Give stochastic components separate recorded seed streams. The event/operational/calendar distinction also follows the companion account [6]; its abstract provides context here, not verification of a fractional two-side solver.

The prior raw Sibuya kernel is

$$K_1 = \alpha_u, \quad K_2 = \frac{1}{2}\alpha_u(\alpha_u - 1), \quad K_k = K_{k-1} \left(1 - \frac{2 - \alpha_u}{k}\right), \quad k \geq 3.$$

The prototype applies survival $e^{-\nu(k-1)\Delta u}$ to lagged transport and $e^{-\nu\Delta u}$ to the current survivor, once each. Cancellation is not already folded into the raw kernel. $\alpha_u = 1$ leaves only $K_1 = 1$. Its scaling is $D_{\alpha_u} = r\Delta x^2/(2\Delta u^{\alpha_u})$; changing α_u while silently retaining an incompatible $r, \Delta u, D$ is invalid.

The current state has no transport-history buffer. A future fractional two-side model must derive the survival and history treatment of nonlinear reaction, cancellation, births and consumed volume. It must not resurrect removed liquidity through old history. This requires a stated formulation and tests against the prior primitive and its Markov limit; inserting the kernel into the existing Laplacian alone is insufficient.

The paper's stationary placement construction requires a finite mean completion lag. A heavy-tailed transport waiting time is not permission to replace the completion density with an infinite-mean law. That would change the equilibrium assumptions and can make pending exposure diverge under sustained input.

The correlation-emergence observation layer acts on completed operational paths. A future projection must select the previous completed state under the declared clock, retain index maps and reject out-of-support queries. Do not interpolate across an execution. Use a coherent book observation for bid, ask and midpoint before testing separate reporting delays. A held last trade price creates neither another trade nor another sign. Empty calendar bins have zero signed flow, not a repeated aggressor; distinguish them from missing pre-first-trade observations. Compare shocked/control paths using the same realized clock. These rules are specified, not yet implemented.

References

- [1] D. Diana and T. Gebbie, *Non-uniformly sampled simulated price impact of an order-book*, Journal of Computational and Applied Mathematics **456** (2025), 116202. doi:10.1016/j.cam.2024.116202.
- [2] C. N. Angstmann and T. Gebbie, *Correlation emergence and the Epps effect in two coupled limit order books*, arXiv:2606.14182, version 3.
- [3] T. Gebbie, *correlation-emergence-reproducibility*, v2.0.0 and v2.2.0. Exact commits and inspected component hashes are in SOURCES.md.
- [4] F. Lillo, S. Mike and J. D. Farmer, *A theory for long-memory in supply and demand*, Physical Review E **71** (2005), 066122. arXiv:cond-mat/0412708.
- [5] J. Donier, J. Bonart, I. Mastromatteo and J.-P. Bouchaud, *A fully consistent, minimal model for non-linear market impact*, arXiv:1412.0141.
- [6] C. Angstmann and T. Gebbie, *Event-Time Order-Flow Memory, Operational-Time Impact, and Subordinated Market Observables*, arXiv:2609.13715 (2026).