

1 Factors affecting inferences on natural mortality and
2 associated environmental effects in state-space
3 age-structured assessment models

4 Timothy J. Miller^{1,2} Gregory L. Britten³ Elizabeth N. Brooks²

5 Gavin Fay⁴ Alexander C. Hansell² Christopher M. Legault²

6 Brandon Muffley⁵ John Wiedenmann⁶

7 ~~18 June, 2025~~31 March, 2026

8 ¹corresponding author: timothy.j.miller@noaa.gov

9 ²Northeast Fisheries Science Center, Woods Hole Laboratory, 166 Water Street, Woods
10 Hole, MA 02543 USA

11 ³Biology Department, Woods Hole Oceanographic Institution, 266 Woods Hole Rd. Woods
12 Hole, MA, USA

13 ⁴Department of Fisheries Oceanography, School for Marine Science and Technology,
14 University of Massachusetts Dartmouth, 836 S Rodney French Boulevard, New Bedford,
15 MA 02744, USA

16 ⁵Mid-Atlantic Fishery Management Council, 800 North State Street, Suite 201, Dover, DE
17 19901 USA

¹⁸ ⁶Department of Ecology, Evolution, and Natural Resources. Rutgers University

¹⁹

Abstract

Treatment of natural mortality is a major consideration in assessment models and state-space approaches allow estimation of temporal variation in this mortality rate as well as effects of specific covariates. However, there has been no investigation of the reliability of inferences made ~~regarding natural mortality, associated covariate effects, and important assessment output~~ from state-space assessment models when there is process error or covariate effects on natural mortality. We conducted a large-scale simulation study ~~that considers models fit to data simulated from operating models with alternative assumptions defined by several factors, but we focus on scenarios where there is temporal contrast in fishing pressure and lower uncertainty in population observations (age composition and indices of abundance)~~with operating models simulating data assuming alternative levels of observation error, types and magnitude of process error, and magnitude of covariate effects on natural mortality. We fit estimating models to ~~simulated observations~~the simulated data with alternative assumptions ~~on inclusion of environmental~~regarding the inclusion of covariate effects, estimation of the median natural mortality rate, and ~~the source of temporal variability in the population demography~~. Our results suggest that estimation of environmental type of process error. We found estimating covariate effects on natural mortality ~~is possible and reliable even when the population process error source was misspecified in some~~required ideal scenarios with lower ~~uncertainty in covariate observations~~observation uncertainty, contrast in fishing pressure, and higher covariate ~~temporal variability~~variability, but the source of process error could be mis-specified. Estimation of spawning stock biomass and fishing mortality was also reliable in a wider range of scenarios even when estimating the median natural mortality rate.

keywords: state-space assessment models, time-varying natural mortality, bias, AIC

Introduction

State-space population models are now used widely for fisheries stock assessment in Europe, the United States, and Canada (Nielsen and Berg, 2014; Cadigan, 2016; Pedersen and Berg, 2017; Stock and Miller, 2021). Because application of these methods are considered best practice and recommended for the next generation of stock assessment models (Hoyle et al., 2022; Punt, 2023), it is expected their use will only grow globally. An appeal of state-space models lies in their separation of sources of biological and measurement variability by treating ~~latent~~ temporal variation in population characteristics as ~~statistical time series~~ latent stochastic processes with periodic observations measured with error. Through advances in computational capacity, we can use sophisticated numerical approaches to estimate ~~model parameters as mixed~~ parameters of these models as a combination of fixed and random effects (Thorson and Minto, 2015; Kristensen et al., 2016).

State-space stock assessment models ~~;~~ with non-linear functions of latent processes and numerous observation types with different probability ~~distribution assumptions~~ distributions, represent one of the most complex classes of state-space models. ~~The literature on~~ Our understanding of the effects of various factors on reliability of inferences from state-space assessment models is growing (Li et al., 2024; Miller et al., In reviewa). The importance of temporal contrast in population size and fishing mortality (F) and quality of data used to fit ~~assessment models including the state-space variety~~ both traditional and state-space assessment models is known (Magnusson and Hilborn, 2007; Miller et al., In reviewa). Furthermore, estimation of natural mortality (M), and even temporal variability in M , is possible in many scenarios (Lee et al., 2011; Cadigan, 2016; Miller and Hyun, 2018; Miller et al., In reviewa).

The effects of temporal variation in recruitment via undefined or explicit environmental factors have been extensively investigated in both traditional assessment models and ~~state space~~ state-space models (Myers, 1998; Haltuch and Punt, 2011; Johnson et al., 2016;

Miller et al., 2016). Reliability of estimating environmental and spawning biomass effects on recruitment in state-space assessment models requires a combination of strong effects, ~~good~~informative age composition data~~quality~~, contrast in the environmental covariate~~and lower recruitment variability (Britten et al., In review; Miller et al., In review)~~, and low variability of recruitment deviations (Britten et al., 2026; Miller et al., In review).

A critical aspect of fisheries assessment models and their use in management is short-term projections that are used to determine catch advice. While understanding drivers of recruitment is important, particularly for subsequent effects on reference points, recruitment in short-term projections typically has little impact on the exploitable biomass in the first few projection years. However, assumptions for M have immediate and larger effects on projected biomass because they affect the abundances at older age classes at the end of the data time series that constitute spawning biomass and catch (Brodziak et al., 2008; Stock et al., 2021).

Because of the effects of M on both biological reference points and ~~short term~~short-term projections, better understanding sources of variation in M would provide more accurate estimation of abundance and productivity and~~therefore~~, therefore, improved management. Temporal variation in M is less studied than recruitment, but its importance for explaining variability in observations has been demonstrated in state-space assessment models for Atlantic cod and yellowtail flounder (Cadigan, 2016; Stock et al., 2021). Deriso et al. (2008) also demonstrated the importance of several factors affecting M for Pacific herring.

Assessment models could include temporal variation in many aspects of population dynamics or how observations are related to the population. For example, the Woods Hole Assessment Model (WHAM) can include process errors treated as random effects for ~~transition in cohorts over time~~annual changes in cohort abundance (hereafter referred to as apparent survival), catchability for indices of abundance, selectivity of fishing fleets or indices, movement between regions, or in M (~~Stock and Miller, 2021; Miller et al., In review~~)

([Stock and Miller, 2021; Miller et al., 2025](#)). However, ~~misspecified temporal population~~
~~process errors~~ [mis-specifying the type of temporal variation](#) could lead to biased ~~population~~
[estimation of population size](#) and stock status ~~estimation~~, and, therefore, poor fisheries
management decisions (Legault and Palmer, 2016; Szuwalski et al., 2018). Studies of the
reliability of ~~inferences regarding the presence of~~ [identifying](#) temporal variability in M are
limited. Miller et al. (In reviewa) found AIC could accurately distinguish process errors
in apparent survival, but not for those specifically due to M except when uncertainty in
population observations (indices, catch, and age composition) was low and there was greater
temporal variation in M . ~~In their simulation studies looking at models with multiple sources~~
~~of process error, Li et al. (2024) found~~ [Li et al. \(2024\) found that](#) including more sources
of process error than existed in the operating model ([OM](#)) was a better model-building
approach than excluding them ~~a-priori~~ [a priori](#).

~~Here we conduct,~~ [we present results from](#) a simulation study with ~~operating models (OMs)~~
[OMs](#) varying by degree of ~~observation error~~ [observation error](#) uncertainty, sources of process
error, fishing history, temporal variation in environmental covariates, and magnitude of the
~~effect of the covariate~~ [covariate effect](#) on M . ~~The~~ [We fit the](#) simulated observations from
these OMs ~~are fitted~~ with estimating models ([EMs](#)) that make alternative assumptions for
sources of process error, and whether median M and covariate effects ~~are~~ [were](#) estimated.
We evaluated the effects of these factors on [1\)](#) convergence of fitted models, [2\)](#) whether
Akaike’s information criterion (AIC) can determine the correct source of process error and
correct assumption about covariate effects on M , ~~and~~; [3\)](#) [estimation bias for median \$M\$](#)
[and covariate effects; and 4\)](#) the accuracy for estimators of ~~relevant parameters and stock~~
~~size and harvest rates derived from the assessment model~~ [important terminal year \$M\$ and](#)
[spawning stock biomass \(SSB\)](#).

Methods

Our analyses used ~~the Woods Hole Assessment Model (WHAM)~~ WHAM to construct both OMs and EMs (~~Miller and Stock, 2020; Stock and Miller, 2021; Miller et al., In reviewb~~). ~~The WHAM package~~ (Miller and Stock, 2020; Stock and Miller, 2021; Miller et al., 2025). WHAM has been used extensively to configure OMs and EMs for several other simulation studies (~~Stock et al., 2021; Legault et al., 2023; Li et al., 2024; Britten et al., In review; Li et al., In review~~) (Stock et al., 2021; Legault et al., 2023; Li et al., 2024; Britten et al., 2026; Li et al., 2026a) and is used to assess many commercially important stocks in the Northeast U.S. (e.g., NEFSC, 2022a,b, 2024). We used version 1.0.6.9000 ~~;~~ (commit 77bbd94) to generate all results.

We completed a simulation study with 288 ~~operating models~~ OMs. The factors defining the configuration of each ~~operating model~~ OM, described in detail in subsequent sections, include source of ~~population~~-process error (3 levels), index and catch observation uncertainty (2 levels), environmental covariate uncertainty (2 levels), temporal variation in the latent environmental covariate (4 levels), size of the covariate effect on M (3 levels), and fishing history (2 levels). We simulated 100 data sets for each ~~operating model~~ OM that included simulations of process errors.

For each simulated data set we fit a set of 12 ~~estimating models (EMs)~~ EMs. The factors that distinguish the ~~estimating models~~ EMs, also described in detail below, include source of population process error type (3 levels) whether ~~(median)~~ median M was estimated or assumed known (2 levels), and whether the ~~effect of the environmental covariate~~ environmental covariate effect on M was estimated or not (2 levels).

The sources of population process error that were used in the OMs or assumed in the EMs were on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and M (R+M). We did not use the log-normal bias-correction feature for process errors or observations described by Stock and Miller (2021) for OMs and EMs (~~Li et al., In reviewb~~).

Using this correction with random effects and estimation of fixed effects via marginal likelihood can result in substantial bias in assessment output such as recruitment and SSB (Li et al., 2026b). Simulations were all carried out on the University of Massachusetts Green High-Performance Computing Cluster. Code for completing the simulations and summarizing results can be found at https://github.com/timjmiller/SSRTWG/ecov_study/mortality.

Operating models

Environmental covariate

In ~~the WHAM model~~ WHAM, environmental covariates are ~~assumed to be described~~ modeled as state-space processes with annual observations of the true latent covariate (Miller et al., 2016; Stock and Miller, 2021). In our simulations, ~~we assumed~~ the latent covariate ~~is assumed~~ to be a stationary first order autoregressive (AR1) process.

$$X_y | X_{y-1} \sim N \left(\mu_E (1 - \rho_E) + \rho_E X_{y-1}, (1 - \rho_E^2) \sigma_E^2 \right),$$

with marginal mean $\mu_E = 0$ and variance σ_E^2 . The four configurations of the latent environmental covariate in the ~~operating models~~ OMs assume one of two values for the marginal standard deviation $\sigma_E \in \{0.1, 0.5\}$ and for the autocorrelation parameter $\rho_E \in \{0, 0.5\}$.

~~The WHAM treats the~~ observations of the latent environmental covariate ~~are assumed to be unbiased and Gaussian~~ (x_y) as unbiased and normal conditional on the true covariate X_y .

$$x_y | X_y \sim N \left(X_y, \sigma_e^2 \right).$$

The standard deviation of the environmental observations in the ~~operating models is~~ OMs ~~was~~ one of two values $\sigma_e \in \{0.1, 0.5\}$. Figure S1 provides example simulations of the latent covariate and observations under the alternative configurations.

Population

Many of the characteristics of the population biology and structure including the age classes (10 age classes (ages: 1 to 10+)), time span (40 years), maturity (Figure S2, top left), growth (Figure S2, top right), time of spawning (1/4 of the year), and recruitment (Figure S2, bottom right) ~~are~~ were identical to Miller et al. (In reviewa). The maturity at age ~~is~~ was a logistic function with age at 50% maturity (a_{50}) = 2.89 and slope = 0.88 and weight at age ~~is~~ was derived from a von Beralanffy growth function where $t_0 = 0$, $L_\infty = 85$, and $k = 0.3$, and a ~~length-weight relationship~~ length-weight relationship

$$W_a = \theta_1 L_a^{\theta_2},$$

where $\theta_1 = e^{-12.1}$ and $\theta_2 = 3.2$.

~~The~~ In WHAM, the general model for M in year y is a log-linear function of both covariate effects X_y and process errors $\varepsilon_{M,y}$ and a parameter β_M that defines median M .

$$\log M_y = \beta_M + \beta_E X_y + \varepsilon_{M,y},$$

where the process errors are modeled as random effects that may, in general, be autocorrelated ~~normal random variables~~ Gaussian random variables.

$$\varepsilon_{M,y} | \varepsilon_{M,y-1} \sim N(\varepsilon_{M,y-1}, (1 - \rho_M^2) \sigma_M^2)$$

(Stock and Miller, 2021), but we assumed ~~ρ_M~~ $\rho_M = 0$ in our R+M OMs. We assumed ~~the~~ the median ~~M~~ M ~~rate~~ $e^{\beta_M} = 0.2$ ~~is~~ was constant across ages. For R and R+S OMs and EMs, $\varepsilon_{M,y} = 0$. For all R+M OMs, we ~~assume the same~~ assumed the standard deviation $\sigma_M = 0.3$, which ~~is~~ was estimated in the R+M EMs. The covariate effect ~~is~~ was one of 3 alternative values in the ~~operating models~~ OMs, $\beta_E \in \{0, 0.25, 0.5\}$. The parameters defining the simulated

covariate time series, size of the covariate effect, and any M random effects resulted in a range of different levels of variation in annual M values (Figure S3).

We assumed expected recruitment each year ~~is from a Beverton-Holt~~ was from a Beverton-Holt stock-recruit relationship (SRR)

$$R_y = \frac{aSSB_{y-1}}{1 + bSSB_{y-1}}.$$

All biological inputs to calculations of ~~spawning biomass~~ SSB per recruit (i.e., weight, maturity, and M at age) ~~are were~~ constant in the R and R+S OM without covariate effects on M . Therefore, steepness and equilibrium unfished recruitment ~~are were~~ also constant over the time period for those OM (Miller and Brooks, 2021). As in Miller et al. (In review), our assumed biological inputs and selectivity (defined below) with constant M ~~result in~~ resulted in the equilibrium F that reduces ~~spawning biomass~~ SSB per recruit to 40% of the unfished level ~~is being~~ $F_{40\%} = 0.348$. With an assumed unfished recruitment of $R_0 = e^{10}$, setting $F_{MSY} = F_{40\%}$ results in a steepness of 0.69 and $a = 0.60$ and $b = 2.4 \times 10^{-5}$. For R+M OM and all OM with covariate effects on M , steepness ~~is not constant~~, was not constant but we used the same a and b parameters as other ~~operating models OM~~ which equates to a steepness and R_0 at the median (or mean on log-scale) of the time series models for M and the covariate.

We also used the same two fishing scenarios as Miller et al. (In review) for OM. In the first scenario, the stock ~~experiences~~ experienced overfishing at $2.5F_{MSY}$ for the first 20 years followed by fishing at F_{MSY} for the last 20 years (denoted $2.5F_{MSY} \rightarrow F_{MSY}$). In the second scenario, the stock ~~is was~~ fished at F_{MSY} for the entire time period (40 years). The magnitude of the overfishing assumptions ~~is was~~ intended to reflect estimates of overfishing for Northeast US groundfish stocks from Wiedenmann et al. (2019).

We configured all R, R+S, and R+M OM with uncorrelated random effects on recruitment with standard deviation on $\log(\text{recruitment})$ $\sigma_R = 0.5$. ~~This same assumption was used~~

~~by Miller et al. (In review)~~ Miller et al. (In review) used this same assumption for R+M
OMs and other OMs with fishery selectivity and index catchability process errors. For R+S
OMs, apparent survival process errors were uncorrelated with $\sigma_{2+} = 0.3$. These standard
deviations were on the lower end of the range of values estimated for groundfish stocks in
the NE US (see Table S1 in Britten et al., 2026).

Catch and index observations

We defined the generation of observations of aggregate (total combined across ages) catch and
indices ~~;~~ and corresponding age composition identical to Miller et al. (In review). There ~~is~~
~~was~~ a single fleet operating year round for catch observations with logistic selectivity ~~for the~~
~~fleet defined~~ with $a_{50} = 5$ and slope = 1 (Figure S2, bottom left). Observations ~~are were~~
generated for all 40 years of the model. There ~~are were~~ two index time series intended to
represent fishery-independent surveys occurring in the spring (0.25 way through the year)
and the fall (0.75 way through the year). ~~Catchability~~ We assumed catchability of both
surveys ~~are assumed to be~~ to be 0.3.

WHAM treats the aggregate catch and index observations as log-normal and we assumed
the standard deviation parameter was 0.1 for all annual catch observations. We assumed
catch and index ~~age composition observations are~~ age composition observations were gener-
ated from a logistic-normal distribution where errors on the multivariate normal scale ~~are~~
~~independent were independent~~ (Aitchison and Shen, 1980; Stock and Miller, 2021; Trijoulet et al., 2023)
. The standard deviation parameter ~~is~~ was also constant across ages.

~~Standard deviation for log-aggregate catch was 0.1.~~ There were two levels of observation
~~error variance uncertainty~~ for aggregate indices and age composition for both indices and fleet
catch. A ~~low uncertainty low uncertainty~~ specification assumed standard deviation ~~of both~~
~~series of log-aggregate index observations was~~ parameter equal to 0.1 for all annual aggregate
index observations and the standard deviation of the logistic-normal for ~~age composition all~~

age-composition observations was 0.3. In the ~~high-uncertainty~~ high-uncertainty specification the standard deviation ~~for log-aggregate indices~~ parameter for the annual aggregate index observations was 0.4 and that for the ~~age-composition~~ age-composition observations was 1.5. ~~For all estimating models, the standard deviation for log-aggregate observations was assumed known at the true value whereas that for the logistic-normal age-composition observations was estimated.~~

Estimating models

~~Estimating models were fit to each of 100 simulated data sets from each operating model.~~ There were three factors defining the configuration of each ~~estimating model~~ EM: 1) whether β_M was estimated or assumed known; 2) whether an environmental effect β_E was estimated or not; and 3) whether the process errors were assumed on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and M (R+M).

~~We fit each EM to each of 100 simulated data sets from each OM.~~ The configuration of the process errors in the ~~estimating models~~ EMs generally matched the corresponding options in the ~~operating models~~ OMs. For example, ~~uncorrelated~~ Random effects were assumed uncorrelated for R EMs and R OMs and for R+S ~~was assumed for both the estimating and operating model~~ EMs and R+S OMs. However, R+M EMs did not assume M random effects were uncorrelated (ρ_M was estimated) although they were simulated without correlation in the R+M OMs. The environmental covariate observations were included in all ~~estimation models~~ EMs, whether effects on M were estimated or not, to ensure comparability of AIC. All ~~fixed-effects~~ fixed-effects parameters for selectivity, catchability, fully-selected F , mean recruitment, initial abundance at age, and ~~variances~~ standard deviations for logistic-normal ~~age composition~~ age-composition distributions were estimated. Any ~~process error~~ process-error variance parameters for recruitment, apparent survival, and M were also estimated. ~~The observation error variance of the environmental observations and aggregate catch and indices~~

~~were all~~ For all EMs, the standard deviation parameters for aggregate catch, index, and environmental covariates were assumed known at the true ~~values~~ value whereas that for the logistic-normal age-composition observations was estimated.

Measures of reliability

EM convergence

We measured the frequency of convergence when fitting each EM to the simulated data sets for each OM. There are various ways to assess convergence of the fit (e.g., Carvalho et al., 2021) ~~;~~ but we defined successful convergence as the Hessian of the marginal log-likelihood being invertible and providing variance estimates for the fixed effects parameters as recommended by Miller et al. (In review). ~~We calculated 95% confidence intervals (CIs) for probability of convergence using the Clopper-Pearson exact method (Clopper and Pearson, 1934; Thulin, 2014).~~

AIC for model selection

We measured the frequency of correct model selection using marginal AIC~~.~~, known to be appropriate for mixed-effects models specifically when the analyst is interested in fixed effects rather than random effects (Vaida and Blanchard, 2005). Other studies have shown the utility of marginal AIC in distinguishing process error and other model structure for state-space assessment models applied to a particular stock (Miller and Hyun, 2018; Stock and Miller, 2021).

For a given ~~operating model~~ OM, the set of ~~models that were considered~~ EMs that were compared all made the same ~~assumptions on whether or not to estimate~~ assumption about β_M ~~or it is assumed~~ (estimated or treated as known at the true value. ~~For model~~). For EM m , the marginal AIC is a function of the marginal log-likelihood maximized with respect

to the estimated fixed effects in the ~~model-EM~~ θ and the number of estimated fixed effects
 $n(\theta)$ ~~estimated~~,

$$\text{AIC}_m = -2 [\text{argmax}_{\theta} \log L_m(\theta) - n(\theta)].$$

All ~~model-EM~~ fits that successfully completed the optimization were used for this set of
 analyses. We ~~used all of these fits~~ did not condition on convergence as defined above because
 some lack of convergence would be expected for the correct behavior of more complicated
 models that include process errors that did not exist in the ~~operating-model~~ OM. For example
 R+M EMs fit to R OMs would be expected to estimate no variance in the M random effects
 and the estimated variance parameter going to zero would cause poor convergence ~~—but~~
have the same marginal log-likelihood (with more fixed effects) and therefore higher AIC
as expected. The correct EM made the correct assumption for the source of process errors
(R, R+S, R+M) and either included a covariate effect on M when an effect was simulated
($\beta_E \in \{0.25, 0.5\}$) or did not include an effect when one was not simulated ($\beta_E = 0$).

Parameter estimation bias and accuracy

All results here used OM simulations with fits that satisfied the Hessian-based convergence
 criterion described above. We used this conditioning to reflect how practitioners would
 proceed in analyses of model fits with real assessment data. ~~That is, practitioners would~~
~~ensure models converged such that Hessian-based standard errors were available for all model~~
~~parameter estimates.~~

We focused on statistical behavior of estimators of the covariate effect on M ($\hat{\beta}_E$), ~~the~~
~~estimator of the~~ the median M parameter ($\hat{\beta}_M$), ~~and estimators of~~ and terminal year
 M ($\hat{M}$), ~~spawning stock biomass (SSB)~~ SSB, and fully-selected ~~F~~ ($\hat{F}$ fishing mortality (F)).
 In preliminary analyses we examined results for the estimators of all the annual values for M ,
 SSB and F over the whole time series, but we found no appreciable differences in patterns
 across the various factors defining the OM and EMs. Furthermore, because results for

terminal year F were generally inversely related to those for ~~spawning stock biomass, and~~
~~are SSB, they are not discussed further but~~ provided in the Supplementary Materials ~~and not~~
~~discussed further.~~

We calculated ~~median errors (ME) errors~~ of $\hat{\beta}_E$ and $\hat{\beta}_M$, and the Hessian-based standard
error estimators of these parameters ($\widehat{SE}(\hat{\beta}_E)$ and $\widehat{SE}(\hat{\beta}_M)$) ~~. We calculated the median~~
~~relative errors (MRE) of terminal year $\hat{M}$, SSB, and $\hat{F}$. We as~~

$$\delta_i = \hat{\theta}_i - \theta_i$$

where θ_i is the true value of a given parameter or attribute for simulation i and $\hat{\theta}_i$ is the
estimate from an EM fitted to data from simulation i . We defined the true standard error as
the standard deviation of estimates across converged fits of a given EM and OM combination.

We calculated the relative errors for terminal year M , SSB, and F as

$$\text{RE}_i(\theta_i) = \frac{\hat{\theta}_i - \theta_i}{\theta_i}. \quad (1)$$

To measure accuracy we also calculated the root mean square error (RMSE) ~~and estimated~~
~~probability of coverage of constructed,~~

$$\text{RMSE}(\hat{\theta}) = \sqrt{\frac{1}{n} \sum (\hat{\theta}_i - \theta_i)^2},$$

where n is the number of simulations with converged fits of a given EM and OM. The true
values for terminal year SSB, and in some OM M (R+M OM and any OM with covariate
effects on M), varied among simulations. Finally, we calculated 95% ~~CIs~~ confidence intervals
for $\hat{\beta}_E$ and $\hat{\beta}_M$ ~~for using~~ Hessian-based standard error estimates for converged EMs that
estimated these parameters ~~. We constructed the CIs for probabilities of CI coverage using~~

~~the same methods as those for probabilities of convergence.~~ in each OM simulation and recorded whether they included the true value.

~~The true values for terminal year SSB and F vary among simulations. For the i th simulated data the relative error~~ Estimation of terminal year M could appear to be unbiased even for EMs that assumed no temporal variation in M because the assumed or estimated β_M parameter can be equal or near the median of the true terminal year values. This centering of the true values at the constant value of M in the EM happens because the mean of the M process error and covariate distributions were assumed equal to 0. Therefore, we calculated the correlation of estimates and simulated values for terminal year ~~value θ_i provided from the fitted estimation model is~~

$$\text{RE}_i(\theta) = \frac{\hat{\theta}_i - \theta_i}{\theta_i}$$

~~We calculated RMSE as~~

$$\text{RMSE}(\hat{\theta}) = \sqrt{\frac{1}{n} \sum^n (\hat{\theta}_i - \theta_i)^2}$$

M for a given EM and OM combination,

$$\text{Cor}(\hat{M}, M) = \frac{\text{Cov}(\hat{M}, M)}{\text{SD}(\hat{M}) \text{SD}(M)}.$$

~~For ME and MRE of estimators, we constructed 95% CIs using the binomial distribution approach as in Stock and Miller (2021) and Miller et al. (In review). Because of the inverse relationship of the CI and the null hypothesis significance test, when the CI contains zero we do not have evidence of bias from the simulation study.~~ For R+M OMs and any other OMs where there were covariate effects on M , high correlation (approaching 1) along with median relative errors close to 0 indicates ability of the EM to estimate annual M . If correlation and median relative errors are both near 0, this indicates no ability to estimate annual M . Note also that the correlation is undefined when either the estimated or true value are constant across simulations because the corresponding standard deviations are 0. This applies to R

and R+S EMs that assumed β_M to be known and $\beta_E = 0$, and R and to R+S OM where $\beta_E = 0$. We also did not use calculated correlations for EMs when there were only two converged fits because they are uninformative. Because true terminal year SSB values also varied across simulations we made the same correlation calculations for them as well.

Summarizing results across OM and EM attributes

~~Miller et al. (In reviewa) found inferences are most reliable for distinguishing process error sources and stock-recruit relationships in scenarios with lower observation error for indices and age composition data and with temporal contrast in fishing pressure. Our expectation is that inferences for covariate effects on~~ There were numerous OM and EM attributes across all simulation scenarios and, like Miller et al. (In reviewa), we used two methods to summarize the major factors explaining differences in results for each measure of reliability. First, we fit regression models with the responses as the measures of reliability described above and the predictor variables were defined based on OM and EM attributes (e.g., MacKinnon et al., 1995; Wang et al., 2017; Harwell et al., 2018). We used logistic regression for binary indicators of convergence and AIC-based selection of the appropriate assumption of the covariate effect on M ~~would require data~~. For indicators of AIC-based selection of EM process error source (multiple categories), we performed multinomial regressions. For other measures of reliability we fit ordinary linear regression models. Relative errors of terminal year M , SSB, and F (Eq. 1) are bounded below at -1, so we used a transformation of these values

$$f(\hat{\theta}_i, \theta_i) = \log \left[\text{RE}(\hat{\theta}_i, \theta_i) + 1 \right]. \quad (2)$$

We omitted fits where estimated terminal year SSB, M , or F was equal to zero because of the undefined transformation. RMSE is bounded below at 0 so we used a log transformation for responses in corresponding regressions. We did not use any transformation for the correlation

of estimated and true values for terminal year M and SSB. Note there was only one RMSE and correlation calculated for an EM from all simulation fits that converged from a given OM scenario. For all regressions we fit separate models with just individual OM and EM factors included, all factors combined, with all second order interactions, and with all third order interactions. For the multinomial regression, we used the `vglm` function from the VGAM package (Yee, 2008, 2015). We calculated percent reduction of residual deviance (from the null model with no factors) for each of the regression fits. As Miller et al. (In reviewa) noted, there is a lack of independence of the results for each EM fitted to the same simulated data set of a given OM and therefore we did not perform formal statistical analyses of effects of OM and EM attributes.

For the second summarization method, we used classification and regression trees (Breiman et al., 1984) to illustrate the primary OM and EM attributes and interactions that partition the values for each measure of reliability (e.g., Gonzalez et al., 2018; Collier et al., 2022). We used classification trees for categorical measures (convergence and AIC model selection) and regression trees for the other measures with continuous scales (errors, relative error, and RMSE). The response variables were the same as the regressions for the deviance reduction analyses. We used the `rpart` function in the `rpart` package (Therneau and Atkinson, 2025) to fit trees. Full trees were determined using default settings except that we increased the number of cross-validations to be at least that informative and inspection of the comprehensive set of results across all OMsgenerally confirms this (See Supplemental Materials). Therefore, we restrict our attention to these OM scenarios with more informative data (temporal contrast in fishing and lower population observation error). 100. For one classification tree fitted to AIC selection of process error configuration results for R+S OMs, we had to make a small changes to the default prior probabilities (the proportions of each class in the data) used in the criterion on whether to allow branches from a given node because the default priors would not produce any branches. For clarity, we manually pruned the full trees to show just the primary branches.

All OM and EM factors we used to examine reductions in deviance and regression trees are given in Table S1. We provide more detailed results for each EM and OM configuration in the Supplementary Materials using similar methods to Miller et al. (In review). The detailed results present probability of convergence, probability of each EM having lowest AIC, median error and RMSE of β_M and β_E and corresponding standard error estimates, and median relative error and RMSE for terminal year SSB, F , and M . There are also detailed results for correlation of estimates and true values for terminal year M and SSB.

Results

EM Convergence

~~In OMs with contrast in fishing pressure and lower uncertainty in population observations (catch, indices and age composition), R EMs generally converged with high frequency except when EMs estimated covariate effects on M and the covariate uncertainty (σ_e) was high (Figure ??).~~ For fitted logistic regression models to indicators of convergence, the EM process error assumption provided the largest percent reduction in deviance (9-25%) of all OM and EM attributes for all three OM process error types (Table 1). Including second order interactions of OM and EM factors also provided further reduction of residual deviance (between 7 and 11%), indicating successful convergence depended on a combination of OM and EM attributes.

Classification trees for each OM process error source all had the primary branches defined using the attribute (EM process error assumption) that also provided the largest reduction in deviance (Figure 1). Across all fits for each OM process error configuration, convergence was best for R+S ~~EMs generally converged with low frequency when the OM process error did not match the EM configuration.~~ OMs, but better convergence occurred when the process error was correct for R and R+S OMs. For R+M ~~EMs generally did not converge with~~

high frequency except for OMs, R EMs converged well (88%), but good convergence of R+S OMs with low covariate uncertainty and where the M EMs occurred for any true process error source if the OMs also had constant fishing pressure and EMs assumed median M as known. There was generally little effect of the treatment of median parameter (β_M) to be known. All branches based on observation error of covariates or other assessment data (indices and age composition) showed better convergence with more precise observations. Better convergence also occurred when EMs assumed no covariate effects (regardless of whether the OM simulated them), when EMs assumed median M (known or estimated) on convergence for any of the other EMs. The largest impact of the size of the was known, and when OMs assumed greater temporal variability in the true environmental covariate.

AIC performance

Process error source selection

For all OM process error types, the level of error in both the population and covariate observations were the attributes that resulted in the largest percent reductions in deviance for multinomial regression fits to indicators of AIC selection of the process error configuration (Table 2). For R+S and R+M OMs, level of error in population observations provided deviance reduction (19% and 13%, respectively) much larger than any other OM and EM factors whereas error in covariate observations provided the largest reduction (8%) for R OMs. Lesser deviance reduction (2-3%) occurred for the true covariate effect (β_E) on convergence was observed for R and size and whether the EM made the correct assumption about including a covariate effect for R OMs and error in covariate observations for R+M EMs that estimated covariate effects in OMs. Including second and third order interactions, provided the largest reductions in residual deviance for R OMs (24%) and little reduction for R and R+M OMs with high uncertainty in covariate observations (σ_e) and high temporal variability in the latent covariate (σ_E). There was an increase in convergence frequency with

~~increased temporal variability in the covariate for many EMs in OMs with large covariate uncertainty~~ S and R+M OMs (4% and 6%, respectively).

The attributes defining the primary branches of classification trees matched those that provided the largest reductions in deviance for all OM process error types (Figure 2). AIC performed poorly in selecting the correct process error for R+M OMs where lowest AIC occurred for the simpler R EMs. However, accuracy of AIC-based selection was high for R and R+S OMs. For R+S OMs, mis-classification occurred to R EMs with the lowest AIC when OM had higher error in population observations. For R OM, there was a small decrease in accuracy when the OM included the largest covariate effect on M , but in those scenarios, AIC was more accurate for the process error assumption when the EMs also incorrectly assumed no covariate effect.

~~AIC performance~~

Covariate effect selection

~~We only present results for EMs where the median M rate parameter was estimated because results differed little whether this parameter was estimated or assumed known (see Figures S13 to S15)~~ The factors explaining the largest reduction in deviance for logistic regression models fitted to indicators of AIC selection of the correct covariate effect assumption depended on the magnitude of the true effect assumed in the OM (Table 3). When ~~there was temporal contrast in fishing pressure and lower uncertainty~~ OM had no covariate effect ($\beta_E = 0$) the OM process error type provided the largest reduction in deviance (4%). When OM assumed a covariate effect ($\beta_E > 0$), level of error in population observations, the only OM scenarios where AIC was consistently accurate for both covariate effect and process error configurations were R OM with high variability in the covariate and low uncertainty in observations of the covariate (Figure ??). R+M OM with those same covariate attributes were only accurate and magnitude of temporal variability in the true

covariate provided the largest reductions in deviance (4% and 8-21%, respectively). The reduction in deviance provided by the magnitude of covariate temporal variability also increased with larger true covariate effect. Including second and third order interactions provided a modest reduction in deviance (5-7%).

The factors defining the primary branches of classification trees for indicators of AIC selecting the correct covariate effect assumption matched the factors providing the largest deviance reductions (Figure 3). Accuracy of AIC-based model selection for the covariate effect (assumption was high overall for OMs with no covariate effect (88%) (low Type I and II errors) and the incorrect process error almost always chosen was R rather than R+S. We also observed high accuracy for both covariate effect error) and low when there was a covariate effect (23% and 35% for OMs with $\beta_E = 0.25$ and process error configurations for 0.5, respectively) (high Type II error), but higher accuracy occurred for certain combinations of OM attributes. When the true covariate effect was greatest ($\beta_E = 0.5$), AIC accuracy was 91% for EMs fit to OMs with high temporal variability in the covariate and low error in population and covariate observations regardless of whether the EM process error was correct. However, when the covariate effect was weaker ($\beta_E = 0.25$), high accuracy only occurred for a subset of those OMs with R or R+S OMs where either no effect of the covariate or the strongest effect of the covariate was simulated. All R and R+S OMs were accurate for process error configuration and across all OMs, AIC accurately selects models without covariate effects when there are none (low Type I error). See Figures S13 to S15 for full results.

M process errors (76%), and the further conditioning on OMs with high autocorrelation of the covariate provided higher accuracy (86%). When there was temporal contrast in fishing pressure, lower uncertainty no covariate effect, AIC accuracy was best for R OMs (95%), and for R+S and R+M OMs, accuracy was improved if the OMs also had lower error in population observations, low uncertainty in environmental observations, and larger temporal contrast in the simulated true environmental covariate, we observed little or no

~~bias in estimation of covariate effects ($\hat{\beta}_E$) across all EM and OM process error assumptions,~~
~~both EM configurations of median M , and all true covariate effect sizes (Figure ??, rows 3~~
~~and 4) . The only exception to this was when R EMs with median M estimated were fit to~~
~~and the EM process error assumption was correct (85%).~~

Covariate effect inferences

Estimation bias

For regression models fit to errors in estimation of the covariate effect β_E , we found very little
 reduction in deviance for any of the OM and EM factors (Table 4). For the normal model in
 these regressions, deviance is the sum of squares and there were extremely large estimates
 of β_E (and errors) for many simulations which resulted in extremely large sums of squares
 and relative differences from the null model were therefore small (<1%). Nevertheless,
 observation error of the covariate provided either the largest or second largest reductions
 in deviance of any of the single OM and EM attributes for OM process error configuration
 types. OM fishing history provided reductions of similar scale for R and R+S OM~~s~~~~with those~~
~~configurations. Decreasing trends in bias of $\hat{\beta}_E$ for EMs fit to many configurations of R and,~~
 Other factors providing similar reductions were true covariate effect size for R OM~~s~~ and level
 of error in population observations and EM process error assumption for R+~~M~~ OM~~s~~~~indicate~~
~~that~~ S OM~~s~~. Including second and third order interactions also provided further reductions
 in residual deviance similar in scale to the individual factors.

Median errors for $\hat{\beta}_E$ ~~values were closer~~ across all simulations for each OM process error
 configuration were negative but close to 0 (-0.1 to ~~zero on average even when the true effect~~
~~increased -0.07~~) and primary branches were based on factors that matched those providing
 the largest reductions in deviance (Figure 4). Branches based on level of error in population
 or covariate observations provided better median estimation error (closer to 0) with lower
 uncertainty. Branches based on level of covariate temporal variation provided better median

estimation error with higher temporal variation. For R OMs with the largest true covariate effect, estimation error was poor unless there was also high temporal variation and low observation error for the covariate. For R+S OMs, ~~EMs with the matching process error configuration showed little evidence of bias in estimating~~ we also found lower estimation error when the EM process error assumption matched. We observed low median error across all simulations for R+M OMs with low error in covariate observations. The detailed results for all OM and EM configurations demonstrate that the median errors get worse with increased values for the true β_E ~~across a range of covariate and covariate observation configurations~~. for many scenarios without low error in population and covariate observations and contrast in fishing pressure, indicating that the estimates remain close to zero when the true $\beta_E > 0$ (Figures S16 to S18). The detailed results also indicated little bias in covariate effect estimation regardless of the EM process error configuration when OMs had low population and covariate observation error and contrast in fishing pressure.

Standard error estimation bias

Factors providing the largest reductions in deviance for errors in estimation of the standard error of the covariate effect size ($\widehat{SE}(\hat{\beta}_E)$) were generally similar to those that were most important for estimation of β_E , but deviance reductions were one or two orders of magnitude larger (Table S2). Level of error in covariate observations and EM process error assumption was important for all OM process error configurations (6-13% and 3-4%, respectively). Level of error in population observations provided relatively large reductions in deviance for R+S and R+M ~~EMs generally performed similarly when fitted to either R+S or R+M OMs except R+S OMs with high covariate uncertainty and low covariate variability (Figure ??, rows 5 and 6 of middle columns)~~ OMs (3% and 8%, respectively). The OM fishing pressure history provided a lesser reduction in deviance for all OM process error types (2-4%). Like estimation of β_E , additional large reductions in deviance also occurred when second and third order interactions of OM and EM attributes were included in the regression models

(11-31%).

~~Bias of Hessian-based standard error estimation~~ Median errors in estimation of the standard error for $\hat{\beta}_E$ ~~was also close to zero in the~~ across all simulations for a given OM process error configuration were strongly negative (-0.8 to -0.6), but combinations of OM and EM attributes resulted in improved values closer to zero (Figure S4). For all OM process error types, branches with lower error in covariate observations provided better median errors in the standard error estimator (closer to 0).

Examining the detailed results for each OM and EM configuration shows little bias of standard error estimation in the same OM scenarios where ~~there was little~~ bias in estimation of β_E ~~itself~~ was close to zero (~~Figure ??, rows 3 and 4~~ Figures S19 to S21). However, in these high information OMs, bias of standard error estimation was observed for several EMs that did not have the matching R+S and R+M process error configuration. ~~Estimation of standard errors was essentially unreliable for most EMs when OMs had higher uncertainty in covariate observations (Figure ??, rows 5 to~~

Confidence interval bias

For logistic regressions of indicators of confidence intervals including the true value of β_E , factors that provided the largest reductions in deviance were true covariate effect size for R and R+M OMs (11% and 3%, respectively) and EM process error configuration for R+S OMs (7%) (Table 5). Covariate observation error also provided relatively large reduction in deviance for R OMs (8) ~~—~~%. Lesser reductions of 1-2% occurred for level of error in population observations and covariate temporal variability for all OM process error types. Including second and third order interactions also provided relatively large further reductions in deviance (5-9%).

~~Despite the reliable estimation of~~ Over all simulations, rates of 95% confidence interval coverage indicated negative bias with rates ranging 85-88% for the different OM process

error configurations (Figure 5). Factors defining the primary branches matched the factors that made the largest reductions in deviance for fitted logistic regression models. For R OMs, the primary branches based on error in covariate observations indicated higher rates of confidence interval coverage with lower error, but when that error was higher, higher rates were associated with OMs where no covariate effect was assumed ($\beta_E = 0$). For R OMs with some covariate effect and higher error in covariate effects and standard errors with low uncertainty in environmental observations, and larger temporal contrast in the simulated true environmental covariate, we found poor CI coverage for many EMs fitted to observations, higher rates of coverage occurred with higher error in the population observations. For R+S OMs, higher rates of confidence interval coverage were observed for EMs that assumed R+S and/or R+M OMs (Figure ??, rows 3 and 4). For example, process errors.

The more detailed results for each OM and EM configuration show that good confidence interval coverage can be obtained for all of the investigated covariate effect sizes when there was low error in covariate and population observations, high covariate temporal variability and contrast in fishing pressure for R and R+S OMs. The EM process error configuration was not a factor for the R OMs, but R+M EMs or R+M EM configurations were necessary for R+S OMs (Figures S22 to S23). However, confidence interval coverage was negatively biased for all EM process error configurations fit to R+M OMs showed little bias with these conditions even though little bias was observed in estimation of the covariate effect and standard errors, but CI coverage was negatively biased (rows 3 and 4 in right two columns of Figures ??, ??, and ??). However β_E or its standard error (Figure S24). Examining the estimates for each simulated data set when R+M EMs were fit to R+M OMs where we observed little estimation bias, there appears to be a positive correlation of the covariate and standard error estimates such that estimates less than the true value had negatively biased standard error estimates which would make CIs confidence intervals too small (Figure S29). CI coverage was most reliable for R+S OMs when the EMs had the matching process error

configuration (Figure ??, middle 2 columns).

Median natural mortality rate inferences

We observed negligible bias in estimating the median

Estimation bias

Across all OM process error types, regression models fit to errors for median M parameter (β_M) over a much wider set of operating models than that for estimating the covariate effect on M . When there is temporal contrast parameter estimation ($\hat{\beta}_M$) showed largest reductions in deviance from the type of OM fishing pressure (4-17%), EM process error assumption (2-12%), and the EM covariate effect assumption (3-7%, Table 6). The level of observation error also provided a similar reduction in deviance for R OMs (3%). Relatively large reductions in deviance also occurred with second and third order interactions (7-16%).

Like $\hat{\beta}_E$, median errors for $\hat{\beta}_M$ were negative but close to 0 (-0.1 to -0.07) across all simulations for each OM process error configuration (Figure 6). The primary branches for all OM process error types were defined by the type of OM fishing pressure, with median errors closer to zero when there was a change in fishing pressure, lower uncertainty for R+S and R+M OMs. Branches based on the level of error in population observations, all EMs fit to R showed median errors closer to zero with more precise observations. For R+S and R+M OMs showed little evidence of bias for, branches that included the matching EM process error assumption provided better median errors.

Detailed results reveal that bias of $\hat{\beta}_M$ (Figure ??, left and right sets of columns). For was negligible for all EM process error assumptions fitted to R and R+S OMs with those conditions, only EMs with the matching M OMs when there was contrast in fishing pressure and low error in population observations, but the matching EM process error assumption

~~showed little evidence of~~ was also important for EMs fit to R+S OMs (Figures S30 to S32). For those OM, any attributes related to covariate process error and effects on M had little effect on bias.

~~We also found little or no bias of Hessian-based standard error estimation for β_M in many of the same OM scenarios where bias~~

Standard error estimation bias

The EM process error assumption provided the largest reduction in deviance for regression models fitted to errors in estimation of the standard errors of $\hat{\beta}_M$ ~~was close to zero except for some EMs with the process error source mis-specified or where the OM simulated the strongest covariate effect (Figure ??) . Like $\hat{\beta}_M$, reliability of standard error estimation in~~ from R (3%) and R+M (7%) OM simulations (Table S3). The type of OM fishing pressure provided the largest reduction (13%) for R+S OM and a lesser reduction in deviance for R+M OM (4%). The EM covariate effect assumption also provided similar reductions in deviance for R+S (9%) and R+M (4%) OM.

The factors defining the primary branches of regression trees matched those providing the largest reductions in deviance for errors in $\widehat{SE}(\hat{\beta}_M)$, but median errors indicated strong negative bias across all OM process error types (Figure S5). Branches with lower error in population observations provided better median errors (closer to 0). For R+S OM, ~~required the EMs to have the matching~~, branches where the EM process error assumption was correct provided better median errors.

~~we found CI coverage to be unreliable for many EM-OM combinations even for Detailed results for each OM and EM combinations where bias in $\hat{\beta}_M$ and its standard error estimators was negligible, much like that for $\hat{\beta}_E$ (Figure ??). In the same EM-OM combination we investigated above for $\hat{\beta}_E$, we observed an opposite negative correlation of $\hat{\beta}_M$ and its standard error estimates (Figure S39) , which would result in CIs being too narrow when~~

~~$\hat{\beta}_M$ values are larger than average. The exceptions to the general poor CI coverage were~~
~~configuration indicate little bias in certain scenarios where OMs had temporal contrast in~~
~~fishing pressure and low error in population observations (Figures S33 to S35). For R and~~
~~R+M OMs with those conditions, all EM process error assumptions provided median errors~~
~~close to zero except for OMs with the true covariate effect strongest and the true covariate~~
~~time series was autocorrelated and had high temporal variation. For R+S OMs with those~~
~~conditions, only EMs with the correct process error assumption exhibited low (absolute)~~
~~median error.~~

Confidence interval bias

~~For logistic regressions of indicators of confidence intervals including β_M , factors that~~
~~provided the largest reductions in deviance were the OM fishing history and level of error in~~
~~population observations for all OM process error types (0.3-6%) (Table 7). Including second~~
~~and third order interactions provided the largest relative increase in deviance reduction~~
~~for all EMs fit to several R OMs with low covariate uncertainty and covariate variability~~
~~(Figure ??, first two columns) and for R+S EMs in many (8-11%) and R+S OMs (Figure~~
~~??, middle two columns) M OMs (5-8%) and lesser increases for R OMs (3-5%).~~

~~The EMs with either R or R~~The factors defining the primary branches of classification
~~trees for indicators of confidence interval inclusion of β_M matched those providing the~~
~~largest reductions in deviance for each of the OM process error types (Figure 7). Across~~
~~all simulations, rates of 95% confidence interval coverage indicated negative bias with rates~~
~~ranging 83-87% for the different OM process error types. Branches based on contrast in OM~~
~~fishing history and lower error in population observations indicated less bias in confidence~~
~~interval coverage except R+S process error configurations with median M assumed known~~
~~and covariate effects not estimated will have terminal year M correctly specified when fitted~~
~~to either R or R+S OMs with no covariate effect because M is constant in both the EM~~

~~and OM at the same value. We exclude those OM-EM combinations from our results here.~~
~~where the R EM configuration was incorrectly assumed.~~

~~In many of the OMs with temporal~~ The more detailed results for each OM and EM
configuration show that good confidence interval coverage can be obtained for many R
and R+S OM scenarios with low error in population observations and contrast in fishing
pressure~~and lower uncertainty in population observations, there was little difference in bias~~
~~for~~, but coverage was generally unreliable for all R+M OM scenarios (Figures S22 to S24).

Terminal year natural mortality rate

Estimation bias

Regressions of errors in estimation of terminal year M ~~among EMs whether the median~~
 ~~M parameter (show largest reductions in deviance from whether the EM treats β_M) was~~
~~estimated (Figure ??). However, when there were differences, bias in terminal year as known~~
~~(Table 8). For R and R+S OMs, annual M was constant, and therefore, terminal M~~
~~was closer to zero when the EM assumed known when the true β_M known, as would be~~
~~expected. For R OMs, all estimating models exhibited little evidence of bias except when~~
~~OMs simulated the largest covariate effect ($\beta_E = 0.5$). However, bias estimates were at worst~~
~~around 10% (Figure ??, left columns) was assumed known and there will be no error in any of~~
~~those EMs. Beyond the β_M assumption, the OM fishing history, level of error in population~~
~~observations, and EM assumptions for covariate effect and process error all provided notable~~
~~deviance reductions.~~

Regression trees for errors in estimation of terminal year M have primary branches based on
the whether β_M was assumed known or estimated for all OM process error types (Figure 8).
For R and R+S OMs, the branches where β_M was assumed known have no error as expected
because yearly M was equal to e^{β_M} , but there was also very little error in terminal year M
for the same branch in R+S EMs ~~generally exhibited the least evidence for bias and M OMs.~~

When β_M was estimated, median errors were near zero for many EMs in R OMs, but median error was problematic for this subset of EMs in R+M OMs. EMs generally estimated terminal $\hat{M}$ lower than the true value, particularly when observation uncertainty in the covariate was lower. For S OMs. On the other hand, detailed results for each OM and EM attribute show that there was little evidence of bias for R+S OMs when error in population observations was low and there was contrast in fishing pressure and the EM process error configuration was correct (Figure S45). In R+M OMs, many of the EM configurations provided little bias in terminal $\hat{M}$, but the biases were generally further from zero than the results for the R OMs (Figure ??, right columns). median errors indicated little bias overall with temporal contrast in fishing pressure, although detailed results demonstrated best reliability when there was also lower error in population observations (Figure S46).

Accuracy (lower RMSE)

Estimation accuracy

Like bias of terminal year $\hat{M}$ was similar for all of the EMs when fitted to R+M OMs (Figure ??). For R+S OMs, accuracy was generally better for $\hat{M}$, whether the EMs treated β_M as known or estimated provided the largest reduction in deviance (22-45%) for accuracy of estimates as measured by RMSE for all OM process error types (Table S4). Including second and third order interactions also provided large further reductions in deviance (21-38%). Also like bias, regression trees fit to log-RMSE of terminal M showed better accuracy when EMs assumed β_M was known rather than estimated (Figure S6). Better accuracy was also generally associated with lower error in population observations and temporal contrast in fishing pressure. For R+S EM than other process error assumptions, particularly when β_E was also estimated. Accuracy for OMs, R+S EMs was also generally as good as other EM provided better accuracy than incorrect process error assumptions even when the OM process error type was different. As would be expected, the

Using correlation to measure accuracy of terminal year $\widehat{M}$ where median M was assumed known was better than when it was estimated, when there were differences estimation provides a different perspective. Treatment of β_M provided relatively large reductions in deviance for all OM process error types (3-12%), but level of variability in the covariate was equally or more important (3-15%) as was whether the covariate effect was included in the EM (12%) for R and R+S OMs (Table S5). Including second and third order interactions also provided relatively large rather reductions in deviance (15 to 24%). Regression trees demonstrated better accuracy with higher covariate variability, lower error in population and covariate observations, and when EMs assumed β_M was known (Figure S7).

Terminal year spawning stock biomass

~~In OMs with temporal contrast in fishing pressure and lower uncertainty~~

Estimation bias

Fits of regression models to log-scale errors of terminal SSB resulted in largest reductions in deviance for OM fishing history (2-3%), error in population observations, ~~we found little bias in estimates of terminal year SSB for all EM configurations when fitted to R or (1% for R and R+M OMs (Figure ??). We also found little or no evidence of bias), and EM β_M assumption (2%; Table 9). The type of EM process error assumed also provided a deviance reduction similar in scale for R+S OMs when the EMs assumed the correct process error configuration. We found the worst bias in terminal year SSB estimation for R EMs with β_M estimated when fitted to R (1%). Including second and third order interactions of OM and EM attributes provided further deviance reductions of 5-10%.~~

The primary branches of regression trees fitted to log-scale errors in terminal SSB were based on type of OM fishing history for all OM process error types, but differences in median errors between branches were small (Figure 9). For R+S OMs, and R+M EMs also had more bias

than OMs, branches that assumed β_M was known indicated less bias than those where it was estimated. For R+S EMs in these OMs. Like the bias results, accuracy, OMs with constant fishing pressure and β_M estimated, less bias was indicated when the EM process error assumption was correct.

Estimation accuracy

Largest reductions in deviance for accuracy of terminal year SSB as measured by RMSE, was similar for many of EMs when fit to log-RMSE were provided by whether β_M was known or estimated (20-21%) and whether there was temporal contrast in fishing pressure (20-33%) across all OM process error types (Table S6). Lesser reductions were also provided by the EM process error assumption for R and R+M OMs. Accuracy was generally worse (higher RMSE) when EMs were fit to R+S OMs, and the worst accuracy occurred when the EM process error assumption did not match that of the OM (Figure ??). Inclusion of second and third order interactions of OM and EM attributes provided further reductions in deviance of 26-40%.

Primary branches of regression trees fit to log-RMSE of terminal year SSB were based on the same factors that provided the largest reduction in deviance (Figure S8). All branches with temporal change in fishing pressure provided better accuracy than branches with constant fishing pressure. Similarly, better accuracy occurred for branches with β_M assumed known rather than estimated, and lower error in population observations.

Using correlation to measure accuracy of terminal year SSB estimation, temporal contrast in fishing pressure and EM treatment of β_M provided relatively important reductions in deviance like RMSE, but level of error in population observations was important rather than EM process error type (Table S7). Including second and third order interactions also provided relatively large ruther reductions in deviance (6 to 20%). Regression trees demonstrated better accuracy with contrast in fishing pressure, lower error in population observations, and

when EMs assumed β_M was known (Figure S9).

Discussion

Our simulation study demonstrated that reliable inferences about effects on natural mortality require low observation error, contrast in fishing pressure, and large temporal variability in the environmental covariate. Under these conditions, reliable AIC-based selection and estimation of environmental effects on M is possible and reliable in certain scenarios can be possible even when the process error was mis-specified (e.g., R and R+S EMs fit to R+M OM). ~~In many of these same OMs, frequency of convergence of fitted models did not appear to suffer when covariate effects on M were estimated even when there was no effect simulated. However, these scenarios are information-rich in that there was contrast in population size (due to changes in fishing pressure) and the covariate affecting the population, and there was low uncertainty in population and covariate observations. Previous research has shown that estimation of a constant M parameter requires contrast in time series and informative data (Lee et al., 2011), so it is no surprise that estimation of these effects also requires relatively good information via more precise observations and higher contrast in the covariate time series.—~~

~~Even though estimation of covariate effects was unbiased in many scenarios, AIC could only reliably detect covariate effects~~ Under those same conditions, reliable estimation of standard errors and confidence interval coverage was also possible for R and R+M OMs ~~with contrast in covariates and low covariate uncertainty. In those scenarios where the covariate effect could be detected, CI coverage for the covariate effect was often biased even when S OMs especially when the EM process error type was correct. However, there was poor confidence interval coverage for R+M OMs even though~~ there was little ~~or no bias in the estimators bias in estimation~~ of the effect ~~and its standard error~~ standard errors, possibly related to correlation of the estimated effects and standard errors. Cadigan et al. (2024) found ~~CI~~

confidence interval coverage to be biased for SSB and F estimation in a state-space model in some scenarios ~~;~~ but they attributed the poor coverage to bias in Hessian-based standard error estimation ~~;~~ and their simulations held any process error random effects constant. ~~The coverage bias we observed, at least for some OM-EM combinations, may be related to correlation of the estimators of the covariate effect and the corresponding standard error and therefore consideration of~~ Consideration of other methods of calculating ~~CIs~~ confidence intervals may be warranted ~~(e.g., those based on profile likelihood and /or Monte Carlo sampling of the log-likelihood surface)~~ (e.g., Boyko and O'Meara, 2024). Previous research has shown that estimation of a constant M parameter requires contrast in time series and informative data (Lee et al., 2011), so it is unsurprising that estimation of covariate effects on M also requires relatively good information via more precise observations and higher contrast in the covariate time series.

We found AIC unreliable for determining the source of process error for EMs fit to R+M OMs where marginal temporal variability was specified as $\sigma_M = 0.3$. Miller et al. (In reviewa) investigated R+M OMs with two levels of M ~~process error~~ process-error variability ($\sigma_M \in \{0.1, 0.5\}$) and only found AIC able to accurately distinguish R+M process errors with the higher level of ~~process error variability ($\sigma_M = 0.5$).~~ We assumed $\sigma_M = 0.3$, intermediate to the values investigated by Miller et al. (In reviewa), and found process error inferences unreliable for the source of process error, indicating that the process-error variability. If AIC accuracy increases with σ_M , our results together would suggest the minimum level of variability required for detecting M process errors must be greater than $\sigma_M = 0.3$, but may still be less than for reliable detection may be with σ_M somewhere between 0.3 and 0.5. In our results and those by Miller et al. (In reviewa), inaccurate selection of AIC typically chose R EMs which would indicate the fitted R+M EMs would estimate no variability in the M process errors. Future studies like ours where R+M OMs are simulated with greater variability in M process errors would better inform reliability of covariate effect inferences under such scenarios.

Deriso et al. (2008) attempted to estimate process errors as well as covariate effects with M for Pacific herring ~~;~~ but similarly found no variability in M , suggesting there was too much uncertainty in the available observations relative to the true temporal variability in M . Given that we found covariate effect inferences using R EMs ~~was~~ were reliable in R+M OMs with apparently little variability in M process errors, the findings of Deriso et al. (2008) on covariate effects for Pacific herring would presumably also be robust to true low variability in M . However, they did not account for uncertainty in covariate observations, some of which would probably have substantial uncertainty (e.g., competition and predation covariates). We found higher covariate observation uncertainty ~~to cause~~ resulted in true covariate effects to be less detectable using AIC ~~;~~ but we did not investigate the implications for incorrectly assuming no covariate uncertainty for covariate inferences.

~~Any bias or poor accuracy for annual SSB estimation was primarily a function of whether or not the median~~ It was important to consider RMSE and correlation of estimates with the true values for reliable estimation of terminal year M parameter was estimated or known and the types of because summaries of relative errors could suggest unbiasedness that is not relevant due to the distribution of true values being centered at an assumed constant value or near an estimated constant value in the EM (e.g., R and R+S EMs that have a temporally constant M fit to OMs with temporally varying M). Despite the inability of AIC to distinguish M process errors, ~~rather than the treatment of the covariate effects~~ reliable estimation of annual M was possible in some scenarios regardless of the EM process error assumption. To achieve little bias and high correlation of terminal M estimates required larger variability in the covariate (when OMs and EMs included the effect on M . For example, we found estimation of SSB was better when the EM had the process error correctly specified.) along with lower population and covariate observation error. It was also more difficult when EMs estimated β_M rather than treating it as known. However, there was little correlation of terminal year M estimates and true values for R+M OMs without covariate effects for any combination of OM and EM attributes (Figure S52). Therefore, our results suggest estimating annual

M is easier when the temporal variation is due to effects of highly variable covariates than unexplained variation in M . Like the results for AIC-based selection, this may be a function of the level of variability in M we assumed for R+M OMs.

Accurate estimation of terminal year SSB, was more robust than terminal year M . When OMs had contrast in fishing pressure and low error in population observations, there was little bias and high correlation of terminal year SSB estimates regardless of whether OMs and EMs included covariate effects. However, for R+S OMs, it was important for the EM process error configuration to be correct. Fortunately, our results and those by Miller et al. (In reviewa) ~~demonstrate that marginal AIC seems to be a good tool for determining whether this source~~, suggest AIC-based model selection is accurate for this type of process errors ~~should be included in the model~~. However, the reliability of the ~~estimation of SSB~~ SSB estimation does break down in the less ideal scenarios when there is higher ~~population observation error~~ error in population observations, and lack of contrast in fishing pressure (e.g., Figures S53 to S55).

The R+S and R+M EMs both include process error for the apparent survival of cohorts and would be expected to perform similarly, and they did in our simulations when the OM and EM matched. However, we found that the ~~biases~~ accuracy for model output like SSB and F that are important for management was generally worse using R+M EMs for R+S OMs ~~were generally worse~~ than using R+S EMs for R+M OMs in the high information OMs with contrast in fishing pressure and low error in population observations. Additionally, there are implications for biological reference points for R+M EMs (e.g., Legault and Palmer, 2016) that are not present with R+S EMs. ~~So, we~~ We therefore recommend following the suggestion from Li et al. (2024) to prefer R+S EMs over R+M ~~OMs~~ EMs unless strong biological evidence is present to support a particular R+M ~~OM~~ EM. Such support could be found through both biological understanding (e.g., Trijoulet et al., 2020) as well as statistical properties such as a large delta AIC for R+M compared to R+S associated with greater temporal variability in ~~natural mortality~~ M (Miller et al., In reviewa).

The pattern of improved inferences using state-space models in the presence of higher contrast in fishing pressure, lower observation error and greater process error variability that emerges from our work here and by Miller et al. (In reviewb) is consistent with expectations from statistical theory. State-space models are more reliable when observation error is lower than process error (Auger-Méthé et al., 2016), and, more generally, maximum likelihood estimation is less biased and more accurate as the information in the data increases (Neyman and Scott, 1948; Kiefer and Wolfowitz, 1956). However, the greater complexity of state-space assessment models over the more well-studied linear and Guassian state-space models makes determining sufficient levels of precision in observations difficult. Statistical information available to the state-space assessment model could be increased with longer time series of observations, more time series (e.g. multiple indices of abundance and corresponding age composition data sets), or more precise observations of a given time series of observations (e.g., greater index observation precision or more sampling for age composition). Therefore, it will be important to conduct self-test simulations (where the OM and EM are the same) on a case-by-case basis to ensure the reliability of models used to manage commercially important fish stocks.

The ability to accurately infer covariate effects on M in some realistic—situations indicates that such investigations may be fruitful. ~~Ability—~~The ability to make inferences could improve further when WHAM is extended to incorporate tagging data (~~Miller et al., In reviewb~~)(Miller et al., 2025). Tagging data can greatly inform ~~natural—mortality—estimation—(Pollock et al., 1991; Hampton, 2000),—~~ M estimation (Pollock et al., 1991; Hampton, 2000) and this impact on M estimation should also apply to estimation of covariate effects or unexplained temporal variation in M . Given our findings and planned future WHAM development, we expect investigations of and accounting for covariate effects on M to become more common within the fisheries stock assessment process. At the same time, it will be equally important to conduct research that will improve our understanding of how best to measure the depletion of stocks and determine

902 catch advice for these stocks with covariate effects on M .

903 Acknowledgements

904 This work was funded by NOAA Fisheries Northeast Fisheries Science Center. We thank the
905 associate editor, two anonymous reviewers, and Emily Liljestrand for constructive comments
906 on previous drafts that led to a much improved manuscript.

References

- Aitchison, J. and Shen, S.M. 1980. Logistic-normal distributions: some properties and uses. *Biometrika* **67**(2): 261–272. doi:10.2307/2335470.
- Auger-Méthé, M., Field, C., Albertsen, C.M., Derocher, A.E., Lewis, M.A., Jonsen, I.D., and Mills Flemming, J. 2016. State-space models’ dirty little secrets: even simple linear Gaussian models can have estimation problems. *Scientific reports* **6**(1): 26677. doi:10.1038/srep26677.
- Boyko, J.D. and O’Meara, B.C. 2024. dentist: Quantifying uncertainty by sampling points around maximum likelihood estimates. *Methods in Ecology and Evolution* **15**(4): 628–638. doi:10.1111/2041-210X.14297.
- Breiman, L., Friedman, J.H., Olshen, R.A., and Stone, C.J. 1984. *Classification and Regression Trees*. Chapman and Hall/CRC, New York, NY USA. doi:10.1201/9781315139470. 368p.
- Britten, G., Brooks, E.N., and Miller, T.J. 2026. Factors affecting the inference on stock-recruitment relationships in state-space assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* doi:10.1139/cjfas-2025-0210.
- Britten, G., Brooks, E.N., and Miller, T.J. In review. Factors affecting the inference on stock-recruitment relationships in state-space assessment models. For *Canadian Journal of Fisheries and Aquatic Sciences*, State-space assessment models special collection.
- Brodziak, J., Cadrin, S.X., Legault, C.M., and Murawski, S.A. 2008. Goals and strategies for rebuilding New England groundfish stocks. *Fisheries Research* **94**(3): 355–366. doi:10.1016/j.fishres.2008.03.008.
- Cadigan, N.G. 2016. A state-space stock assessment model for northern cod, including under-

- reported catches and variable natural mortality rates. *Canadian Journal of Fisheries and Aquatic Sciences* **73**(2): 296–308. doi:10.1139/cjfas-2015-0047.
- Cadigan, N.G., Albertsen, C.M., Zheng, N., and Nielsen, A. 2024. Are state-space stock assessment model confidence intervals accurate? case studies with sam and barents sea stocks. *Fisheries Research* **272**: 106950. doi:10.1016/j.fishres.2024.106950.
- Carvalho, F., Winker, H., Courtney, D., Kapur, M., Kell, L., Cardinale, M., Schirripa, M., Kitakado, T., Yemane, D., Piner, K.R., Maunder, M.N., Taylor, I., Wetzel, C.R., Doering, K., Johnson, K.F., and Methot, R.D. 2021. A cookbook for using model diagnostics in integrated stock assessments. *Fisheries Research* **240**: 105959. doi:10.1016/j.fishres.2021.105959.
- Clopper, C.J. and Pearson, E.S. 1934. The use of confidence or fiducial limits illustrated in the case of the binomial. *Biometrika* **26**(4): 404–413. doi:10.1093/biomet/26.4.404.
- Collier, Z.K., Zhang, H., and Soyoye, O. 2022. Alternative methods for interpreting Monte Carlo experiments. *Communications in Statistics - Simulation and Computation* pp. 1–16. doi:10.1080/03610918.2022.2082474.
- Deriso, R.B., Maunder, M.N., and Pearson, W.H. 2008. Incorporating covariates into fisheries stock assessment models with application to pacific herring. *Ecological Applications* **18**(5): 1270–1286. doi:10.1890/07-0708.1.
- Gonzalez, O., O'Rourke, H.P., Wurpts, I.C., and Grimm, K.J. 2018. Analyzing Monte Carlo simulation studies with classification and regression trees. *Structural Equation Modeling: A Multidisciplinary Journal* **25**(3): 403–413. doi:10.1080/10705511.2017.1369353.
- Haltuch, M.A. and Punt, A.E. 2011. The promises and pitfalls of including decadal-scale climate forcing of recruitment in groundfish stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* **68**(5): 912–926. doi:10.1139/f2011-030.

- Hampton, J. 2000. Natural mortality rates in tropical tunas: size really does matter. *Canadian Journal of Fisheries and Aquatic Sciences* **57**(5): 1002–1010. doi:10.1139/f99-287.
- Harwell, M., Kohli, N., and Peralta-Torres, Y. 2018. A survey of reporting practices of computer simulation studies in statistical research. *The American Statistician* **72**(4): 321–327. doi:10.1080/00031305.2017.1342692.
- Hoyle, S.D., Maunder, M.N., Punt, A.E., Mace, P.M., Devine, J.A., and A’mar, Z.T. 2022. Preface: Developing the next generation of stock assessment software. *Fisheries Research* **246**: 106176.
- Johnson, K.F., Councill, E., Thorson, J.T., Brooks, E., Methot, R.D., and Punt, A.E. 2016. Can autocorrelated recruitment be estimated using integrated assessment models and how does it affect population forecasts? *Fisheries Research* **183**: 222–232. doi:10.1016/j.fishres.2016.06.004.
- Kiefer, J. and Wolfowitz, J. 1956. Consistency of the maximum likelihood estimator in the presence of infinitely many incidental parameters. *The Annals of Mathematical Statistics* **27**(4): 887–906.
- Kristensen, K., Nielsen, A., Berg, C.W., Skaug, H., and Bell, B.M. 2016. TMB: Automatic differentiation and Laplace approximation. *Journal of Statistical Software* **70**(5): 1–21.
- Lee, H.H., Maunder, M.N., Piner, K.R., and Methot, R.D. 2011. Estimating natural mortality within a fisheries stock assessment model: An evaluation using simulation analysis based on twelve stock assessments. *Fisheries Research* **109**(1): 89–94. doi:10.1016/j.fishres.2011.01.021.
- Legault, C.M. and Palmer, M.C. 2016. In what direction should the fishing mortality target change when natural mortality increases within an assessment? *Canadian Journal of Fisheries and Aquatic Sciences* **73**(3): 349–357. doi:10.1139/cjfas-2015-0232.

- Legault, C.M., Wiedenmann, J., Deroba, J.J., Fay, G., Miller, T.J., Brooks, E.N., Bell, R.J., Langan, J.A., Cournane, J.M., Jones, A.W., and Muffley, B. 2023. Data-rich but model-resistant: an evaluation of data-limited methods to manage fisheries with failed age-based stock assessments. *Canadian Journal of Fisheries and Aquatic Sciences* **80**(1): 27–42. doi:10.1139/cjfas-2022-0045.
- Li, C., Deroba, J.J., Miller, T.J., Legault, C.M., and Perretti, C.T. 2024. An evaluation of common stock assessment diagnostic tools for choosing among state-space models with multiple random effects processes. *Fisheries Research* **273**: 106968. doi:10.1016/j.fishres.2024.106968.
- Li, C., Deroba, J.J., Berger, A.M., Goethel, D.R., Langseth, B.J., Schueller, A.M., and Miller, T.J. 2026a. Random effects on numbers-at-age transitions implicitly account for movement dynamics and improve performance within a state-space stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* **83**: 1–18. doi:10.1139/cjfas-2025-0092.
- Li, C., Deroba, J.J., Miller, T.J., Legault, C.M., and Perretti, C. 2026b. Guidance on bias-correction of log-normal random effects and observations in state-space assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* **83**: 1–11. doi:10.1139/cjfas-2025-0093.
- Li, C., Deroba, J.J., Berger, A.M., Goethel, D.R., Langseth, B.J., Schueller, A.M., and Miller, T.J. In reviewa. Random effects on numbers-at-age transitions implicitly account for movement dynamics and improve performance within a state-space stock assessment. For *Canadian Journal of Fisheries and Aquatic Sciences*, State-space assessment models special collection.
- Li, C., Deroba, J.J., Miller, T.J., Legault, C.M., and Perretti, C. In reviewb. Guidance on bias-correction of log-normal random effects and observations in state-space assessment models. For *Canadian Journal of Fisheries and Aquatic Sciences*, State-space assessment models special collection.

1003 MacKinnon, D.P., Warsi, G., and Dwyer, J.H. 1995. A simulation study of medi-
 1004 ated effect measures. *Multivariate Behavioral Research* **30**(1): 41–62. doi:10.1207/
 1005 s15327906mbr3001_3.

1006 Magnusson, A. and Hilborn, R. 2007. What makes fisheries data informative? *Fish and*
 1007 *Fisheries* **8**(4): 337–358. doi:https://doi.org/10.1111/j.1467-2979.2007.00258.x.

1008 Miller, T.J. and Brooks, E.N. 2021. Steepness is a slippery slope. *Fish and Fisheries* **22**(3):
 1009 634–645. doi:10.1111/faf.12534.

1010 Miller, T.J. and Hyun, S.Y. 2018. Evaluating evidence for alternative natural mortality and
 1011 process error assumptions using a state-space, age-structured assessment model. *Canadian*
 1012 *Journal of Fisheries and Aquatic Sciences* **75**(5): 691–703. doi:10.1139/cjfas-2017-0035.

1013 Miller, T.J. and Stock, B.C. 2020. The Woods Hole Assessment Model (WHAM).

1014 Miller, T.J., Hare, J.A., and Alade, L. 2016. A state-space approach to incorporating environ-
 1015 mental effects on recruitment in an age-structured assessment model with an application
 1016 to Southern New England yellowtail flounder. *Canadian Journal of Fisheries and Aquatic*
 1017 *Sciences* **73**(8): 1261–1270. doi:10.1139/cjfas-2015-0339.

1018 Miller, T.J., Curti, K.L., and Hansell, A.C. 2025. Space for WHAM: a multi-region, multi-
 1019 stock generalization of the Woods Hole Assessment Model with an application to black sea
 1020 bass. *Canadian Journal of Fisheries and Aquatic Sciences* **82**. doi:10.1139/cjfas-2025-0097.

1021 Miller, T.J., Britten, G., Brooks, E.N., Fay, G., Hansell, A., Legault, C.M., Li, C., Muffley,
 1022 B., Stock, B.C., and Wiedenmann, J. In reviewa. An investigation of factors affecting
 1023 inferences from and reliability of state-space age-structured assessment models. For *Cana-*
 1024 *dian Journal of Fisheries and Aquatic Sciences*, State-space assessment models special
 1025 collection.

1026 Miller, T.J., Curti, K.L., and Hansell, A.C. In reviewb. Space for WHAM: a multi-region,
 1027 multi-stock generalization of the Woods Hole Assessment Model with an application to
 1028 black sea bass. For Canadian Journal of Fisheries and Aquatic Sciences, State-space
 1029 assessment models special collection.

1030 Myers, R.A. 1998. When do environment–recruitment correlations work? Reviews in Fish
 1031 Biology and Fisheries **8**(3): 229–249. doi:10.1023/A:1008828730759.

1032 NEFSC 2022a. Final report of the haddock research track as-
 1033 sessment working group. Available at [https://s3.us-east-](https://s3.us-east-1.amazonaws.com/nefmc.org/14b_EGB_Research_Track_Haddock_WG_Report_DRAFT.pdf)
 1034 [1.amazonaws.com/nefmc.org/14b_EGB_Research_Track_Haddock_WG_Report_DRAFT.pdf](https://s3.us-east-1.amazonaws.com/nefmc.org/14b_EGB_Research_Track_Haddock_WG_Report_DRAFT.pdf).

1035 NEFSC 2022b. Report of the american plaice research track working group. Available at
 1036 https://s3.us-east-1.amazonaws.com/nefmc.org/2_American-Plaice-WG-Report.pdf.

1037 NEFSC 2024. Butterfish research track assessment report. US Dept Commer Northeast Fish
 1038 Sci Cent Ref Doc. 24-03; 191 p.

1039 Neyman, J. and Scott, E.L. 1948. Consistent estimates based on partially consistent obser-
 1040 vations. *Econometrica* **16**(1): 1–32. doi:10.2307/1914288.

1041 Nielsen, A. and Berg, C.W. 2014. Estimation of time-varying selectivity in stock assessments
 1042 using state-space models. *Fisheries Research* **158**: 96–101. doi:10.1016/j.fishres.2014.01.
 1043 014.

1044 Pedersen, M.W. and Berg, C.W. 2017. A stochastic surplus production model in continuous
 1045 time. *Fish and Fisheries* **18**(2): 226–243. doi:10.1111/faf.12174.

1046 Pollock, K., Hoenig, J., and Jones, C. 1991. Estimation of fishing and natural mortality when
 1047 a tagging study is combined with a creel survey or port sampling. *American Fisheries*
 1048 *Society Symposium* **12**: 423–434.

1049 Punt, A.E. 2023. Those who fail to learn from history are condemned to repeat it: A per-
 1050 spective on current stock assessment good practices and the consequences of not following
 1051 them. *Fisheries Research* **261**: 106642.

1052 Stock, B.C. and Miller, T.J. 2021. The Woods Hole Assessment Model (WHAM): A general
 1053 state-space assessment framework that incorporates time- and age-varying processes via
 1054 random effects and links to environmental covariates. *Fisheries Research* **240**: 105967.
 1055 doi:10.1016/j.fishres.2021.105967.

1056 Stock, B.C., Xu, H., Miller, T.J., Thorson, J.T., and Nye, J.A. 2021. Implementing two-
 1057 dimensional autocorrelation in either survival or natural mortality improves a state-space
 1058 assessment model for Southern New England-Mid Atlantic yellowtail flounder. *Fisheries*
 1059 *Research* **237**: 105873. doi:10.1016/j.fishres.2021.105873.

1060 Szuwalski, C.S., Ianelli, J.N., and Punt, A.E. 2018. Reducing retrospective patterns in stock
 1061 assessment and impacts on management performance. *ICES Journal of Marine Science*
 1062 **75**(2): 596–609. doi:10.1093/icesjms/fsx159.

1063 Therneau, T. and Atkinson, B. 2025. *rpart*: Recursive Partitioning and Regression Trees.
 1064 R package version 4.1.25.

1065 Thorson, J.T. and Minto, C. 2015. Mixed effects: a unifying framework for statistical
 1066 modelling in fisheries biology. *ICES Journal of Marine Science* **72**(5): 1245–1256. doi:
 1067 10.1093/icesjms/fsu213.

1068 Thulin, M. 2014. The cost of using exact confidence intervals for a binomial proportion.
 1069 *Electronic Journal of Statistics* **8**(1): 817–840. doi:10.1214/14-EJS909.

1070 Trijoulet, V., Fay, G., and Miller, T.J. 2020. Performance of a state-space multispecies model:
 1071 What are the consequences of ignoring predation and process errors in stock assessments?
 1072 *Journal of Applied Ecology* **57**(1): 121–135. doi:10.1111/1365-2664.13515.

1073 Trijoulet, V., Albertsen, C.M., Kristensen, K., Legault, C.M., Miller, T.J., and Nielsen, A.
1074 2023. Model validation for compositional data in stock assessment models: Calculating
1075 residuals with correct properties. *Fisheries Research* **257**: 106487. doi:10.1016/j.fishres.
1076 2022.106487.

1077 Vaida, F. and Blanchard, S. 2005. Conditional Akaike information for mixed-effects models.
1078 *Biometrika* **92**(2): 351–370. doi:10.1093/biomet/92.2.351.

1079 Wang, S., Cadigan, N.G., and Benoît, H.P. 2017. Inference about regression parameters
1080 using highly stratified survey count data with over-dispersion and repeated measurements.
1081 *Journal of Applied Statistics* **44**(6): 1013–1030. doi:10.1080/02664763.2016.1191622.

1082 Wiedenmann, J., Free, C.M., and Jensen, O.P. 2019. Evaluating the performance of data-
1083 limited methods for setting catch targets through application to data-rich stocks: A case
1084 study using northeast U.S. fish stocks. *Fisheries Research* **209**(1): 129–142. doi:10.1016/
1085 j.fishres.2018.09.018.

1086 Yee, T.W. 2008. The VGAM package. *R News* **8**(2): 28–39.

1087 Yee, T.W. 2015. *Vector Generalized Linear and Additive Models: With an Implementation*
1088 in R. Springer, New York, NY USA. doi:10.1007/978-1-4939-2818-7. 589p.




Fig. 1. ~~Estimated probability of fits~~ Classification trees indicating primary factors determining convergence as defined by providing Hessian-based standard errors for EMs operating models (OMs) with ~~alternative~~ process error assumptions, ~~treatment of median natural mortality on recruitment only~~ (β_M known or estimated), ~~recruitment and treatment of covariate effect apparent survival~~ ($\beta_E = 0$ or estimated). The OMs have R(left) and R+S(middle), or recruitment and natural mortality (R+M(right)) process error structures, alternative configurations of covariate time series structure and levels of observation uncertainty. Nodes denote percent convergence (rowstop) - and three levels number of true covariate effect on median natural




Fig. 2. ~~For each OM~~ Classification trees indicating primary factors determining which process error assumption in estimating models (EMs) provides the lowest AIC for operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M). Each node shows the ~~proportion percentage of simulated data sets where the EM type (treatment of environmental covariate effect and assumed fits with a given process error type) had assumption having the lowest AIC and the number of observations for the corresponding subset of fits.~~ All OMs had low observation. Lower or higher accuracy of the process error assumption are indicated by more red or green polygons, respectively. See Table S1 for ~~fish population observations description of OM and temporal contrast in fishing pressure~~ EM factors defining nodes and branches. ~~All EMs estimated median natural mortality rate.~~




Fig. 3. Median-error Classification trees indicating primary factors determining whether the correct EM assumption for covariate effect on M (MEno effect with OM $\beta_E = 0$ or estimated effect when OM $\beta_E > 0$) provides the lowest AIC for OMs assuming values for β_E of estimates 0, 0.25, and 0.5. Nodes denote the percentage of environmental effect on natural mortality β_E from fitting estimating models (EMs) with alternative process error assumptions correct assumption and treatment lowest AIC (top), and number of median natural mortality observations (β_M known or estimated bottom) for the corresponding subset. All OMs had low observation Lower or higher accuracy of the process error assumption are indicated by more red or green polygons, respectively. See Table S1 for description of OM and contrast in fishing mortality EM factors defining nodes and branches. Vertical lines represent 95% confidence intervals.

Fig. 4. **Probability** Regression trees indicating primary factors determining reductions in sums of 95%-confidence interval for β_E -containing squares of errors in estimation of the true value covariate effect on M for EMs operating models (OMs) with alternative process error assumptions on recruitment only (R), recruitment and treatment of median apparent survival (R+S), or recruitment and natural mortality (β_M known or estimated R+M). All OMs had low observation Each node shows the median error (top) and contrast in fishing mortality number of observations (bottom) for the corresponding subset. Vertical lines represent 95%-confidence intervals Lower or higher median absolute errors of the process error assumption are indicated by more green or red polygons, respectively. See Table S1 for description of OM and EM factors defining nodes and branches.




Fig. 5. Median error Classification trees indicating primary factors determining whether confidence intervals for estimated covariate effects on M in estimating models (ME) of estimates of β_M from fitting EMs) fitted to operating models (OMs) with alternative process error assumptions on recruitment only (R), recruitment and treatment of covariate effect apparent survival ($\beta_E = 0$ R+S), or estimated recruitment and natural mortality (R+M) included the true value. All OMs had low observation Error. Each node shows the median error (top) and contrast in fishing mortality number of observations (bottom) for the corresponding subset. Vertical lines represent 95% confidence intervals. Lower or higher median absolute errors of the process error assumption are indicated by more green or red polygons, respectively. See Table S1 for description of OM and EM factors defining nodes and branches.




Fig. 6. Probability-Regression trees indicating primary factors determining reductions in sums of 95% confidence interval for β_M containing squares of errors in estimation of the true value for median M parameter (β_M) in estimating models (EMs) fitted to operating models (OMs) with alternative-process error assumptions on recruitment only (R), recruitment and treatment of covariate effect apparent survival ($\beta_E = 0$ R+S), or estimated recruitment and natural mortality (R+M). All OMs had low observation Each node shows the median error (top) and contrast in fishing mortality number of observations (bottom) for the corresponding subset. Vertical lines represent 95% confidence intervals Lower or higher median absolute errors of the process error assumption are indicated by more green or red polygons, respectively. See Table S1 for description of OM and EM factors defining nodes and branches.




Fig. 7. ~~Median relative error~~ Classification trees indicating primary factors determining whether confidence intervals for ~~estimated median M parameter (MRE_{β_M}) of estimates of natural mortality rate in the terminal year for estimating models (EMs) fitted to operating models (OMs) with alternative process error assumptionson recruitment only (R), treatment of covariate effect recruitment and apparent survival ($\beta_E = 0$ or estimated R+S), or recruitment and treatment of median natural mortality parameter (β_M estimated or known R+M) included the true value. All OMs had low observation~~ Each node shows the median error (top) and ~~contrast in fishing mortality~~ number of observations (bottom) for the corresponding subset. Lower or higher median absolute errors of the process error assumption are indicated by more green or red polygons, respectively. See Table S1 for description of OM and EM factors defining nodes and branches.




Fig. 8. ~~Median relative error (MRE)~~ Regression trees indicating primary factors determining reductions in sums of ~~estimates~~ squares of spawning stock biomass (SSB) errors in estimation measured by Eq. 2 for the terminal year ~~for~~ M in estimating models (EMs) fitted to operating models (OMs) with ~~alternative~~ process error assumptions on recruitment only (R), ~~treatment of~~ covariate effect recruitment and apparent survival ($\beta_E = 0$ or estimated R+S), or recruitment and ~~treatment of~~ median natural mortality parameter (β_M estimated or known R+M). All OMs had low observation ~~Each node shows the median error (top) and contrast in fishing mortality~~ number of observations (bottom) for the corresponding subset. Lower or higher median absolute errors of the process error assumption are indicated by more green or red polygons, respectively. See Table S1 for description of OM and EM factors defining nodes and branches.




Fig. 9. Regression trees indicating primary factors determining reductions in sums of squares of errors in estimation measured by Eq. 2 for the terminal year SSB in estimating models (EMs) fitted to operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M). Each node shows the median error (top) and number of observations (bottom) for the corresponding subset. Lower or higher median absolute errors of the process error assumption are indicated by more green or red polygons, respectively. See Table S1 for description of OM and EM factors defining nodes and branches.

Table 1. Percent reduction in deviance for logistic regression models fit to indicators of convergence (providing Hessian-based standard errors). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	0.86	<0.01	0.24
OM Obs. Error	0.56	0.03	0.26
OM σ_e	0.66	2.99	0.93
OM σ_E	0.53	2.22	0.62
OM ρ_E	<0.01	0.02	<0.01
OM β_E	0.02	<0.01	<0.01
EM Process Error	24.53	9.33	23.06
EM β_E assumption	0.16	2.96	0.51
EM M Assumption	0.18	2.83	0.40
All factors	28.92	22.67	27.34
+ All Two Way	36.15	33.54	34.03
+ All Three Way	37.95	36.58	35.57

Table 2. Percent reduction in deviance for multinomial logistic regression models fit to indicators of process error assumption in the estimating models (EMs) with lowest AIC. Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	0.23	0.06	0.30
OM Obs. Error	0.54	18.58	13.43
OM σ_e	8.41	0.05	2.23
OM σ_E	0.17	0.08	0.59
OM ρ_E	1.29	0.03	0.06
OM β_E	3.42	0.04	0.38
EM M Assumption	0.35	0.34	0.48
EM β_E Assumption Correct	2.35	0.22	0.78
All factors	21.97	19.29	19.07
+ All Two Way	38.25	21.13	24.73
+ All Three Way	46.26	22.85	26.97

Table 3. Percent reduction in deviance for logistic regression models fit to indicators of correct estimating model (EM) covariate effect assumption (no effect with OM $\beta_E = 0$ or estimated effect when OM $\beta_E > 0$) with lowest AIC. Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	$\beta_E = 0$	$\beta_E = 0.25$	$\beta_E = 0.5$
OM F history	<0.01	0.09	0.18
OM Obs. Error	1.19	4.06	4.64
OM Process Error	4.11	0.71	0.25
OM σ_e	0.66	2.00	1.97
OM σ_E	0.35	7.97	20.66
OM ρ_E	0.21	1.23	1.48
EM M Assumption	0.23	0.15	0.15
EM Process Error Assumption Correct	1.95	0.39	0.06
All factors	6.95	17.65	33.44
+ All Two Way	10.58	23.21	38.36
+ All Three Way	11.83	24.95	39.79

Table 4. Percent reduction in deviance for linear regression models fit to errors in estimation of the covariate effect on natural mortality rate ($\hat{\beta}_E$). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	0.03	0.06	0.02
OM Obs. Error	<0.01	0.06	0.02
OM σ_e	0.04	0.08	0.06
OM σ_E	<0.01	0.02	0.01
OM ρ_E	<0.01	0.01	<0.01
OM β_E	0.05	<0.01	<0.01
EM Process Error	0.01	0.06	0.01
EM M assumption	0.02	0.02	<0.01
All factors	0.18	0.37	0.14
+ All Two Way	0.50	1.11	0.35
+ All Three Way	1.38	1.88	0.66

Table 5. Percent reduction in deviance for logistic regression models fit to indicators of whether confidence intervals for covariate effect estimates includes the true value. Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	0.24	0.05	0.01
OM Obs. Error	1.36	1.91	1.78
OM σ_e	8.26	0.24	0.92
OM σ_E	0.92	1.70	1.56
OM ρ_E	0.03	0.05	0.06
OM β_E	11.10	0.68	2.72
EM Process Error	0.01	6.51	0.27
EM M assumption	0.25	0.29	0.06
All factors	23.61	12.16	7.78
+ All Two Way	30.17	18.08	12.63
+ All Three Way	32.58	20.62	14.87

Table 6. Percent reduction in deviance for linear regression models fit to errors in estimation of the median natural mortality rate parameter ($\hat{\beta}_M$). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	4.03	16.52	6.51
OM Obs. Error	2.74	0.54	0.64
OM σ_e	0.01	<0.01	0.01
OM σ_E	0.18	0.02	0.15
OM ρ_E	0.07	0.05	<0.01
OM β_E	0.37	0.04	0.10
EM Process Error	2.31	11.99	2.65
EM β_E assumption	2.83	7.20	3.25
All factors	12.14	33.71	12.81
+ All Two Way	21.74	46.46	19.84
+ All Three Way	26.38	49.41	22.24

Table 7. Percent reduction in deviance for logistic regression models fit to indicators of whether confidence intervals for estimates of the median natural mortality rate parameter ($\hat{\beta}_M$) includes the true value. Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	6.08	1.46	0.14
OM Obs. Error	1.48	0.41	0.29
OM σ_e	0.01	0.06	0.01
OM σ_E	0.13	0.02	0.04
OM ρ_E	<0.01	<0.01	0.02
OM β_E	0.08	0.11	0.09
EM Process Error	0.53	0.08	0.04
EM β_E assumption	<0.01	0.19	0.01
All factors	8.46	2.32	0.62
+ All Two Way	11.97	10.57	5.99
+ All Three Way	13.35	13.51	8.44

Table 8. Percent reduction in deviance for linear regression models fit to errors in estimation measured by Eq. 2 for the terminal year natural mortality rate (M). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
Convergence	<0.01	0.18	0.01
OM F history	0.66	2.69	1.00
OM Obs. Error	0.29	0.43	0.16
OM σ_e	<0.01	<0.01	<0.01
OM σ_E	0.02	<0.01	<0.01
OM ρ_E	<0.01	0.01	<0.01
OM β_E	0.03	0.01	0.01
EM Process Error	0.22	1.42	0.43
EM β_E assumption	0.35	1.07	0.39
EM M assumption	0.84	5.10	1.22
All factors	2.56	10.83	3.37
+ All Two Way	5.41	18.75	6.38
+ All Three Way	7.93	22.49	8.76

Table 9. Percent reduction in deviance for linear regression models fit to errors in estimation measured by Eq. 2 for the terminal year spawning stock biomass (SSB). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
Convergence	0.02	0.13	<0.01
OM F history	2.08	2.74	1.75
OM Obs. Error	1.09	0.18	1.19
OM σ_e	0.01	<0.01	<0.01
OM σ_E	<0.01	<0.01	<0.01
OM ρ_E	<0.01	<0.01	<0.01
OM β_E	<0.01	0.01	<0.01
EM Process Error	0.52	1.05	0.57
EM β_E assumption	<0.01	0.05	0.01
EM M assumption	1.76	2.12	1.51
All factors	5.97	6.58	5.21
+ All Two Way	12.81	13.00	11.97
+ All Three Way	16.41	15.57	15.73

1089 Supplemental Materials

1090 Referenced Figures and Tables



Fig. S1. Example simulations of environmental covariate latent processes and observations with different levels of observation error, and different assumptions about variability of the latent process.




Fig. S2. The proportion mature at age, weight at age, fleet and index selectivity at age, and Beverton-Holt stock-recruit relationship assumed for the population in all **operating models**OMs. For **operating models**-OMs with random effects on fleet selectivity, this represents the selectivity at the mean of the random effects.

Example simulations of environmental covariate latent processes and observations with different levels of observation error, and different assumptions about variability of the latent process.

69




Fig. S3 Example simulations of annual M that may be a function of a temporally varying environmental covariate and

Example simulations of annual natural mortality rates that may be a function of a temporally varying environmental covariate and autoregressive random effects.

70




Fig. S4. Regression trees indicating primary factors determining reductions in sums of squares of errors in estimation of the Hessian-based standard error estimates for covariate effect on M ($SE(\hat{\beta}_E)$) for operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M). Each node shows the median error (top) and number of observations (bottom) for the corresponding subset. Lower or higher median absolute errors of the process error assumption are indicated by more green or red polygons, respectively.




Fig. S5. Regression trees indicating primary factors determining reductions in sums of squares of errors in estimation of the Hessian-based standard error estimates for median M parameter ($\widehat{SE}(\hat{\beta}_M)$) in EMs fitted to operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M). Each node shows the median error (top) and number of observations (bottom) for the corresponding subset. Lower or higher median absolute errors of the process error assumption are indicated by more green or red polygons, respectively.




Fig. S6. Regression trees indicating primary factors determining reductions in sums of squares for the log-transformed RMSE of terminal year M in EMs fitted to operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M). Each node shows the median error (top) and number of observations (bottom) for the corresponding subset. Lower or higher median RMSE are indicated by more green or red polygons, respectively.




Fig. S7. Regression trees indicating primary factors determining reductions in sums of squares for the correlation of terminal year M estimates and true values in EMs fitted to operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M). Each node shows the median error (top) and number of observations (bottom) for the corresponding subset. Lower or higher median correlations are indicated by more red or green polygons, respectively.




Fig. S8. Regression trees indicating primary factors determining reductions in sums of squares for the log-transformed RMSE of terminal year SSB in EMs fitted to operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M). Each node shows the median error (top) and number of observations (bottom) for the corresponding subset. Lower or higher median RMSE are indicated by more red or green polygons, respectively.




Fig. S9. Regression trees indicating primary factors determining reductions in sums of squares for the correlation of terminal year SSB estimates and true values in EMs fitted to operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M). Each node shows the median error (top) and number of observations (bottom) for the corresponding subset. Lower or higher median correlations are indicated by more red or green polygons, respectively.

Table S1. Operating (OM) and estimating model (EM) attributes used in analyses of deviance reduction and classification and regression trees. R, R+S, and R+M, denote models (OM or EM) with process error on recruitment only, recruitment and apparent survival, and recruitment and natural mortality, respectively. SD denotes standard deviation and F_{MSY} is the fishing mortality rate that achieves maximum sustainable yield.

Factor	Levels
OM Process error	R, R+S, R+M
OM Fishing History	$F_{\text{MSY}}, 2.5F_{\text{MSY}} \rightarrow F_{\text{MSY}}$
OM Covariate effect size (β_E)	0, 0.25, 0.5
OM Population Observation Error	Low, High
OM Covariate Observation Error	Low ($\sigma_e = 0.1$), High ($\sigma_e = 0.5$)
OM Covariate Process Error SD (σ_E)	0.1, 0.5
OM Covariate Process Error Correlation (ρ_E)	0, 0.5
EM Process Error	R, R+S, R+M
EM Covariate Effect	None ($\beta_E = 0$), β_E estimated
EM Median Natural Mortality Rate Parameter (β_M)	Known, Estimated

Table S2. Percent reduction in deviance for linear regression models fit to errors in estimation of the Hessian-based standard error of the estimates of the covariate effect on M ($\widehat{\text{SE}}(\hat{\beta}_E)$). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	1.84	2.68	4.00
OM Obs. Error	0.06	2.74	8.04
OM σ_e	6.27	7.28	12.83
OM σ_E	0.48	1.05	0.46
OM ρ_E	0.17	0.01	0.02
OM β_E	0.37	0.01	0.05
EM Process Error	2.90	3.89	2.98
EM M assumption	0.15	0.34	0.47
All factors	13.01	20.95	29.24
+ All Two Way	28.58	32.25	42.77
+ All Three Way	44.44	42.83	51.84

Table S3. Percent reduction in deviance for linear regression models fit to errors in estimation of the Hessian-based standard error of the estimates of the median M parameter ($\widehat{\text{SE}}(\hat{\beta}_M)$). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	0.55	12.91	3.65
OM Obs. Error	0.98	0.01	0.07
OM σ_e	<0.01	0.03	0.11
OM σ_E	0.13	0.03	0.19
OM ρ_E	<0.01	0.02	<0.01
OM β_E	0.12	0.07	0.23
EM Process Error	3.16	8.79	6.83
EM β_E assumption	0.18	8.96	3.81
All factors	5.32	28.39	14.71
+ All Two Way	11.80	44.67	22.70
+ All Three Way	17.83	51.54	29.46

Table S4. Percent reduction in deviance for linear regression models fit to log-transformed RMSE of terminal year natural mortality (M). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	0.26	2.21	14.65
OM Obs. Error	1.65	0.71	4.68
OM σ_e	0.02	0.16	0.01
OM σ_E	1.06	1.23	0.84
OM ρ_E	0.02	0.01	<0.01
OM β_E	3.73	1.94	1.14
EM Process Error	2.04	3.85	0.87
EM β_E assumption	0.19	0.59	0.07
EM M assumption	22.28	45.15	44.70
All factors	32.83	56.71	66.99
+ All Two Way	57.65	84.17	87.96
+ All Three Way	74.93	94.57	91.94

Table S5. Percent reduction in deviance for linear regression models fit to correlation of terminal year natural mortality (M) estimates and true values. Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	0.05	0.02	0.37
OM Obs. Error	2.00	0.89	3.73
OM σ_e	2.30	2.56	0.37
OM σ_E	14.91	13.78	3.03
OM ρ_E	1.98	0.33	0.14
OM β_E	0.41	3.76	3.87
EM Process Error	0.43	1.52	0.62
EM β_E assumption	12.72	12.30	1.56
EM M assumption	12.34	7.16	3.03
All factors	42.11	38.31	15.76
+ All Two Way	63.38	62.40	31.99
+ All Three Way	72.79	75.98	46.62

Table S6. Percent reduction in deviance for linear regression models fit to log-transformed RMSE of terminal year spawning stock biomass (SSB). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	19.88	32.67	20.92
OM Obs. Error	2.21	0.11	1.48
OM σ_e	0.05	0.16	0.03
OM σ_E	0.08	0.16	0.02
OM ρ_E	0.03	<0.01	0.02
OM β_E	0.05	0.03	0.14
EM Process Error	5.72	0.13	5.73
EM β_E assumption	0.03	0.79	0.05
EM M assumption	20.05	21.11	19.43
All factors	48.09	55.17	47.82
+ All Two Way	79.09	80.98	78.40
+ All Three Way	88.30	89.49	87.67

Table S7. Percent reduction in deviance for linear regression models fit to correlation of terminal year spawning stock biomass (SSB) estimates and true values. Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	19.27	25.70	19.59
OM Obs. Error	12.53	3.60	9.46
OM σ_e	0.08	0.44	0.09
OM σ_E	0.10	0.44	0.11
OM ρ_E	<0.01	<0.01	0.09
OM β_E	0.09	0.05	<0.01
EM Process Error	0.07	0.92	0.71
EM β_E assumption	<0.01	1.32	0.38
EM M assumption	28.94	19.74	25.36
All factors	61.19	52.21	56.09
+ All Two Way	81.48	72.16	74.98
+ All Three Way	85.23	83.22	80.48




Fig. S10. Estimated probability of fits providing Hessian-based standard errors for EMs assuming alternative process error, that estimate or assume known median natural mortality $\underline{M}$, and that estimate or assume no covariate effect on median natural mortality $\underline{M}$ when fitted to R OMs and three levels of true covariate effect on median natural mortality $\underline{M}$ (x axis). Vertical lines represent 95% confidence intervals




Fig. S11. Estimated probability of fits providing Hessian-based standard errors for EMs assuming alternative process error, that estimate or assume known median natural mortality $\underline{M}$, and that estimate or assume no covariate effect on median natural mortality $\underline{M}$ when fitted to R+S OMs and three levels of true covariate effect on median natural mortality $\underline{M}$ (x axis). Vertical lines represent 95% confidence intervals




Fig. S12. Estimated probability of fits providing Hessian-based standard errors for EMs assuming alternative process error, that estimate or assume known median natural mortality $\underline{M}$, and that estimate or assume no covariate effect on median natural mortality $\underline{M}$ when fitted to R+M OMs and three levels of true covariate effect on median natural mortality $\underline{M}$ (x axis). Vertical lines represent 95% confidence intervals




Fig. S13. Proportion of simulated data sets for R OMs where the EM type (treatment of environmental covariate and assumed process error type) had the lowest AIC.




Fig. S14. Proportion of simulated data sets for R+S OMs where the EM type (treatment of environmental covariate and assumed process error type) had the lowest AIC.

Fig. S15. Proportion of simulated data sets for R+M OMs where the EM type (treatment of environmental covariate and assumed process error type) had the lowest AIC.

Table S8. Percent reduction in deviance for linear regression models fit to errors in estimation of the covariate effect on M (β_E). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	<0.01	0.01	<0.01
OM Obs. Error	<0.01	<0.01	<0.01
OM σ_ϵ	<0.01	<0.01	<0.01
OM σ_E	<0.01	<0.01	<0.01
OM ρ_E	<0.01	<0.01	<0.01
OM β_E	<0.01	0.01	<0.01
EM Convergence	<0.01	<0.01	<0.01
EM Process Error	<0.01	0.01	0.01
EM M assumption	<0.01	<0.01	<0.01
All factors	0.02	0.03	0.03
+ All Two Way	0.16	0.20	0.18
+ All Three Way	0.58	0.73	0.59




Fig. S16. For R OMs, median error (ME) of estimates of environmental effect on natural mortality $M(\beta_E)$ from fitting EMs with alternative process error assumptions and treatment of median natural mortality M parameter (e_M^β known or estimated β_M). Vertical lines represent 95% confidence intervals.




Fig. S17. For R+S OMs, median error (ME) of estimates of environmental effect on natural mortality M (β_E) from fitting EMs with alternative process error assumptions and treatment of median natural mortality M parameter (e_M^{β} known or estimated β_M). Vertical lines represent 95% confidence intervals.




Fig. S18. For R+M OMs, median error (ME) of estimates of environmental effect on natural mortality $M(\beta_E)$ from fitting EMs with alternative process error assumptions and treatment of median natural mortality M parameter (e_M^β known or estimated β_M). Vertical lines represent 95% confidence intervals.

Median error (ME) of Hessian-based estimates of standard error for covariate effect on natural mortality β_E from fitting EMs with alternative process error assumptions and treatment of median natural mortality (e_M^β known or estimated). All OMs had low observation error and contrast in fishing mortality. True standard error is defined as the mean of the standard error estimates across converged fits to simulated data sets for a given OM scenario.




Fig. S16. E, R, OM results (ME) of Hessian-based estimates of standard error for covariate effect on natural mortality.




Fig. S20. For R+S OM, median error (ME) of Hessian-based estimates of standard error for covariate effect on natural mortality $M(\beta_E)$ from fitting EMs with alternative process error assumptions and treatment of median natural mortality M parameter (e_M^β known or estimated β_M). True standard error is defined as the mean of the standard error estimates across converged fits to simulated data sets for a given OM scenario.




Fig. S21. For R+M OMs, median error (ME) of Hessian-based estimates of standard error for covariate effect on natural mortality $\hat{M}(\beta_E)$ from fitting EMs with alternative process error assumptions and treatment of median natural mortality $\hat{M}$ parameter (e_M^β known or estimated β_M). True standard error is defined as the mean of the standard error estimates across converged fits to simulated data sets for a given OM scenario.




Fig. S22. For R OMs, probability of 95% confidence interval for β_E containing the true value for EMs with alternative process error assumptions and treatment of median natural mortality M parameter (e_M^β known or estimated β_M). Vertical lines represent 95% confidence intervals.




Fig. S23. For R+S OMs, probability of 95% confidence interval for β_E containing the true value for EMs with alternative process error assumptions and treatment of median natural mortality M parameter (e_M^β known or estimated β_M). Vertical lines represent 95% confidence intervals.




Fig. S24. For R+M OMs, probability of 95% confidence interval for β_E containing the true value for EMs with alternative process error assumptions and treatment of median natural mortality- M parameter (e_M^β known or estimated β_M). Vertical lines represent 95% confidence intervals.




Fig. S25. ~~Root-mean-square-error (Regression trees indicating primary factors determining reductions in sums of squares for the log-transformed RMSE) of estimates of covariate effect on natural mortality β_E from fitting EMs M ($RMSE(\hat{\beta}_E)$) for operating models (OMs) with alternative process error assumptions on recruitment only (R), recruitment and treatment of median apparent survival (R+S), or recruitment and natural mortality (e_M^β known or estimated R+M). All OMs had low observation~~ Each node shows the median error (top) and ~~contrast in fishing mortality~~ number of observations (bottom) for the corresponding subset. Lower or higher median RMSE are indicated by more green or red polygons, respectively.

Table S9. Percent reduction in deviance for linear regression models fit to the log-transformed RMSE for estimates of the covariate effect on M ($\text{RMSE}(\hat{\beta}_E)$). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	0.50	0.11	0.12
OM Obs. Error	12.04	5.66	0.36
OM σ_e	0.87	5.78	2.60
OM σ_E	15.85	24.18	24.26
OM ρ_E	0.36	0.82	0.34
OM β_E	6.72	0.07	0.96
EM Process Error	6.49	11.31	11.98
EM M assumption	0.04	0.64	0.07
All factors	42.87	48.58	40.65
+ All Two Way	67.34	74.09	61.74
+ All Three Way	79.42	82.75	71.92




Fig. S26. For R OMs, ~~root-mean-square-error (RMSE)~~ of estimates of covariate effect on ~~natural mortality~~ $\hat{M}(\beta_E)$ from fitting EMs with alternative process error assumptions and treatment of median ~~natural mortality~~ $\hat{M}$ parameter (e_M^β ~~known or estimated~~ $\hat{\beta}_M$).




Fig. S27. For R+S OMs, ~~root-mean-square-error (RMSE)~~ of estimates of covariate effect on ~~natural mortality~~ $\hat{M}(\beta_E)$ from fitting EMs with alternative process error assumptions and treatment of median ~~natural mortality~~ $\hat{M}$ parameter (e_M^β ~~known or estimated~~ $\hat{\beta}_M$).




Fig. S28. For R+M OMs, ~~root-mean-square-error (RMSE)~~ of estimates of covariate effect on ~~natural mortality~~ $\hat{M}(\beta_E)$ from fitting EMs with alternative process error assumptions and treatment of median ~~natural mortality~~ $\hat{M}$ parameter (e_M^β ~~known or estimated~~ $\hat{\beta}_M$).




Fig. S29. Positive correlation of covariate effect estimates and Hessian-based standard error estimates for EM that also estimates the median ~~natural mortality~~ $\mathcal{M}$ parameter ($\beta_{\mathcal{M}}$) and has correct R+M process error assumption fitted to simulated data from the OM with R+M process errors, temporal contrast in fishing pressure, low observation uncertainty for both population (*LowOE*) and covariate observations ($\sigma_e = 0.1$), high and uncorrelated temporal variability in the true covariate ($\sigma_E = 0.5$ and $\rho_E = 0$), and the strongest covariate effect on ~~natural mortality~~ $\mathcal{M}$ ($\beta_E = 0.5$).

Table S10. Percent reduction in deviance for linear regression models fit to errors in estimation of the median M parameter (β_M). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	<0.01	<0.01	<0.01
OM Obs. Error	<0.01	<0.01	<0.01
OM σ_e	<0.01	<0.01	<0.01
OM σ_E	<0.01	<0.01	<0.01
OM ρ_E	<0.01	<0.01	<0.01
OM β_E	<0.01	<0.01	0.01
EM Convergence	<0.01	0.01	0.01
EM Process Error	<0.01	0.01	0.01
EM β_E assumption	<0.01	<0.01	<0.01
All factors	0.02	0.02	0.05
+ All Two Way	0.15	0.15	0.24
+ All Three Way	0.57	0.54	0.73




Fig. S30. For R OMs, median error (ME) of estimates of β_M from fitting EMs with alternative process error assumptions and treatment of covariate effect ($\beta_E = 0$ or estimated). Vertical lines represent 95% confidence intervals.




Fig. S31. For R+S OMs, median error (ME) of estimates of β_M from fitting EMs with alternative process error assumptions and treatment of covariate effect ($\beta_E = 0$ or estimated). Vertical lines represent 95% confidence intervals.





Fig. S32. For R+M OMs, median error (ME) of estimates of β_M from fitting EMs with alternative process error assumptions and treatment of covariate effect ($\beta_E = 0$ or estimated). Vertical lines represent 95% confidence intervals.

Median-error (ME) of Hessian-based estimates of standard error for median natural mortality parameter β_M from fitting EMs with alternative process error assumptions and treatment of the covariate effect ($\beta_E = 0$ or estimated). All OM had low observation error and contrast in fishing mortality. True standard error is defined as the mean of the standard error estimates across converged fits to simulated data sets for a given OM scenario.






Fig. S34. For R+S OM, median error (ME) of Hessian-based estimates of standard error for median ~~natural mortality~~ $\hat{M}$ parameter ($\hat{\beta}_M$) from fitting EMs with alternative process error assumptions and treatment of the covariate effect ($\beta_E = 0$ or estimated). True standard error is defined as the mean of the standard error estimates across converged fits to simulated data sets for a given OM scenario.




Fig. S35. For R+M OM, median error (ME) of Hessian-based estimates of standard error for median ~~natural mortality~~ $\hat{\beta}_M$ parameter ($\hat{\beta}_M$) from fitting EMs with alternative process error assumptions and treatment of the covariate effect ($\beta_E = 0$ or estimated). True standard error is defined as the mean of the standard error estimates across converged fits to simulated data sets for a given OM scenario.




Fig. S36. For R OMs, probability of 95% confidence interval for β_M containing the true value for EMs with alternative process error assumptions and treatment of covariate effect ($\beta_E = 0$ or estimated). Vertical lines represent 95% confidence intervals.




Fig. S37. For R+S OMs, probability of 95% confidence interval for β_M containing the true value for EMs with alternative process error assumptions and treatment of covariate effect ($\beta_E = 0$ or estimated). Vertical lines represent 95% confidence intervals.




Fig. S38. For R+M OMs, probability of 95% confidence interval for β_M containing the true value for EMs with alternative process error assumptions and treatment of covariate effect ($\beta_E = 0$ or estimated). Vertical lines represent 95% confidence intervals.

1101 Median natural mortality parameter estimate and standard error
1102 example




Fig. S39. Negative correlation of β_M estimates and Hessian-based standard error estimates for EM that also estimates the covariate effect and has correct R+M process error assumption fitted to simulated data from the OM with R+M process errors, temporal contrast in fishing pressure, low observation uncertainty for both population (*LowOE*) and covariate observations ($\sigma_e = 0.1$), high and uncorrelated temporal variability in the true covariate ($\sigma_E = 0.5$ and $\rho_E = 0$), and the strongest covariate effect on ~~natural mortality~~ $\mathcal{M}$ ($\beta_E = 0.5$).




Fig. S40. ~~Root-mean-square-error~~ (Regression trees indicating primary factors determining reductions in sums of squares of errors for the log-transformed RMSE) of estimates of β_M from fitting for the median M parameter (RMSE(β_M)) in EMs fitted to operating models (OMs) with ~~alternative~~ process error assumptions on recruitment only (R), recruitment and ~~treatment of covariate effect apparent survival~~ ($\beta_E = 0$ R+S), or ~~estimated~~ recruitment and natural mortality (R+M). ~~All OMs had low observation~~ Each node shows the median error for population observations (top) and ~~contrast in fishing mortality~~ number of observations (bottom) for the corresponding subset. Lower or higher median RMSE are indicated by more green or red polygons, respectively.

Table S11. Percent reduction in deviance for linear regression models fit to the log-transformed RMSE for estimates of the median M parameter ($\text{RMSE}(\hat{\beta}_M)$). Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	34.70	24.62	34.91
OM Obs. Error	25.55	17.82	12.77
OM σ_ϵ	0.01	≤ 0.01	≤ 0.01
OM σ_E	0.19	0.02	0.07
OM ρ_E	0.05	0.26	0.03
OM β_E	0.58	0.23	0.16
EM Process Error	10.75	18.12	14.50
EM β_E assumption	3.96	6.39	2.93
All factors	75.79	67.47	65.26
+ All Two Way	90.21	80.65	77.59
+ All Three Way	93.72	88.68	86.44




Fig. S41. For R OMs, ~~root-mean-square-error~~ (RMSE) of estimates of β_M from fitting EMs with alternative process error assumptions and treatment of covariate effect ($\beta_E = 0$ or estimated).




Fig. S42. For R+S OMs, ~~root-mean-square-error~~ (RMSE) of estimates of β_M from fitting EMs with alternative process error assumptions and treatment of covariate effect ($\beta_E = 0$ or estimated).




Fig. S43. For R+M OMs, root-mean-square-error (RMSE) of estimates of β_M from fitting EMs with alternative process error assumptions and treatment of covariate effect ($\beta_E = 0$ or estimated).




Fig. S44. For R OM, median relative error (MRE) of estimates of ~~natural mortality rate in the~~ terminal year $\hat{M}$ for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ $\hat{M}$ parameter (β_M estimated or known).




Fig. S45. For R+S OM, median relative error (MRE) of estimates of ~~natural mortality rate in the~~ terminal year $\hat{M}$ for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ $\hat{M}$ parameter (β_M estimated or known).




Fig. S46. For R+M OM, median relative error (MRE) of estimates of ~~natural mortality rate in the~~ terminal year $\hat{M}$ for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ $\hat{M}$ parameter (β_M estimated or known).




Fig. S47. Root mean square error (For R OMs, RMSE) of estimates of natural mortality rate in the terminal year $\hat{M}$ for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median natural mortality $\hat{M}$ parameter (β_M estimated or known). All OMs had low population observation error and contrast in fishing mortality.




Fig. S48. For R+S OMs, root-mean-square-error (RMSE) of estimates of natural mortality rate in the terminal year $\hat{M}$ for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median natural mortality M parameter (β_M estimated or known).




Fig. S49. For R+S-M OM, root-mean-square-error (RMSE) of estimates of natural mortality rate in the terminal year M for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median natural mortality M parameter (β_M estimated or known).




Fig. S50. For R +M-OMs, ~~root-mean-square-error (RMSE)~~ correlation of estimates ~~and true values~~ of ~~natural mortality rate in~~ the terminal year $\hat{M}$ for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ $\hat{M}$ parameter (β_M estimated or known).




Fig. S51. For R+S OM, correlation of estimates and true values of terminal year M for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median M parameter (β_M estimated or known).




Fig. S52. For R+M OMs, correlation of estimates and true values of terminal year M for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median M parameter (β_M estimated or known).




Fig. S53. For R OMs, median relative error (MRE) of estimates of ~~spawning-stock-biomass~~(SSB) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural-mortality~~ M parameter (β_M estimated or known).




Fig. S54. For R+S OMs, median relative error (MRE) of ~~estimates of spawning stock biomass (SSB)~~ in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ M parameter (β_M estimated or known).




Fig. S55. For R+M OMs, median relative error (MRE) of estimates of spawning-stock-biomass (SSB) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median natural mortality (β_M estimated or known).




Fig. S56. ~~Root-mean-square-error (For R OMs, RMSE)~~ of estimates of ~~spawning stock biomass~~ SSB in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ M parameter (β_M estimated or known). ~~All OMs had low population observation error and contrast in fishing mortality.~~




Fig. S57. For R+S OMs, root-mean-square-error (RMSE) of estimates of spawning stock biomass (SSB) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median natural mortality M parameter (β_M estimated or known).




Fig. S58. For R+S-M OM, root-mean-square-error (RMSE) of estimates of spawning stock biomass (SSB) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median natural mortality (M) parameter (β_M estimated or known).




Fig. S59. For R +M-OMs, root-mean-square-error (RMSE)-correlation of estimates and true values of spawning-stock-biomass in the terminal year SSB for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median natural-mortality-M parameter (β_M estimated or known).




Fig. S60. For R+S OM, correlation of estimates and true values of terminal year SSB for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median M parameter (β_M estimated or known).




Fig. S61. For R+M OM, correlation of estimates and true values of terminal year SSB for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median M parameter (β_M estimated or known).




Fig. S62. **Median relative error (MRE)** Regression trees indicating primary factors determining reductions in sums of **estimates squares** of **fully-selected-fishing-mortality (F) errors** in estimation measured by Eq. 2 for the terminal year **for-fishing-mortality rate** in EMs fitted to operating models (OMs) with **alternative-process error assumptions** on recruitment only (R), **treatment-of-covariate-effect-recruitment and apparent survival** ($\beta_E = 0$ or estimated R+S), or recruitment and **treatment-of median-natural-mortality parameter** (β_M estimated or known R+M). All-OMs had low-population-observation-Each node shows the median error (top) and **contrast-in-fishing-mortality** number of observations (bottom) for the corresponding subset. Lower or higher median absolute errors of the process error assumption are indicated by more green or red polygons, respectively.

Table S12. Percent reduction in deviance for linear regression models fit to errors in estimation measured by Eq. 2 for the terminal year fishing mortality rates. Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
Convergence	<0.01	0.19	0.01
OM F history	1.92	2.42	1.94
OM Obs. Error	0.74	0.12	1.12
OM σ_e	0.01	<0.01	<0.01
OM σ_E	<0.01	<0.01	<0.01
OM ρ_E	<0.01	<0.01	<0.01
OM β_E	<0.01	0.01	<0.01
EM Process Error	0.40	0.92	0.48
EM β_E assumption	0.01	0.03	0.02
EM M assumption	1.58	1.85	1.68
All factors	4.95	5.82	5.35
+ All Two Way	10.91	11.03	12.23
+ All Three Way	14.16	13.24	16.19




Fig. S63. For R OMs, median relative error (MRE) of estimates of fully-selected fishing mortality (F) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ M parameter (β_M estimated or known). All OMs had low observation error and contrast in fishing mortality.




Fig. S64. For R+S OMs, median relative error (MRE) of estimates of fully-selected fishing mortality (F) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median natural-mortality- M parameter (β_M estimated or known). All OMs had low observation error and contrast in fishing mortality.




Fig. S65. For R+M OMs, median relative error (MRE) of estimates of fully-selected fishing mortality (F) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ $\mathcal{M}$ parameter (β_M estimated or known). All OMs had low observation error and contrast in fishing mortality.




Fig. S66. **Root mean square error (RMSE)** Regression trees indicating primary factors determining reductions in sums of estimates squares of fully-selected fishing mortality (F) in errors for the log-transformed RMSE of terminal year for fishing mortality rate in EMs fitted to operating models (OMs) with alternative process error assumptions on recruitment only (R), treatment of covariate effect recruitment and apparent survival ($\beta_E = 0$ or estimated R+S), or recruitment and treatment of median natural mortality parameter (β_M estimated or known R+M). All OMs had low population observation. Each node shows the median error (top) and contrast in fishing mortality number of observations (bottom) for the corresponding subset. Lower or higher median RMSE are indicated by more green or red polygons, respectively.

Table S13. Percent reduction in deviance for linear regression models fit to log-transformed RMSE of terminal year fishing mortality rate. Table rows give each operating model (OM) and estimating model (EM) factor included individually, combined, and with second and third order interactions. Table columns are operating models (OMs) with process error on recruitment only (R), recruitment and apparent survival (R+S), or recruitment and natural mortality (R+M).

Factor	R	R+S	R+M
OM F history	11.29	25.63	13.46
OM Obs. Error	37.27	21.42	35.47
OM σ_ϵ	<0.01	0.02	0.02
OM σ_E	0.09	0.03	0.14
OM ρ_E	0.02	0.01	0.04
OM β_E	0.15	0.03	0.21
EM Process Error	0.42	0.75	0.63
EM β_E assumption	0.04	<0.01	0.04
EM M assumption	25.38	26.10	24.68
All factors	74.65	73.99	74.76
+ All Two Way	92.28	91.29	91.64
+ All Three Way	94.31	95.51	93.73




Fig. S67. For R OMs, ~~root-mean-square-error~~ (RMSE) of estimates of fully-selected fishing mortality (F) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ M parameter (β_M estimated or known).




Fig. S68. For R+S OMs, ~~root-mean-square-error~~ (RMSE) of estimates of fully-selected fishing mortality (F) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ M parameter (β_M estimated or known).




Fig. S69. For R+M OMs, ~~root-mean-square-error~~ (RMSE) of estimates of fully-selected fishing mortality (F) in the terminal year for EMs with alternative process error assumptions, treatment of covariate effect ($\beta_E = 0$ or estimated), and treatment of median ~~natural mortality~~ M parameter (β_M estimated or known).