

Formulation

Let:

$x_1, x_2 \dots x_m$ = amount (in grams) of each food item in the USDA rolled-up list

α_{ij} = amount (in grams) of nutrient j in food item i

L_j = lower nutritional bound for all nutrients $j = 1 \dots 32$

U_j = upper nutritional bound for all nutrients $j = 1 \dots 32$

Linear Programming Model:

For low-carb diet:

minimize: $z = \sum_{i=1}^{i=m} x_{ij_1}$ where j_1 = carbohydrates

For low-carb, low-sodium, low-cholesterol diet:

minimize: $z = \sum_{i=1}^{i=m} x_{ij_1} + x_{ij_2} + x_{ij_3}$ where j_1 = carbohydrates, j_2 = sodium, j_3 = cholesterol

Subject to:

$x_1, x_2 \dots x_m \geq 0$ non-negativity constraint

$\sum_{i=1}^{i=m} \alpha_{ij} x_{ij} \geq L_j, j = 1 \dots 32$ sum of nutrient j for all foods i must meet the minimum nutritional requirement L

$\sum_{i=1}^{i=m} \alpha_{ij} x_{ij} \leq U_j, j = 1 \dots 32$ sum of nutrient j for all foods i must not exceed the maximum nutritional requirement U