

Individual-level causes and population-level consequences of variation in fitness in an alpine rodent

Dissertation
zur
Erlangung der naturwissenschaftlichen Doktorwürde
(Dr. sc. nat.)
Mathematisch-naturwissenschaftlichen Fakultät
der
Universität Zürich
von

Timothée Bonnet

aus
Frankreich

Promotionskomitee

Prof. Dr. Lukas Keller (Vorsitz)
Dr. Erik Postma (Leitung der Dissertation)
Prof. Dr. Ben Sheldon
Prof. Dr. Barbara Tschirren
Prof. Dr. Arpat Ozgul
Dr. Jarrod Hadfield
Dr. Marc Kéry

Zürich, 2016

©Timothée Bonnet

timotheebonnet@gmail.com

Photos: Timothée Bonnet

This is a L^AT_EX document using a customized version of the KOMA-Script's class
scrbook

Summary

This thesis investigates the stochastic and selective causes of variation in fitness components, and the evolutionary consequences of this variation in a wild rodent population. It shows the contemporary genetic evolution of body mass and decouples classic estimates of selection from adaptive evolution.

The heart of evolutionary biology is understanding the variation in organisms. For over 150 years, researchers have documented the causes of within-species variation and how it contributes to speciation and explains the fit between organisms and their environment. Recently, increasing concerns regarding rapid anthropogenic changes have driven renewed investigation of how wild populations adapt to environmental change. This new focus has revealed the difficulties measuring natural selection, disentangling evolution from plastic changes, and predicting evolutionary trajectories. For instance, there are few robust examples of contemporary evolution in wild populations, casting doubt on the possibility that evolution can rescue populations from rapid environmental change. In this thesis, I investigate the causes of natural selection and evolution in a wild population of snow voles (*Chionomys nivalis*). Thanks to 10 years of intensive individual-based monitoring and genotyping, knowledge of this population includes life-history, morphological data, and a high-resolution pedigree. This population is therefore among the best available worldwide to measure selection and evolution in action.

The population is nevertheless relatively small and recent publications suggest that the evolutionary potential in small populations is effectively cancelled by stochasticity in fitness components. I assess the methods used in those publications and demonstrate that the variation in fitness components is not purely stochastic. Small populations, including these snow voles, show evolutionary potential.

With collaborators, I then compare four common methodological frameworks to disentangle the contributions to phenotypic change of evolution, plasticity, and demography. We identify important discrepancies between the frameworks, partly originating from using different definitions, but also possessing intrinsically different capabilities. Among the considered frameworks only quantitative genetics can measure genetic change.

Applying methods from quantitative genetics to the snow vole population, I demonstrate that body mass evolved adaptively over the study period. I show that phenotypic estimates of selection are not predictive of genetic evolution: neither the mean selection nor its temporal variation are related to the rate of genetic evolution. This demonstrates that the dominant purely-phenotypic method used to measure selection risks measuring variation in nutritional status instead. Nevertheless, I employed quantitative genetics to identify the target of selection and obtain selection estimates in line with the observed genetic change

This thesis establishes contemporary evolution in a wild population and shows that



evolutionary responses to environmental change cannot be reliably estimated nor understood from purely-phenotypic methods; an explicit genetic approach is necessary.



Zusammenfassung

Diese Doktorarbeit untersucht die stochastischen und selektiven Ursachen der Variation in Fitnesskomponenten und deren evolutionären Konsequenzen in einer freilebenden Nagetierpopulation. Sie zeigt die gegenwärtige, genetische Evolution von Körpermasse und entkoppelt klassische Selektionsschätzungen von adaptiver Evolution.

Das Herzstück der Evolutionsbiologie liegt im Verständnis der Vielfalt von Organismen. Während über 150 Jahren haben Forscher die Ursachen von intraspezifischer Variation dokumentiert, wie sie zur Artbildung beiträgt und zum Zusammenpassen von Organismen mit ihrer Umwelt. Zunehmende Bedenken wegen der schnellen anthropogenischen Veränderungen haben in letzter Zeit eine erneute Erforschung, wie sich freilebende Populationen an Umweltveränderungen anpassen, vorangetrieben. Dieser neue Fokus offenbart die Schwierigkeiten im Messen von natürlicher Selektion, die Entflechtung von Evolution und plastischen Veränderungen und dem Vorhersagen von evolutionären Entwicklungsverläufen. Unter anderem gibt es nur wenige robuste Beispiele von gegenwärtiger Evolution in freilebenden Populationen, die, die Möglichkeit, dass Evolution Populationen bei schnellen Veränderungen der Umweltbedingungen rettet, fraglich erscheinen lässt. In dieser Doktorarbeit erforsche ich die Ursachen natürlicher Selektion und Evolution in einer freilebenden Population von Schneemäusen (*Chionomys nivalis*). Dank 10 Jahren intensivem Individuenbasiertem Monitoring und Genotypisierung, beinhaltet der Erkenntnisstand dieser Population Lebensweise, morphologische Daten und einen hochaufgelösten Stammbaum. Deswegen ist diese Population unter den besten weltweit verfügbaren um Selektion und Evolution in Aktion zu messen.

Trotzdem ist die Population ziemlich klein und neuerliche Publikationen legen nahe, dass das evolutionäre Potential in kleinen Populationen effektiv von Stochastik in Fitnesskomponenten aufgehoben wird. Ich beurteile diese Methoden, die in diesen Publikationen benutzt wurden und demonstriere, dass die Variation in Fitnesskomponenten nicht ausschliesslich stochastisch ist. Kleine Populationen, einschliesslich diese Schneemäuse, zeigen evolutionäres Potential.

Mit Kollaboratoren vergleiche ich dann vier häufig benutzte methodologische Ansätze um die Anteile von Evolution, Plastizität und Demographie an der phänotypischen Veränderung zu entflechten. Wir identifizieren wichtige Unstimmigkeiten zwischen den Ansätzen, die teilweise vom Gebrauch von unterschiedlichen Definitionen, aber auch vom Besitz von intrinsisch unterschiedlichen Fähigkeiten stammen. Unter den in Betracht gezogenen Ansätzen kann nur quantitative Genetik genetische Veränderungen messen.

Durch die Anwendung von quantitativ-genetischen Methoden an der Schneemaus-Population demonstriere ich, dass sich Körpermasse über die Studiendauer adaptiv entwickelt. Ich zeige auf, dass phänotypische Schätzungen von Selektion nicht

genetische Evolution hervorsagen: weder die durchschnittliche Selektion noch die temporale Variation hängen mit der Rate genetischer Evolution zusammen. Das legt dar, dass die dominante, rein phänotypische Methode zur Messung von Selektion stattdessen die Messung von Variation im Ernährungszustand riskiert. Dennoch habe ich quantitative Genetik zur Identifikation von Selektion verwendet und Selektionsschätzungen erhalten, die mit der beobachteten genetischen Veränderung übereinstimmen.

Diese Doktorarbeit weist gegenwärtige Evolution in einer freilebenden Population nach und zeigt dass evolutionäre Reaktionen auf Umweltveränderungen von rein phänotypischen Methoden weder zuverlässig eingeschätzt noch verstanden werden können; ein expliziter, genetischer Ansatz ist notwendig.

Contents

Summary	i
Zusammenfassung	iii
Chapter 1	
General introduction	1
Addresses of co-authors	17



Chapter 1

General introduction

One can't understand everything at once, we can't begin with perfection all at once! In order to reach perfection one must begin by being ignorant of a great deal. And if we understand things too quickly, perhaps we shan't understand them thoroughly.

— Fyodor Dostoyevsky, *The Idiot* (1868–9)

Si nous ne trouvons pas des choses agréables, nous trouverons du moins des choses nouvelles. / If we do not find anything very pleasant, at least we shall find something new.

— Voltaire a.k.a. François-Marie Arouet, *Candide* (1759)

1.1 Variation in fitness

1.1.1 The origin of variation in evolutionary biology

The heart of evolutionary questioning is understanding variation among living beings (Lynch and Walsh 1998; Wayne and Miyamoto 2006; Kruuk, Charmantier, and Garant 2014). It is in fact its very starting point. Darwin opens his book *the Origin of Species* with two chapters describing variability in domestic and wild organisms (Darwin 1859). Building on these observations, Darwin then summarizes the evidence showing that variation within species is the fuel generating the astonishing diversity among species, and the striking fit between organisms and their environment.

These great answers immediately opened many more questions about the causes and consequences of variation, some of which remain not fully answered more than 150 years later. In particular, nineteenth century biologists struggled with the sources of variation within species, as is clear in *the Origin of Species* itself: “*Variability is governed by many unknown laws, of which correlated growth is probably the most important. Something, but how much we do not know, may be attributed to the definite action of the conditions of life. Some, perhaps a great, effect may be attributed to the increased use or disuse of parts*” (p. 31 Darwin 1859)¹. The environment was sometimes thought to play a predominant role, through what would be nowadays called *plastic* effects, and the effect of ageing was acknowledged (Wilkins 2009), but these did not always provide a satisfactory explanation for variation that appears within a population. Further-

¹ Alternatively, biologists dismissed this within-species variation by considering that species were arbitrary boundaries in a set or continuum of variation. Darwin did not attempt to define species, but by explaining how they originate, he made some definitions indefensible (Wilkins 2009, pp. 129-163).

more, Darwinian arguments build on the observation of this special kind of inherited variation that can appear among siblings of a same litter, clutch, or pod, and that is subsequently transmitted from parent to offspring (Darwin 1859, Chapter 1). The late nineteenth century was utterly ignorant of the sources of inherited variation within species. Only at the beginning of the twentieth century were the laws of inheritance progressively discovered and spread to the scientific community (Dietrich 2006). Four more decades saw these laws formalized into a unified scientific theory to understand variation within populations (R. Fisher 1930), and explained at the molecular level (Oswald, MacLeod, and McCarty 1944; Watson and Crick 1953), thus closing the logical gap in Darwin's argument: Relatives resemble each other because they share similar gene versions on long strands of DNA, a molecule that is copied with high fidelity and transmitted from parent to offspring; There is variation among siblings because of the reshuffling and segregation of parental genes and, on occasions, because DNA mutates.

The understanding of the causes of variation within species and populations has made terrific progresses and now fits elegantly in the broader evolutionary theory (Pigliucci and Müller 2010). Nevertheless, many aspects of the causes of variation are still to be refined or newly explored, especially in natural populations (Kruuk, Charmantier, and Garant 2014). In particular, the relative importance of genes and the environment in the wild remains studied in only a few populations of a few species, taxonomically biased, and concerns a limited set of traits (Lynch and Walsh 1998; Postma 2014). Additional open research questions relate to the consequences of within-species variation. In particular, a lot of attention is paid to how genetic variation translates into adaptive evolution (Brookfield 2016). Any trait that possesses genetic variation can evolve, but it will evolve in an adaptive way only if the trait is subject to selection, be it artificial or natural. Selection occurs when the variation in the trait causes variation in *fitness*. Before we discuss the specificity of the causes and consequences of variation in fitness, we must introduce this difficult concept.

1.1.2 Variation in the definition of fitness

There has been a great deal written about the concept of fitness, including multiple conflicting definitions, which “*is hardly surprising as every important scientific concept is difficult to understand from first principles, as for instance the notions of space and time, or energy and force*” (p. 1358 Wagner 2010). I will not solve the question of the definition of fitness here, but I will try to make clear how the word is used in this thesis.

To start with, in the past, there has been some confusion on whether fitness is a realized reproductive outcome or a propensity to reproduce (Brandon and Beatty 1984). It is now rather consensual that the concept of fitness is more useful when it is defined as a propensity, that is, as an expected value that cannot be measured directly because of stochasticity (Brandon and Beatty 1984; Price 1996; Krimbas 2004) and we will follow this consensus. Fitness has been defined at the level of the genetic lineage (e.g. Akçay and Van Cleve 2016), of the individual (e.g. Cam and Monnat 2000), of the genotype (e.g. Steiner and Tuljapurkar 2012), or of the population (e.g. Tienterden 2000). A propensity definition partly dissolves the problem of the level of the definition, since the expected reproductive outcome of a genotype is the same as the

expected reproductive outcome of the individuals bearing this genotype, and the expected reproductive outcome of a population is the sum of the expectation of the reproductive outcome of its individuals. Here, we will consider fitness at the level of individuals, because they are the unit most easily observable and the primary target of natural selection.

More confusion on fitness comes from it being alternatively defined as the asymptotic number of descendants or as the contribution to the next generation (Wade 2006). Since we consider fitness at the level of individuals and because most of the work carried out is based on data covering about ten generations only, it is intuitive to consider fitness as the contribution to the next generation. Besides practical considerations, this choice allows for a clear, and conceptually crucial, distinction between selection, inheritance and evolution, that is blurred in asymptotic definitions (R. Fisher 1930; Arnold and Wade 1984). Indeed, imagine the simplistic case of a genetic locus with two non-recombining haplotypes in a closed population. Assuming there is no balancing selection, one of the haplotypes will eventually go to fixation, its asymptotic fitness is one, and the other one will go extinct, its asymptotic fitness is zero. Fixation might have happened due to genetic drift, due to selection on the initial differences between the haplotypes, or due to selection on new mutations. Observing only the starting point (two haplotypes), and the end point (one haplotype), perfectly describes the long-term evolution, but does not reveal its mechanisms. To understand the respective roles of chance (drift), selection, and inheritance (mutations), one must describe the generation-to-generation changes in haplotype frequencies with respect to the allelic state of both haplotypes.

A slightly contentious point is whether fitness should be defined as an absolute number of offspring (Wade 2006) or a relative one (Rousset 2004), that is, whether “relative fitness” is a meaningful phrase or a tautological one. The relative definition avoids appending *relative* to every occurrence of *fitness*, and seems closer to the interest of evolutionary biologists. Nevertheless, the field massively favours the absolute definition, possibly because it has a concrete and observable meaning. For the sake of consistency I attempted to yield to the convention (possibly with some inconsistencies).

Finally, instead of a measure of reproductive success, relative fitness has recently been defined as the amount of information about the environment that populations accumulate by selection (Frank 2012). The first and secondary theorems of natural selection can be rigorously written in term of gain and loss of bits of information about the environment. This shows that populations gain information by selection, and lose it by imperfect transmission or environmental change. I see great conceptual promises in this view, that brings together an intuitive meaning of the word *fitness* and the scientific field of information theory, with all its powerful tools and concepts. An information interpretation of fitness did not directly influence the work and is not necessary to understand it, but it might enlighten some of the results presented here, and evolutionary biology in general. To sum up, we define the fitness of an individual as its expected number of descendant in the next generation.

1.1.3 Causes and consequences of fitness variation in the wild

Why is there variation in individual fitness? This question attracts a lot of research attention, because (i) genetic variation in fitness controls the pace of evolution within a population, and because (ii) an intuitive consequence of evolution is the erosion of genetic variation in fitness, thus making the presence of genetic variation in fitness paradoxical (Jones 1987). In this thesis, we only deal briefly with the second point, the fundamental question of appearance and maintenance of genetic variation in fitness, in chapter ???. On the other hand, all the chapters deal with the proximal sources of variation in fitness, and we will consider these proximal sources from two complementary angles.

First in a descriptive approach, one can decompose variation in fitness, i.e. the opportunity for selection, into components of variation. Apart from genetic variation, variation in fitness can also originate from variation in early-life, micro-environment (Turner 2009), or maternal effects (Wolf and Wade 2009). In addition, when working with wild sexual organisms, individual fitness as we defined it cannot be observed directly. Indeed, individuals are unique and their realized reproductive success does not equal their expected reproductive success. Therefore, researchers have to rely on fitness proxies, often realized reproductive success and survival, that contain a large stochastic component. Additive genetic variation in fitness is also the rate of evolution in fitness and sets the maximal rate of evolution (R. Fisher 1930). A variance decomposition approach is therefore useful to ascertain whether variation in fitness proxies is all stochastic and environmental or whether it hides genetic variation in fitness. In doing so, it determines how much adaptive evolution can be expected to happen within a population.

Second, in a more mechanistic approach, one can investigate what characteristics make some individuals fitter than others, that is what traits are under natural selection². The study of natural selection in the wild took-off with the development of regression-based methods to accurately measure its strength and predict its effects (Lande 1979; Lande and Arnold 1983). Under some assumptions, the genetic change in response to selection on a trait is the product of a selection gradient and of additive genetic variation in that trait (Lush 1937). Therefore, by understanding what traits cause variation in fitness, one can predict what traits should evolve, as well as their direction and speed of evolution.

The study of natural selection and adaptive evolution in the wild is very topical in the context of unprecedented rates of environmental changes induced by human activities (Parmesan 2006). Anthropogenic changes provide the opportunity of natural experiments to evolutionary biologists (Altermatt, Ebert, and Altermatt 2016; Brookfield 2016), but also come with societal concerns and an ever increasing urge to better understand and predict how living things respond to the selective pressures imposed by environmental changes (Mc Carty 2001; Shaw and Shaw 2014). This regain of focus

²In this thesis, unless mentioned otherwise, we consider *sexual selection* as part of *natural selection* and of *selection*. Measuring sexual and natural selection separately, would certainly provide a finer understanding of the mechanisms of selection in the study population, but this was beyond the scope of this thesis. Nevertheless, the question was partly explored by García-Navas, Bonnet, Waldvogel, et al. 2016 and García-Navas, Bonnet, Bonal, et al. 2016.

has highlighted the gaps in the understanding of adaptation in natural populations: it is still challenging to predict, or even understand retrospectively, how natural populations respond to selective pressures (Merilä, Sheldon, and Kruuk 2001; Tafani et al. 2013; Shaw and Shaw 2014; Brookfield 2016).

In order to study the evolutionary potential of wild populations and their response to selective pressure, it is necessary to measure genetic parameters. More specifically, one must determine whether the traits under selection are heritable, whether there is heritable variation in fitness and how what is the rate of genetic change for the traits of interest.

1.2 Measuring genetic variation

1.2.1 Looking up or down? Two philosophies

How to measure and make sense of genetic variation? For over a century, there have been two main approaches (Liedvogel, Cornwallis, and Sheldon 2012), that can grossly be traced back to the scientific controversy that opposed the Mendelians to the biometricalists (Dietrich 2006), and summarized as “bottom-up” and “top-down”. Bottom-up approaches, embodied by candidate gene and genome wide association studies, start from molecular data to infer the phenotypic effects of individual genetic loci. Top-down approaches, encompassed within quantitative genetics, attempt to decompose phenotypic variation into genetic variation and other sources of variation, based solely on phenotypic data and on some knowledge of the relatedness between individuals (Lynch and Walsh 1998). Some pros and cons of both approaches are nicely illustrated by the confrontation of the quantitative genetics of mass with the genotyping of a candidate gene for mass. The former will be further developed in chapters ?? and ?? and we present it in a minimalist nutshell here: using a quantitative genetics *animal model* (Henderson 1950; Kruuk 2004), we estimated additive genetic variation in body mass and lifetime reproductive success. The candidate gene approach is a side project of this PhD that does not appear in the other chapters, and we take the opportunity to present it below.

1.2.2 A candidate gene for body mass: insights and limits

We used a candidate gene approach (Fitzpatrick et al. 2005) to uncover a molecular mechanism underlying variation in body mass. To date, the only candidate gene we fully analysed is an intronic region of the gene *lepr*, which codes for the receptor to leptin. Leptin is a hormone known to regulate fat metabolism, energy expenditure and food intake, including in rodents (Houseknecht et al. 1998).

We found a recessive allele (let call the recessive allele *a*, and the dominant allele *A*) associated with lighter individuals (Fig. 1.1A). Homozygotes *aa* were -2.9 g lighter (95% credibility interval [0.6; 5.1]), that is, 8% lighter than the mean. On average, during their lifetime, these *aa* individuals produced one third less offspring than the *AA* individuals (Fig. 1.1B). This strong difference in fitness is however not statistically significant. These results suggest that some of the genetic variation in body mass is due

to food intake and/or fat metabolism, which could not be sensed from the estimation of genetic variances and covariances. Based on such a strong phenotypic effect, *lepr* could be called a major locus, but how much of the genetic variation does it explain?




Figure 1.1: Body mass and lifetime reproductive success (LRS) as a function of *lepr* genotypes. (A) Expected body mass of snow voles bearing the three *lepr* genotypes. The expectations and 95% confidence intervals were predicted from a linear mixed model fitted to the 2311 mass measurement of 532 snow voles. The model accounted for sex, age, date of capture and their two-ways interactions, as well as year of capture and multiple measurements of the same individual. (B) Expected LRS of snow voles bearing the three *lepr* genotypes. The expectations and 95% confidence intervals were predicted from a Poisson generalized linear mixed model fitted to the LRS of 611 snow voles. The model accounted for inbreeding coefficient, year of birth and over-dispersion (using an observation-level random effect). For both panels, the dashed horizontal line projects the expected value of genotype AA to ease comparison with *Aa* and *aa*.

Knowing the effect of the three genotypes and the allele frequencies one can compute analytically the additive genetic variances associated with a bi-allelic locus (R. A. Fisher 1941; Lynch and Walsh 1998, p77). Thus, the additive genetic variances associated with *lepr* are 0.052g^2 for body mass and 0.006pup^2 for lifetime reproductive success. For both traits, *lepr* explains about 1% of the additive genetic variation as estimated from an animal model. This is rather large for a single locus given that quantitative traits loci typically explain a fraction of a percent to a few percent of additive genetic variance (V_A), when they have a large enough sample size to mitigate Beavis effect (Flint and Mackay 2009; Jensen, Szulkin, and Slate 2014). Still, 1% of V_A is not sufficient to infer the evolutionary potential of the trait. Finally, genotyping many more markers, for instance using high-throughput sequencing (Goodwin, McPhereson, and McCombie 2016), is unlikely to improve this situation in the snow vole pop-



ulation. Generally, very large sample sizes and high-quality genomic resources are necessary to explain a biologically proportion of genetic variances (Bloom et al. 2013; Jensen, Szulkin, and Slate 2014). For instance 183,727 individuals were necessary to find 180 QTL that jointly explained only 13% of additive genetic variation in human body height (Lango Allen et al. 2010). High-throughput sequencing can also been used in a top-down way, that does not identify causal genetic variants, but instead quantifies the phenotypic variation jointly explained by all the genotyped markers. Thus, 3,925 individuals and 294,831 markers were able to explain 45% of the genetic variation in human height (Yang et al. 2010). This is much better, but given knowledge on the relatedness between individuals, quantitative genetics can estimate all the genetic variation in a phenotype, without any genotyping effort.

To conclude, bottom-up approaches can better unravel the molecular mechanisms underlying phenotypes. By opening the black box of what mechanisms make up a phenotype, they can identify what is most crucial in a phenotype, how it is linked to the environment and what is the target of natural selection (Jong et al. 2014). Moreover, they contribute to building a genotype-phenotype map, a long lasting challenge in evolutionary biology (Kirschner and Gerhart 2010). On the other hand, quantitative genetics lump all the effects of individual genes and their interactions into only a few parameters, non-informative about the underlying genetic architecture (Mackay 2001; Nietlisbach and Hadfield 2015; Huang and Mackay 2016). This summarized estimation provides simple and direct measures of genetic parameters. Quantitative genetics work directly on the phenotype which is the target of selection, and the source of ecological interactions. They therefore provide simple measures of genetic parameters that can directly be interpreted within the ecology of organisms. This thesis is concerned with the genetics and evolution at the level of organisms, in relation to their environment, and accordingly, most of my work relies on quantitative genetics.

1.3 This thesis

1.3.1 Objectives

In this thesis, I investigate the causes of individual-level variation in fitness, and the consequences of this variation at the population level. This thesis aims at better measuring and better understanding selection and evolution in the wild. It examines the relative importance of stochasticity and selection in shaping reproductive success and survival, disentangles evolutionary from plastic changes and explores the link between selection and evolution. These questions are addressed using a combination of computer simulations and of data from the long-term individual-based monitoring of a snow vole population.

1.3.2 Snow voles in Churwalden

The snow vole (*Chionomys nivalis*, Martins 1842) is a medium-sized rodent, its adult body size ranges from 10 to 14 cm, without the tail (5 to 7.5 cm long). Contrary to a widespread intuition, snow voles are not white (Fig. 1.2). Instead, the fur colour of



the upper-parts varies from light to dark taupe grey, sometimes tinted with brown or dark red. The misconception about color highlights that the species could favourably be renamed *rock vole*: it is a rock, rather than a snow, specialist (Luque-larena, López, and Gosálbez 2002) and might be associated with high elevations only because rocky areas are more widespread there. It is sparsely distributed across southern Europe and Asia Minor, from sea level up to 4000 m of elevation (Janeau and Aulagnier 1997).



Figure 1.2: Juvenile (left picture) and adult (right picture) snow voles in their habitat in Churwalden, Switzerland. Juveniles always lack the brown hue generally found in adults. Neither adults nor juveniles are white.

Snow voles excavate burrows under the rocks, but can also use natural clefts between rocks, sometimes carrying small stones to build walls (Niederer 2008). A burrow consists of tunnels connecting chambers, one for the nest and multiple ones to stock dry plants (Janeau and Aulagnier 1997). The species is not known to hibernate and is therefore exposed to harsh winter conditions in its high-elevation range. Adult females actively defend small territories against non-relatives, and tend to form matrilineal clusters of territories, whereas adult males wander, and fight, across large overlapping home-ranges (Luque-larena, López, and Gosálbez 2004; García-Navas, Bonnet, Waldvogel, et al. 2016). The mating system is promiscuous and a same litter can be sired by multiple males. Females normally produce 1 to 4 litters of 1 to 5 pups between May and September. Juveniles generally do not reproduce in their first civil year. Although they can eat flour worms in the lab, there is no evidence that snow voles are not strictly herbivorous in the wild (Janeau and Aulagnier 1997). In the Swiss Alps, snow voles suffer predation from red foxes, stoats, various owls and corvids, and parasitism from fleas, lice and ticks (Janeau and Aulagnier 1997; Martinoli et al. 2001).

The study area is located by the Churer Joch, Churwalden, in the Swiss canton Graubünden (Fig. 1.3; coordinates 46°48' N, 9°34' E), and covers about 5 ha between 1980 m and 2100 m above sea level. It consists of a west-exposed scree interspersed with small coniferous trees and with patches of alpine meadow. The study area is demarcated by extensive meadows to the south and to the north, by a coniferous forest to the west and by cliff to the east (Fig. 1.4).

Another scree offers about 1 ha of favourable habitat, starting 300 m north-east to




Figure 1.3: Location of the study area in Switzerland. The yellow triangle indicates the study area, with coordinates $46^{\circ}48' N$, $9^{\circ}34' E$, by Churwalden, in the Swiss canton Graubünden. Countries are filled with different shades of grey, Austria and Lichtenstein are not labelled.

the monitored area. This area was trapped in 2008 and 2013. The snow vole density was rather low, with on average five captures per night of trapping, versus 18 on the main study area. More habitat favourable to snow voles can be found 2 Km to the south. The study population is moderately isolated and receives 5 to 10 immigrants per year, on a total of 60 to 180 individuals (García-Navas, Bonnet, Waldvogel, et al. 2016).

The monitoring of this snow vole population was initiated in 2006 by Dr. Peter W. Wandeler. Dr. Erik Postma took the monitoring over in 2012, but the protocol has remained practically unchanged. This thesis contains data collected up to the year 2015. Every year from 2006 to 2016, snow voles were life-trapped multiple times between late May and early October. Traps were set during the day, opened around sunset and checked the next morning. For every snow vole capture³, we recorded sex, age, body mass, body length, tail length, date, location and signs of reproductive activity (pregnancy, lactation, swollen scrotum). In addition, all newly-captured snow voles were individually marked and genotyped for 18 microsatellites (Wandeler, Ravaioli, and Bucher 2008). Based on the autosomal microsatellite genotypes, we reconstruct the pedigree of the population. This pedigree is the raw material for most of the work carried out during this thesis. In particular, the pedigree is used to define reproductive

³Other species (bank voles, pine voles, wood mice, stoats, black salamanders, slugs...) were released without taking measurements.



Figure 1.4: Distant view of the field site, taken from the west. The trapped area covers about a fifth of the width and a tenth of the height of the picture and is located in the centre. This scree is surrounded by a forest, a cliff and meadows.

success, as well as to estimate the relatedness between all pairs of individuals. These two statistics are essential to estimate selection, fitness and genetic variation.

1.3.3 Thesis outline

In natural populations, fitness is generally measured using individual measures of reproductive success and survival. These proxies are not fitness itself and their variation is largely stochastic, leading some authors to doubt that there is any significant variation in fitness in natural populations. Recent methodological developments appeared to support the view that variation in reproduction and survival was purely stochastic, and suggested that the potential for selection and evolution in the wild was largely over-estimated. In chapter ?? I examine these methods and, based on computer simulations, demonstrate that they lack statistical power to detect latent variation in fitness components. Using an alternative approach we show the presence of significant variation in the propensity of reproductive success in the snow vole population, thereby showing some potential for selection and adaptive evolution in this population. We also attempt to clarify some conceptual misunderstandings between the proponents of the two methodological schools.

A similar attempt motivated chapter ??, where, with collaborators from different methodological schools, we review and compare four frameworks to disentangle the causes of phenotypic changes. In particular, these frameworks differ in their estimation of the relative roles of plasticity, demography and genetic change. Based on com-




Figure 1.5: Orthophoto of the study site, from 2008. The red shading indicates the approximate area where traps are set.

puter simulations and on mathematical comparisons, we show that the discrepancies between the frameworks primarily originates from different definitions of the components of change. Nevertheless, one of these frameworks, the quantitative genetics *animal model*, stands out as the only framework able to estimate genetic change and the response to selection (that is, the trans-generational consequence of variation in fitness). I relied heavily on this framework for the two next chapters.

In chapter ??, I explore the reasons of the mismatch between apparent phenotypic selection, phenotypic change and genetic change for body mass. I describe one of the first case of contemporary evolution of a quantitative trait in the wild and show that this genetic change is adaptive. Both the evolution and the selective pressure responsible for it are invisible to purely phenotypic approaches, however. Using multivariate animal models, we identify the main component of selection as juvenile viability. I then infer that the target of selection is potential adult mass in juveniles and that selection is related to a recent change in climatic conditions.

The previous chapter considered selection and evolution averaged over the whole study period, without considering their temporal dynamic within the period. The fluctuation of selection is thought to be a major determinant of the rate of evolution, and a process to consider to understand adaptation in the wild. Nevertheless, unbiased measures of the variation of selection are rare and of the coupling between variation in selection and variation in evolution has been largely ignored. Chapter ?? shows that selection fluctuates in the study population, mainly due to variation in fertility selection. The rate of adaptive evolution is, however, remarkably constant, because viability selection, the driver of body mass evolution, does not vary. In this case the fluctuation of selection is evolutionary irrelevant. These two last chapters highlight the dangers of relying on phenotypic estimates of selection to understand the evolutionary dynamics of natural populations.

Finally, in chapter ??, I summarize the progresses made during this PhD on the

understanding of natural populations, and discuss some of the remaining challenges and future working directions.

References

- Akçay, E. and Van Cleve, J. 2016. "There is no fitness but fitness, and the lineage is its bearer." *Philosophical Transactions of the Royal Society B: Biological Sciences* **371** (1687): 20150085.
- Altermatt, F., Ebert, D., and Altermatt, F. 2016. "Reduced flight-to-light behaviour of moth populations exposed to long-term urban light pollution." *Biology Letters* **12**:20160111.
- Arnold, S. J. and Wade, M. J. 1984. "On the measurement of natural and sexual selection: Theory." *Evolution* **38** (4): 709–719.
- Bloom, J. S., Ehrenreich, I. M., Loo, W. T., Lite, T.-L. V., and Kruglyak, L. 2013. "Finding the sources of missing heritability in a yeast cross." *Nature* **494**:234–239.
- Brandon, R. and Beatty, J. 1984. "The propensity interpretation of 'fitness'. No interpretation is no substitute." *Philosophy of Science* **51** (2): 342–347.
- Brookfield, J. F. 2016. "Why are estimates of the strength and direction of natural selection from wild populations not congruent with observed rates of phenotypic change?" *BioEssays* **38**:1–8.
- Cam, E. and Monnat, J. Y. 2000. "Stratification based on reproductive state reveals contrasting patterns of age-related variation in demographic parameters in the kittiwake." *Oikos* **90** (3): 560–574.
- Darwin, C. 1859. *On the origin of species by means of natural selection, or the preservation of favoured races in the struggle for life*. John Murray, London.
- Dietrich, M. R. 2006. "From Mendel to molecules: A brief history of evolutionary genetics." In *Evolutionary genetics: concepts and case studies*. Edited by C. W. Fox and J. B. Wolf, 3–13. Oxford University Press, Oxford.
- Fisher, R. A. 1941. "Average excess and average effect of a gene substitution." *Annals of Eugenics* **11** (1): 53–63.
- Fisher, R. 1930. *The genetical theory of natural selection*. First. Oxford university press, Oxford, Great Britain.
- Fitzpatrick, M. J., Ben-shahar, Y., Smid, H. M., Vet, L. E. M., Robinson, G. E., and Sokolowski, M. B. 2005. "Candidate genes for behavioural ecology." *Trends in Ecology & Evolution* **20** (2): 96–104.
- Flint, J. and Mackay, T. F. C. 2009. "Genetic architecture of quantitative traits in flies, mice and humans." *Genome Research* **19**:723–733.

- Frank, S. a. 2012. "Natural selection. V. How to read the fundamental equations of evolutionary change in terms of information theory." *Journal of Evolutionary Biology* **25** (12): 2377–2396.
- García-Navas, V., Bonnet, T., Bonal, R., and Postma, E. 2016. "The role of fecundity and sexual selection in the evolution of size and sexual size dimorphism in New World and Old World voles (Rodentia: Arvicolinae)." *Oikos* **early view**.
- García-Navas, V., Bonnet, T., Waldvogel, D., Camenisch, G., and Postma, E. 2016. "Consequences of natal philopatry for reproductive success and mate choice in an Alpine rodent." *Behavioral Ecology* **27** (4): 1158–1166.
- Goodwin, S., McPherson, J. D., and McCombie, W. R. 2016. "Coming of age: ten years of next-generation sequencing technologies." *Nature Reviews Genetics* **17** (6): 333–351.
- Henderson, C. 1950. "Estimation of genetic parameters." *Annals of Mathematical Statistics* **21** (2): 309–310.
- Houseknecht, K. L., Baile, C. A., Matteri, R. L., and Spurlock, M. E. 1998. "The biology of leptin : a review." *Journal of Animal Science* **76**:1405–1420.
- Huang, W. and Mackay, T. F. 2016. "The genetic architecture of quantitative traits cannot be inferred from variance component analysis." *bioRxiv*. eprint: <http://biorxiv.org/content/early/2016/02/26/041434.full.pdf>.
- Janeau, G. and Aulagnier, S. 1997. "Snow vole - *Chionomys nivalis* (Martins 1842)." *IBEX Journal of Mountain Ecology* **4**:1–11.
- Jensen, H., Szulkin, M., and Slate, J. 2014. "Molecular quantitative genetics." In *Quantitative Genetics in the Wild*, edited by A. Charmentier, D. Garant, and L. E. B. Kruuk. Oxford: Oxford University Press.
- Jones, J. S. 1987. "The heritability of fitness: bad news for good genes?" *Trends in Ecology & Evolution* **2** (2): 35–36.
- Jong, M. a. de, Wong, S. C., Lehtonen, R., and Hanski, I. 2014. "Cytochrome P450 gene Cyp337 and heritability of fitness traits in the Glanville fritillary butterfly." *Molecular Ecology* **23** (8): 1994–2005.
- Kirschner, M. W. and Gerhart, J. C. 2010. "Facilitated variation." In *Evolution. The extended synthesis*, edited by M. Pigliucci and G. B. Müller. The MIT press, Cambridge, Massachusetts, USA.
- Krimbas, C. B. 2004. "On fitness." *Biology & Philosophy* **19** (2): 185–203.
- Kruuk, L. E. B. 2004. "Estimating genetic parameters in natural populations using the 'animal model'." *Philosophical Transactions of the Royal Society B: Biological Sciences* **359** (1446): 873–890.
- Kruuk, L. E. B., Charmantier, A., and Garant, D. 2014. "The study of quantitative genetics in wild populations." In *Quantitative Genetics in the Wild*, edited by A. Charmentier, D. Garant, and L. E. B. Kruuk. Oxford: Oxford University Press.

Chapter 1 General introduction

- Lande, R. and Arnold, S. J. 1983. "The measurement of selection on correlated characters." *Evolution* **37** (6): 1210–1226.
- Lande, R. 1979. "Quantitative genetic analysis of multivariate evolution, applied to brain: body size allometry." *Evolution*: 402–416.
- Lango Allen, H., Estrada, K., Lettre, G., Berndt, S. I., Weedon, M. N., Rivadeneira, F., Willer, C. J., et al. 2010. "Hundreds of variants clustered in genomic loci and biological pathways affect human height." *Nature* **467** (7317): 832–8.
- Liedvogel, M., Cornwallis, C. K., and Sheldon, B. C. 2012. "Integrating candidate gene and quantitative genetic approaches to understand variation in timing of breeding in wild tit populations." *Journal of Evolutionary Biology* **25** (5): 813–23.
- Luque-larena, J. J., López, P., and Gosálbez, J. 2002. "Microhabitat use by the snow vole *Chionomys nivalis* in alpine environments reflects rock-dwelling preferences." *Canadian Journal of Zoology* **80**:36–41.
- . 2004. "Spacing behavior and morphology predict promiscuous mating strategies in the rock- dwelling snow vole, *Chionomys nivalis*." *Canadian Journal of Zoology* **82**:1051–1060.
- Lush, J. 1937. *Animal breeding plans*. Ames, Iowa: Iowa State College Press.
- Lynch, M. and Walsh, B. 1998. *Genetics and Analysis of Quantitative Traits*. Sinauer Associates.
- Mackay, T. 2001. "The genetic architecture of quantitative traits." *Annual Review of Genetics* **35**:303–309.
- Martinoli, A., Preatoni, D. G., Chiarenzi, B., Wauters, L. a., and Tosi, G. 2001. "Diet of stoats (*Mustela erminea*) in an Alpine habitat:The importance of fruit consumption in summer." *Acta Oecologica* **22** (1): 45–53.
- Mc Carty, J. P. 2001. "Ecological Consequences of Recent Climate Change." *Conservation Biology* **15** (2): 320–331.
- Merilä, J., Sheldon, B. C., and Kruuk, L. E. B. 2001. "Explaining stasis: microevolutionary studies in natural populations." *Genetica* **112**:199–222.
- Niederer, A. 2008. "Das Verhalten der Schneemaus." PhD diss., Universität Basel.
- Nietlisbach, P. and Hadfield, J. D. 2015. "Heritability of heterozygosity offers a new way of understanding why dominant gene action contributes to additive genetic variance." *Evolution*: 1–16.
- Oswald, T., MacLeod, C., and McCarty, M. 1944. "Studies on the chemical nature of the substance inducing transformation of pneumococcal types: induction of transformation by a desoxyribonucleic acid fraction isolated from pneumococcus type III." *Journal of Experimental Medicine* **79** (2).
- Parmesan, C. 2006. "Ecological and evolutionary responses to recent climate change." *Annual Review of Ecology, Evolution, and Systematics*: 637–669.

- Pigliucci, M. and Müller, G. B. 2010. *Evoution. The extended synthesis*. The MIT press, Cambridge, Massachusetts, USA.
- Postma, E. 2014. "Four decades of estimating heritabilities in wild vertebrate populations: improved methods, more data better estimates?" In *Quantitative Genetics in the Wild*, edited by A. Charmentier, D. Garant, and L. E. B. Kruuk. Oxford: Oxford University Press.
- Price, P. W. 1996. *Biological evolution*. Saunders College Publishing, Philadelphia, PA.
- Rousset, F. 2004. *Genetic structure and selection in subdivided populations*. Princeton, New Jersey, USA.
- Shaw, R. G. and Shaw, F. H. 2014. "Quantitative genetic study of the adaptive process." *Heredity* **112** (1): 13–20.
- Steiner, U. K. and Tuljapurkar, S. 2012. "Neutral theory for life histories and individual variability in fitness components." *Proceedings of the National Academy of Sciences, USA* **109** (12): 4684–4689.
- Tafani, M., Cohas, A., Bonenfant, C., Gaillard, J.-M., and Allainé, D. 2013. "Decreasing litter size of marmots over time: a life history response to climate change ?" *Ecology* **94** (3): 580–586.
- Tienderen, P. H. van. 2000. "Elasticities and the link between demographic and evolutionary dynamics." *Ecology* **81** (3): 666–679.
- Turner, B. M. 2009. "Epigenetic responses to environmental change and their evolutionary implications." *Philosophical Transactions of the Royal Society B: Biological Sciences* **364** (1534): 3403–18.
- Wade, M. J. 2006. "Natural selection." In *Evolutionary genetics: concepts and case studies*. Edited by C. W. Fox and J. B. Wolf, 399–413. Oxford University Press, Oxford.
- Wagner, G. P. 2010. "The measurement theory of fitness." *Evolution* **64** (5): 1358–76.
- Wandeler, P., Ravaioli, R., and Bucher, T. B. 2008. "Microsatellite DNA markers for the snow vole (*Chionomys nivalis*)." *Molecular Ecology Resources* **8**:637–639.
- Watson, J. and Crick, F. 1953. "Genetical implications of the structure of deoxyribonucleic acid." *Nature* **171** (4361): 964–967.
- Wayne, M. L. and Miyamoto, M. M. 2006. "Genetic variation." In *Evolutionary genetics: concepts and case studies*. Edited by C. W. Fox and J. B. Wolf, 399–413. Oxford University Press, Oxford.
- Wilkins, J. S. 2009. *Species. A history of the idea*. Berkeley, California, USA: University of California Press.
- Wolf, J. B. and Wade, M. J. 2009. "What are maternal effects (and what are they not)?" *Philosophical Transactions of the Royal Society B: Biological Sciences* **364** (1520): 1107–15.

Chapter 1 General introduction

Yang, J., Benyamin, B., McEvoy, B. P., Gordon, S., Henders, A. K., Nyholt, D. R., Madden, P. A., et al. 2010. "Common SNPs explain a large proportion of the heritability for human height." *Nature Genetics* **42** (7): 565–569.

Addresses of co-authors

Koen J. van Benthem, Arpat Ozgul, Glauco Camenisch and Erik Postma

Department of Evolutionary Biology and Environmental Studies
University of Zurich-Irchel
Winterthurerstrasse 190
CH-8057 Zurich
Switzerland

Marjolein Bruijning and Eelke Jongejans

Animal Ecology and Physiology Department
Radboud University
Heyendaalseweg 135
6525 AJ Nijmegen
The Netherlands

Peter Wandeler

Musée d'histoire naturelle de Fribourg
Chemin du Musée 6
CH-1700 Freiburg
Switzerland