

Exercise 1 Use the Fundamental Theorem of Calculus, Part 1 to find the derivative of the function.

$$(a) \quad g(x) = \int_1^x \ln(1+t^2) dt$$

$$(b) \quad h(u) = \int_0^u \frac{\sqrt{t}}{t+1} dt$$

$$(c) \quad R(y) = \int_y^2 t^3 \sin(t) dt$$

$$(d) \quad h(x) = \int_1^{\sqrt{x}} \frac{z^2}{z^4+1} dz$$

$$(e) \quad y = \int_0^{x^4} \cos^2(\theta) d\theta$$

$$(f) \quad y = \int_{\sin(x)}^1 \sqrt{1+t^2} dt$$

Exercise 2 Evaluate the integral.

$$(a) \quad \int_0^1 (1 - 8v^3 + 16v^7) dv$$

$$(b) \quad \int_{-5}^5 e dx$$

$$(c) \quad \int_{-1}^2 (3u-2)(u+1) du$$

$$(d) \quad \int_{\pi/4}^{\pi/3} \csc^2(\theta) d\theta$$

$$(e) \quad \int_0^1 \cosh(t) dt$$

$$(f) \quad \int_1^3 \frac{y^3 - 2y^2 - y}{y^2} dy$$

Exercise 3 Find the derivative of the function.

$$g(x) = \int_{1-2x}^{1+2x} t \sin(t) dt$$

Exercise 4 If $f(x) = \int_0^x (1 - t^2)e^{t^2} dt$, on what interval is f increasing?

Exercise 5 The **error function** is defined as follows:

$$\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$$

Show that the function $y = e^{x^2} \operatorname{erf}(x)$ satisfies the differential equation $y' = 2xy + \frac{2}{\sqrt{\pi}}$.