

## Lecture 8: Lagrange Interpolation

Solving via linear system.

$$\begin{bmatrix} 1 & x_0 & x_0^2 & \cdots & x_0^n \\ 1 & x_1 & x_1^2 & \cdots & x_1^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_n & x_n^2 & \cdots & x_n^n \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} f(x_0) \\ f(x_1) \\ \vdots \\ f(x_n) \end{bmatrix}$$

Solving via basis polynomials. Let

$$L_k(x) = \frac{(x-x_0)\cdots(x-x_{k-1})(x-x_{k+1})\cdots(x-x_n)}{(x_k-x_0)\cdots(x_k-x_{k-1})(x_k-x_{k+1})\cdots(x_k-x_n)} \text{ for } k=0,1,\dots,n. \text{ Then } P_n(x) = y_0 L_0(x) + y_1 L_1(x) + \cdots + y_n L_n(x).$$

Error analysis.

$\forall x \in [a, b], \exists \xi \in (\min\{x, x_0, x_1, \dots, x_n\}, \max\{x, x_0, x_1, \dots, x_n\})$  such that

$$f(x) = P_n(x) + \frac{f^{(n+1)}(\xi)}{(n+1)!} (x-x_0)(x-x_1)\cdots(x-x_n)$$

## Tutorial 3: Lagrange Interpolation

**Example 1.** Construct the LIP for  $f(x) = \log_2 x$  using  $x_0 = 1/2, x_1 = 1, x_2 = 2, x_3 = 4$ . Find a bound of the absolute error for any  $x \in [1/2, +\infty)$ .

**Solution.**  $f(0.5) = -1, f(1) = 0, f(2) = 1, f(4) = 2$ . Solve using linear system to get  $a_0 = -\frac{52}{21}, a_1 = \frac{7}{2}, a_2 = -\frac{7}{6}, a_3 = \frac{1}{7}$ .

So  $P(x) = \frac{1}{7}x^3 - \frac{7}{6}x^2 + \frac{7}{2}x - \frac{52}{21}$ . To get an u.b for error, first find  $f^{(4)}(x) = -\frac{6}{x^4 \ln 2}$ . Note monotonicity. Hence

$$|f^{(4)}(x)| \leq \frac{6}{(1/2)^4 \ln 2} = \frac{96}{\ln 2}. \text{ Thus an u.b for absolute error}$$

$$|P(x) - f(x)| \text{ is } \frac{1}{4!} \times \frac{96}{\ln 2} |(x-1/2)(x-1)(x-2)(x-4)| = \frac{4}{\ln 2} |(x-1/2)(x-1)(x-2)(x-4)|. \square.$$

**Result from Example 2.** If nodes are equidistributed, the maximum value of  $g(x) = |(x-x_0)(x-x_1)\cdots(x-x_n)|$  must be attained in  $(x_0, x_1)$  and  $(x_{n-1}, x_n)$  (due to the symmetry).  $|g(x^*)| \leq \frac{1}{4} N! h^{N+1}$ .

**Error estimation for equidistributed nodes.**

$$|P_N(x) - f(x)| \leq \frac{h^{N+1}}{4(N+1)} \max_{\xi \in [a, b]} |f^{(N+1)}(\xi)|, \text{ for all } x \in [a, b]$$

**Exercise 2.** Let  $P_n(x)$  be the LIP for  $f(x) = \cos x$  with

$x_k = kh, k = 0, 1, \dots, n$  where  $h = \pi/(2n)$ . **1.** Find a positive integer  $N$  such that  $|P_N(x) - f(x)| < 0.005$ , for all  $x \in [0, \pi/2]$ .

**Solution.** For  $f(x) = \cos(x)$ ,  $\max_{\xi \in [0, \pi/2]} |f^{(N+1)}(\xi)| = 1$ . Hence it suffices to find  $\frac{h^{N+1}}{4(N+1)} < 0.005 \implies N \geq 3$ .

## Lecture 9: Divided Differences

How to find the Lagrange polynomial.

$$P_n(x) = a_0 + a_1(x-x_0) + a_2(x-x_0)(x-x_1) + \cdots + a_n(x-x_0)(x-x_1)\cdots(x-x_{n-1}) \text{ where } a_k = f[x_0, x_1, \dots, x_k]$$

$$\text{and } f[x_0, x_1, \dots, x_n] = \frac{f[x_1, x_2, \dots, x_n] - f[x_0, x_1, \dots, x_{n-1}]}{x_n - x_0}.$$

$$a_0 = f(x_0).$$

## Lecture 10: Cubic Spline Interpolation (CSI)

How to find  $\mu_k$  and  $\lambda_k$ .

$$\mu_k = \frac{x_k - x_{k-1}}{x_{k+1} - x_{k-1}}, \quad \lambda_k = \frac{x_{k+1} - x_k}{x_{k+1} - x_{k-1}}, \quad k = 1, 2, \dots, n-1$$

**Natural Boundary Conditions.**  $M_0 = M_n = 0$ .

$$\begin{bmatrix} 2 & \lambda_1 & & & \\ \mu_2 & 2 & \lambda_2 & & \\ & \mu_3 & 2 & \ddots & \\ & & \ddots & \ddots & \lambda_{n-2} \\ & & & \mu_{n-1} & 2 \end{bmatrix} \begin{bmatrix} M_1 \\ M_2 \\ M_3 \\ \vdots \\ M_{n-2} \\ M_{n-1} \end{bmatrix} = \begin{bmatrix} 6f[x_0, x_1, x_2] \\ 6f[x_1, x_2, x_3] \\ 6f[x_2, x_3, x_4] \\ \vdots \\ 6f[x_{n-3}, x_{n-2}, x_{n-1}] \\ 6f[x_{n-2}, x_{n-1}, x_n] \end{bmatrix}$$

**Clamped Boundary Conditions.**

$$2M_0 + M_1 = 6f[x_0, x_0, x_1], \quad M_{n-1} + 2M_n = 6f[x_{n-1}, x_n, x_n].$$

$$\begin{bmatrix} 2 & \lambda_0 & & & \\ \mu_1 & 2 & \lambda_1 & & \\ & \ddots & \ddots & \ddots & \\ & & \mu_{n-1} & 2 & \lambda_{n-1} \\ & & & \mu_n & 2 \end{bmatrix} \begin{bmatrix} M_0 \\ M_1 \\ \vdots \\ M_{n-1} \\ M_n \end{bmatrix} = \begin{bmatrix} 6f[x_0, x_0, x_1] \\ 6f[x_0, x_1, x_2] \\ \vdots \\ 6f[x_{n-2}, x_{n-1}, x_n] \\ 6f[x_{n-1}, x_n, x_n] \end{bmatrix}$$

How to find  $S_k$ .

$$S_k(x) = M_{k-1} \frac{(x-x_k)^3}{6(x_{k-1}-x_k)} + M_k \frac{(x-x_{k-1})^3}{6(x_k-x_{k-1})} + A_k x + B_k.$$

$$A_k = \frac{f(x_k) - f(x_{k-1})}{x_k - x_{k-1}} - \frac{1}{6} (M_k - M_{k-1}) (x_k - x_{k-1}).$$

$$B_k = \frac{x_k f(x_{k-1}) - x_{k-1} f(x_k)}{x_k - x_{k-1}} + \frac{1}{6} (M_k x_{k-1} - M_{k-1} x_k) (x_k - x_{k-1}).$$

**Piecewise Linear Interpolation (PLI).** If

$$S(x) = S_k(x), \quad \text{for } x \in [x_{k-1}, x_k], k = 1, 2, \dots, n \text{ then}$$

$$S_k(x) = f(x_{k-1}) \frac{x-x_k}{x_k-x_{k-1}} + f(x_k) \frac{x-x_{k-1}}{x_k-x_{k-1}}$$

**Error analysis for PLI on equidistributed nodes.** If

$$x_k = x_0 + kh, \text{ then for } x \in [x_0, x_n],$$

$$|f(x) - S(x)| \leq \frac{1}{8} h^2 \max_{\xi \in [x_0, x_n]} |f''(\xi)|$$

## Tutorial 4: Divided Diff and CSI

**Example 2: Quadratic spline interpolation.** Given  $n+1$  nodes  $x_0 < x_1 < \cdots < x_{n-1} < x_n$  and a continuous function  $f(x)$ , find a function  $S(x)$  such that **1.**  $S(x)$  is first-order differentiable on  $(x_0, x_n)$  **2.**  $S(x)$  is a quadratic polynomial on  $(x_{k-1}, x_k)$  for any  $k = 2, 3, \dots, n$ ; **3.**  $S(x)$  is a linear function on  $(x_0, x_1)$ . **4.**  $S(x_k) = f(x_k)$  for all  $k = 0, 1, \dots, n$ .

$$\text{Solution. } S_1(x) = f(x_0) \frac{x-x_1}{x_0-x_1} + f(x_1) \frac{x-x_0}{x_1-x_0}.$$

$$S_k(x) = \frac{1}{2} M_k \frac{(x-x_{k-1})^2}{x_k-x_{k-1}} + \frac{1}{2} M_{k-1} \frac{(x-x_k)^2}{x_{k-1}-x_k} + C_k, \quad k = 2, 3, \dots, n.$$

$$C_k = f(x_k) - \frac{1}{2} M_k (x_k - x_{k-1}). \quad M_k = 2f[x_{k-1}, x_k] - M_{k-1}.$$

Since  $M_1 = S'_1(x_1) = f[x_0, x_1]$ ,  $M_k$  for  $k = 2, 3, \dots, n$  can be obtained iteratively.

## Lecture 11: Least Squares Approximation

Finding the coefficients. Let

$$X = \begin{pmatrix} 1 & x_0 & x_0^2 & \cdots & x_0^n \\ 1 & x_1 & x_1^2 & \cdots & x_1^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_m & x_m^2 & \cdots & x_m^n \end{pmatrix} \text{ and}$$

$$\mathbf{a} = (a_0, a_1, \dots, a_n)^T, \quad \mathbf{y} = (y_0, y_1, \dots, y_m)^T. \text{ Solve } X^T X \mathbf{a} = X^T \mathbf{y}.$$

**Proof for Exercise 1.** **TODO**

**Weighted LSA.**  $W = \text{diag}\{w_0, w_1, \dots, w_m\}$ . Solve  $X^T W X \mathbf{a} = X^T W \mathbf{y}$ .

## Lecture 12: Newton-Cotes Formulae

Ascertaining number of points for error  $< \epsilon$ . **TODO**

$$\text{Trapezoidal Rule. } \int_a^b f(x) dx \approx \frac{b-a}{2} [f(a) + f(b)].$$

**Error for Trapezoidal Rule.**

$$\int_a^b f(x) dx = \frac{b-a}{2} [f(a) + f(b)] - \frac{1}{12} (b-a)^3 f''(\xi)$$

**Simpson's Rule.** Given  $f(a), f\left(\frac{a+b}{2}\right)$  and  $f(b)$ ,  $P(x) =$

$$f(a) + \frac{f(b)-f(a)}{b-a} (x-a) + \left[ 2f\left(\frac{a+b}{2}\right) - f(b) - f(a) \right] \frac{2(x-a)(x-b)}{(b-a)^2}$$

whose integral is  $\left[ \frac{2}{3} f\left(\frac{a+b}{2}\right) + \frac{1}{6} f(b) + \frac{1}{6} f(a) \right] (b-a)$ .

**Error for Simpson's Rule.**

$$\int_a^b f(x) dx = \frac{b-a}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{90} \left(\frac{b-a}{2}\right)^5 f^{(4)}(\xi)$$

**Theorem 3 on Page 4.** **TODO**

**Gaussian Elimination to find weights.** **TODO**

## Lecture 13: Composite Numerical Integration

**Composite Trapezoidal Rule (CTR).** **Confusing use of h.**

**Read carefully first**

**Error analysis for CTR.** **TODO**

**Composite Simpson's Rule (CSR).** **TODO**

**Error analysis for CSR.** **TODO**

## Tutorial 5: LSA and Integration

**TODO**

Miscellaneous

The Gamma function.  $\Gamma(x) = \int_0^{+\infty} t^{x-1}e^{-t}dt$  and  $\Gamma(n) = (n-1)!$  for any positive integer  $n$ .

Linear Algebra and Calculus. TODO

General formulae for higher derivatives. Higher order derivatives

Error proofs from lectures/tutorials. LSA proof of optimality, the proof of errors, tutorial identities

Finding the set of  $x$  and  $y$  values for the lowest error LIP.

First find the lowest-error polynomial (LEP), which is  $p(x) = a_0 + a_1x + a_2x^2 + \cdots + a_mx^m$ . From Homework Q3,

$$M = \begin{pmatrix} b-a & \frac{b^2-a^2}{2} & \cdots & \frac{b^{m+1}-a^{m+1}}{m+1} \\ \frac{b^2-a^2}{2} & \frac{b^3-a^3}{3} & \cdots & \frac{b^{m+2}-a^{m+2}}{m+2} \\ \vdots & \vdots & & \vdots \\ \frac{b^{m+1}-a^{m+1}}{m+1} & \frac{b^{m+2}-a^{m+2}}{m+2} & \cdots & \frac{b^{2m+1}-a^{2m+1}}{2m+1} \end{pmatrix}$$
and

$b = \left(\int_a^b x^0 f(x)dx, \cdots, \int_a^b x^m f(x)dx\right)^T$ . Solve  $Ma = b$  to get the coefficients of the LEP. Then reverse-engineer the Gaussian Elimination process for finding  $a$  when constructing the LIP to get  $x$  and  $y$  (possibly non-unique).