

Masters Theory I

Homework #2

Out: Thursday

Due / Quiz following Thursday

- C&B 1.47
- C&B 1.51
- C&B 1.52
- C&B 1.53
- C&B 1.54
- C&B 2.1
- C&B 2.2
- C&B 2.3

- (Deb Glueck's problem from BIOS 160, UNC-Chapel Hill, 1989):

For families with four children, it is known that the probability of a child being a boy is equal to π . Assuming that the sex of a child is independent of the sex distribution of the other children in the family, find the probabilities of the following sex distributions in a randomly selected family of four children.

1. All children are of the same sex.
2. The first $k \leq 4$ children are boys and the rest are girls.
3. The first $k \leq 4$ children are the same sex, and the rest are the opposite sex.
4. There are at least $k \leq 4$ boys in the family.
5. The first s children are boys, and there are k ($s \leq k \leq 4$) boys in the family.
6. The first s children are boys, and there are at least k ($s \leq k \leq 4$) boys in the family.

Review problems (won't be on quiz, but could be on exam)

- C&B 1.44
- C&B 1.55
- C&B 2.6

- (Deb Glueck's problem from BIOS 160, UNC-Chapel Hill, 1989):

In a cancer study, 10 mice of strain A and 15 of strain B were treated. Each mouse has probability θ of developing cancer, independently of other mice, and altogether 8 mice developed cancer. What is the probability that exactly 4 of the mice who developed cancer were of strain B ?