KOSTAT-UNFPA Summer Seminar on Population

Workshop 1: Demographic Analysis: Methods and Tools in R

Mortality and Fertility

2 July, 2024

Instructor: Tim Riffe

Assistant: Inchan Hwang









Workshop plan, July 1-5, 2024

1: Monday Intro concepts, and R setup

2: Tuesday Mortality and fertility


3: Wednesday Structure

4: Thursday Growth


5: Friday Projection


Review of session 1


- age, period, cohort
- rates, probabilities, counts, ratios, stocks, flows
- Crude rate versus age-specific rate
- R projects, R markdown
- Objects, functions, grammar




$$M(x) = D(x) / E(x)$$









It makes a big difference for data in wider age groups!




The survivorship curve is inferred by chaining survival probabilities together over successive ages. This is the main trick of the period lifetable and why we call it synthetic. Further quantities are derived from this.




The survivorship curve is inferred by chaining survival probabilities together over successive ages. This is the main trick of the period lifetable and why we call it synthetic. Further quantities are derived from this, such as the lifetable death distribution. These two functions go hand-in-hand.



The survivorship curve is inferred by chaining survival probabilities together over successive ages. This is the main trick of the period lifetable and why we call it synthetic. Further quantities are derived from this, such as the lifetable death distribution. These two functions go hand-in-hand. You might even try drawing them together.




You might wonder how lifetable deaths relate to observed deaths (here rescaled to sum to 100000). They are quite different.



You might wonder how lifetable deaths relate to observed deaths (here rescaled to sum to 100000). They are quite different.






You might wonder how lifetable deaths relate to observed deaths (here rescaled to sum to 100000). They are quite different.


Life expectancy (e0) can be thought of as the **average age at death**, where **lifetable deaths are the weights**.

• equivalent to saying e0 is equal to the **area under the survival curve**, lx. In practice we do this, but first understand the dx-weighted age at death.

All other details are one of three kinds:

- 1. Adjustments to approximate the area under lx given discrete data.
- Adjustments to calculate remaining life expectancy at each age rather than just for age 0.
- 3. Adjustments for what values to use for the highest age (closeout)







Fertility analysis starts with the registration of births, Here birth records have been tabulated to birth counts by age of mother.

$$F(x) = B(x) / E(x)$$






Birth rates are often denoted Fx or ASFR (age-specific fertility rates). Unless otherwise stated, age means age of mother.



Birth rates are often denoted Fx or ASFR (age-specific fertility rates). Unless otherwise stated, age means age of mother. The total fertility rate TFR is the sum of these, in this case 1.2



Birth rates are often denoted Fx or ASFR (age-specific fertility rates). Unless otherwise stated, age means age of mother. The total fertility rate TFR is the sum of these, in this case 1.2. The mean age of childbearning (MAC or MAB) sometimes uses these rates as weights. Here the MAB matches the mode, 31.2.

Time for us to move to R!