

MAX-PLANCK-INSTITUT
FÜR DEMOGRAFISCHE
FORSCHUNG

MAX PLANCK INSTITUTE
FOR DEMOGRAPHIC
RESEARCH





MAX-PLANCK-INSTITUT
FÜR DEMOGRAFISCHE
FORSCHUNG

MAX PLANCK INSTITUTE
FOR DEMOGRAPHIC
RESEARCH


Morbidity concentration and dispersion

Tim Riffe, Aïda Solé Auro, Maarten J. Bijlsma



Fries' diagrams are a nice prop

Figure 1. Possible scenarios for future morbidity and longevity.





Pattern indifference within lifespan





Pattern indifference within lifespan






Patterns matter if lifespans are mixed






Patterns matter if lifespans are mixed






Patterns matter if lifespans are mixed






Patterns matter if lifespans are mixed






Patterns matter if lifespans are mixed






Patterns matter if lifespans are mixed






Patterns matter if lifespans are mixed





Patterns matter if lifespans are mixed





Compression

Compression definition

The level of morbidity compression is the average proportion of life in good health, $\mathbb{C} = \frac{HLE}{LE}$.

Objective:

Separate morbidity levels and morbidity dispersion.



Compression

Compression definition

The level of morbidity compression is the average proportion of life in good health, $\mathbb{C} = \frac{HLE}{LE}$.

Objective:

Separate morbidity levels and morbidity dispersion.



Dispersion

Dispersion definition

Morbidity dispersion, \mathbb{D} , is the average time-to-death of late-life morbidity prevalence.

Formal definition

$$\mathbb{D} = \frac{\int_0^{\omega} y \pi^*(y) \, dy}{\int_0^{\omega} \pi^*(y) \, dy} \quad (1)$$

where a is age, y is time until death, and $\pi^*(y)$ is morbidity prevalence by time to death.

Or one might rather weight a lifespan-varying $\pi(y, l)$, by the length-of-life distribution.



Dispersion

Dispersion definition

Morbidity dispersion, \mathbb{D} , is the average time-to-death of late-life morbidity prevalence.

Formal definition

$$\mathbb{D} = \frac{\int_0^{\omega} y \pi^*(y) \, dy}{\int_0^{\omega} \pi^*(y) \, dy} \quad (1)$$

where a is age, y is time until death, and $\pi^*(y)$ is morbidity prevalence by time to death.

Or one might rather weight a lifespan-varying $\pi(y, l)$, by the length-of-life distribution.



Dispersion

Dispersion definition

Morbidity dispersion, \mathbb{D} , is the average time-to-death of late-life morbidity prevalence.

Formal definition

$$\mathbb{D} = \frac{\int_0^{\omega} y \pi^*(y) \, dy}{\int_0^{\omega} \pi^*(y) \, dy} \quad (1)$$

where a is age, y is time until death, and $\pi^*(y)$ is morbidity prevalence by time to death.

Or one might rather weight a lifespan-varying $\pi(y, l)$, by the length-of-life distribution.



Scenarios

Scenario

ULE

HLE

LE

℄



℄

Base







Scenarios

Scenario	ULE	HLE	LE	\mathbb{C}	\mathbb{D}
Base					
					
	=	=	=	=	↓





Scenarios

Scenario		ULE	HLE	LE	\mathbb{C}	\mathbb{D}
Base						
		=	=	=	=	↑





Scenarios

Scenario		ULE	HLE	LE	\mathbb{C}	\mathbb{D}
Base						
		↓	↑	=	↑	=





Scenarios

Scenario		ULE	HLE	LE	\mathbb{C}	\mathbb{D}
Base						
		↓	↑	=	↑	↓





Scenarios

Scenario		ULE	HLE	LE	\mathbb{C}	\mathbb{D}
Base						
		\uparrow	\downarrow	$=$	\downarrow	$=$





Scenarios

Scenario	ULE	HLE	LE	\mathbb{C}	\mathbb{D}
Base					
	$=$	\uparrow	\uparrow	\uparrow	$=$
					





Scenarios

Scenario	ULE	HLE	LE	\mathbb{C}	\mathbb{D}
Base					
	\uparrow	\uparrow	\uparrow	\downarrow	\downarrow
					




Scenarios

Scenario	ULE	HLE	LE	\mathbb{C}	\mathbb{D}	
Base						
		$=$	\uparrow	\uparrow	\uparrow	\downarrow



Observed time-to-death patterns (HRS, ADL3)


1915





Observed time-to-death patterns (HRS, ADL3)


1916





Observed time-to-death patterns (HRS, ADL3)


1917





Observed time-to-death patterns (HRS, ADL3)


1918





Observed time-to-death patterns (HRS, ADL3)


1919





Observed time-to-death patterns (HRS, ADL3)


1920





Observed time-to-death patterns (HRS, ADL3)


1921





Observed time-to-death patterns (HRS, ADL3)


1922





Observed time-to-death patterns (HRS, ADL3)


1923





Observed time-to-death patterns (HRS, ADL3)


1924







Observed time-to-death patterns (HRS, ADL3)

1925







Results from HRS (RAND, vP), 82 measures







Results from HRS (RAND, vP), 82 measures





Results from HRS (RAND, vP), 82 measures





Summary

Measures

Dispersion measures an aspect of the shape of morbidity prevalence, and it is a complementary index to compression.

Trends

Dispersion of most morbidities changed less than 10% between the 1905 and 1925 cohorts, mostly in the direction of increased concentration of poor health conditions at the end of life.

Limitations

This measure requires follow-up until death (but stable patterns might make prevalence less risky to extrapolate).



Summary

Measures

Dispersion measures an aspect of the shape of morbidity prevalence, and it is a complementary index to compression.

Trends

Dispersion of most morbidities changed less than 10% between the 1905 and 1925 cohorts, mostly in the direction of increased concentration of poor health conditions at the end of life.

Limitations

This measure requires follow-up until death (but stable patterns might make prevalence less risky to extrapolate).



Summary

Measures

Dispersion measures an aspect of the shape of morbidity prevalence, and it is a complementary index to compression.

Trends

Dispersion of most morbidities changed less than 10% between the 1905 and 1925 cohorts, mostly in the direction of increased concentration of poor health conditions at the end of life.

Limitations

This measure requires follow-up until death (but stable patterns might make prevalence less risky to extrapolate).