


# Life lost, lifesaving, and causes of death

Tim Riffe<sup>1</sup> & Aïda Solé Auró<sup>2</sup>

<sup>1</sup> Department of Demography,  
University of California, Berkeley

<sup>2</sup> Universitat Pompeu Fabra

PAA 2015 Annual Meeting  
Session 234, Saturday, May 2



# Mortality measurement

Years Lost

Riffe & Solé

**lifetable:** purged of structure

**counts:** structure  $\times$  intensity

**combo:** Person Years of Life Lost (PYLL), \_\_\_\_\_, ...

# PYLL

Years Lost

Riffe & Solé

Person years of life lost

$$\text{PYLL} = \int_0^{\omega} D(x) \times e(x) \, dx \quad (1)$$

$$\text{PYLL}^c = \int_0^{\omega} D^c(x) \times e(x) \, dx \quad (2)$$

(mention that there are decreasing returns to  $\mu(x)$  improvements, since  $D(x)$  decreases, but  $e(x)$  increases = ambiguous change in PYLL?)

# PYLL

Years Lost

Riffe & Solé

PYLL has an age pattern that we're aware of but don't often look at:

1)  $D_x$  2)  $W_x$  in this case person years of life lost is assigned to the age in which death occurred, but it makes more sense to plot them through the ages in which the years will be lived.

Here a plot of the years of life lived through (foregone). In order to do this properly, you can't just do  $x + e(x)$ , but you need to use the whole conditional dist. Show eqn?

Years Lost

Riffe & Solé

Now the same thing for causes: no formula necessary, just figs.

Years Lost

Riffe & Solé

And now jump to causes.

# Life lost, lifesaving, and causes of death

Tim Riffe<sup>1</sup> & Aïda Solé Auró<sup>2</sup>

Thank you!

[tim.riffe@gmail.com](mailto:tim.riffe@gmail.com)

Aida's email here

