

IB Physics Topic E2 Quantum Physics; HL

By timthede07, M25 Cohort

Table of Contents

1	The Ultraviolet Catastrophe	1
2	The Photoelectric Effect	2
2.1	The Photoelectric Equation	4
2.2	Photon Momentum	4
3	Millikan's Photoelectric Experiment	5
4	The Compton Effect	8
4.1	Experimental Setup	8
4.2	Failure of EM Wave Theory	9
4.3	Explanation by Quantum Theory	9
4.4	Quantifying the Model	9
4.5	Explaining the Peaks	10
5	Alternative Photon-Interaction	11
5.1	Matter Waves and the de Broglie Hypothesis	12
5.2	The Davisson-Germer Experiment	12
6	Exam Questions	14
6.1	Intensity and Frequency	14
6.2	Evidence for Wave/Particle	15
6.3	Photoelectric Experiment — IV Graphs	16
6.4	Energy Transfer — Electron and Photon	18
6.5	De Broglie Wavelength and Momentum	19
6.6	Compton Scattering	20
6.7	Photoelectric Effect	22
6.8	Millikan's Experiment	24

1 The Ultraviolet Catastrophe

Rayleigh-Jeans Law, following from the principles of classical physics, stated that intensity of radiation emitted by a black body satisfies the following relationship

$$\text{intensity} \propto \text{frequency}^2$$

This model actually broke down at higher frequencies (lower wavelengths).

Max Planck proposed that the energy of the oscillators in the black body was quantized in integer values of hf , where f is the frequency of the oscillator and h is Planck's constant. His new model fixed the problem of the ultraviolet catastrophe, but he was not able to explain why the quantization was necessary.

2 The Photoelectric Effect

The *photoelectric effect* is a phenomenon where electrons are ejected from the surface of a material (usually a metal) when it is exposed to light of a certain frequency or higher. It is an evidence that light are particles (photons).

The classical predictions have the following implications

- Light was only understood as a wave. The classical idea was that with enough intensity, waves of any frequency should be able to transfer the binding energy to electrons and free them.
- The time delay when lower-frequency waves are used occurs because the electrons need to accumulate sufficient energy over time to reach the binding energy to be freed.

The observations made in reality were

- Threshold frequency: There is a minimum frequency of light below which no electrons are emitted.
- Intensity independence: The intensity of light does not affect the kinetic energy of the emitted electrons. If the light is below the threshold frequency, no electrons would be emitted no matter how intense the light is. However, the intensity is proportional to the number of electrons emitted per second.
- Instantaneous emission of electrons: There was no time lag between the light being shone and the electrons being emitted.
- Frequency dependence: The KE of the ejected electrons increases with the frequency of the light.

Einstein adopted Planck's idea of quantization and proposed the following

- EM radiation consists of **photons**. Each photon has an energy equal to $E = hf$.

- Each photon interacts with only a single electron. The photon disappears as its energy is entirely transferred to the electron.
- There exists a threshold frequency f_0 , which corresponds to the minimum energy required to free an electron from the material. Below this frequency, no electrons are emitted.
- The minimum energy required to free an electron is the **work function** $\phi = hf_0$ of the material. This is the energy required to overcome the forces holding the electron back onto the surface of the material.
- When the photon supplies an excess of energy above the work function to the electron, the excess energy is transferred to the electron as KE.
- Increasing the intensity **does not increase the KE** of individual electrons, but it increases the number of photons incident on the surface of the material per second, and thus the number of electrons ejected.
- The ejected electrons will eventually be attracted back to the surface of the plate that was left positive, even with high energies supplied.
- The energy $E = hf$ in the photon hitting the electron must be **strictly greater** than the work function for the electron to be ejected; i.e. $hf > \phi$.




Figure 1: The work function ϕ of a material

2.1 The Photoelectric Equation

The maximum KE of the ejected electrons is given by

$$E_{\max} = hf - \phi \quad (1)$$

2.2 Photon Momentum

Any object moving at the speed of light must have 0 mass. But this does not mean that one can use $p = mv$ to simply calculate the momentum of a photon as 0. Einstein has a relativistic equation for momentum

For a moving particle at a relativistic speed, the total energy is given by

$$E = \sqrt{p^2c^2 + m_0^2c^4}$$

where

- p is the momentum of the particle
- m_0 is the rest mass of the particle

A photon has rest mass 0, so the equation simplifies to $E = pc$. Then, as per Einstein, the energy of a photon is given by $E = \frac{hc}{\lambda}$, we can make a substitution and obtain

$$p = \frac{E}{c} = \frac{h}{\lambda} \quad (2)$$

3 Millikan's Photoelectric Experiment

This experiment is to test Einstein's photoelectric equation. The setup is as follows:

- **Vacuum Chamber:** The experiment was conducted in a vacuum to prevent collisions between the ejected electrons and air molecules, which could affect their motion and the experimental results.
- **Metal Surface (Photocathode):** A clean metal surface, such as sodium or potassium, was used as the target for the incident light. Different metals have different *work functions* (the minimum energy required to eject an electron), so the experiment needed to account for the material's response to varying light frequencies.
- **Light Source:** A monochromatic light source (emitting light of a single frequency) was directed at the metal surface. The frequency of the light was varied, while the intensity was kept constant in specific trials of the experiment.
- **Collector Electrode (Anode):** A positively charged electrode was placed opposite the metal surface. Electrons ejected from the photocathode traveled through the vacuum to the anode, provided they had enough kinetic energy to overcome any applied opposing voltage.
- **Adjustable Stopping Potential (Voltage):** A voltage source was used to apply a reverse potential to the anode. This opposing voltage could stop the electrons from reaching the anode, thereby allowing Millikan to measure the maximum kinetic energy of the ejected electrons.
- **Ammeter:** The current generated by the ejected electrons, as they reached the anode and completed the circuit, was measured using an ammeter.

Millikan's goal was to study the relationship between the frequency of the incident light and the kinetic energy of the emitted electrons. The experiment proceeded in the following steps:

1. **Illumination of the metal surface:** Monochromatic light was shone onto the metal surface, causing electrons to be ejected due to the photoelectric effect.
2. **Electron ejection and current measurement:** The ejected electrons traveled across the vacuum toward the positively charged anode. The movement of the electrons created a measurable current, as the circuit is now complete.
3. **Application of stopping potential:** Millikan varied the stopping potential (a negative voltage) applied to the anode. This potential created an opposing electric field that slowed the ejected electrons. When the stopping potential was sufficiently high, it completely stopped the electrons from reaching the anode.
4. **Measuring the stopping potential:** The stopping potential was a measure of the *maximum kinetic energy* of the ejected electrons. Since the kinetic energy of an electron is given by

$$E_{\max} = eV_s$$

where e is the electron charge and V_s is the stopping potential, Millikan could determine the kinetic energy of the ejected electrons by **measuring the voltage at which the current stopped**.

The expected results per Einstein's photoelectric equation are as follows (this is **verified by the experiment**):

- We combine Equation (1) and $E_{\max} = eV_s$ to obtain

$$eV_s = hf - hf_0$$

- Since photons travel at the speed of light, $f = \frac{c}{\lambda}$ substitution gives

$$eV_s = \frac{hc}{\lambda} - \frac{hc}{\lambda_0}$$

- Rearrangement gives

$$V_s = \frac{hc}{e} \left(\frac{1}{\lambda} - \frac{1}{\lambda_0} \right)$$

- This is a linear relationship with

1. V_s on the y-axis
2. $\frac{1}{\lambda}$ on the x-axis
3. $\frac{hc}{e}$ as the gradient
4. $-\frac{hc}{e\lambda_0} = -\frac{\phi}{e}$ as the y-intercept (the work function divided by the elementary charge)
5. $\frac{1}{\lambda_0}$ as the x-intercept

- This results shows that

1. The gradient is a constant, and is equal to $\frac{hc}{e}$. It does not depend on anything.
2. Different metal surfaces just give parallel lines with different axes intercepts.

- We may also rewrite it in terms of frequencies

$$V_s = \frac{h}{e} (f - f_0)$$

- Also note that the ideas also apply to graphs of E_k against f , since $E_k = eV_s$.

4 The Compton Effect

4.1 Experimental Setup

An incident beam of X-ray is incident on a graphite target. For various angles, he measured the wavelength of the scattered X-ray. The results are as follows.




Figure 2: The Compton Effect

For $\theta \neq 0$, the graph of λ against θ is a straight line has two peaks. One of which is simply the wavelength of the incident X-ray, and the other is a scaled-up version of the original wavelength.

4.2 Failure of EM Wave Theory

Following from the classical theory of EM waves, the X-ray causes the electrons in the graphite to oscillate at the same frequency as the X-ray, and thereby emit radiation at the same wavelength as the incident X-rays. This would force

$$\text{Incoming X-ray wavelength} = \text{Outgoing wavelength}$$

But this contradicts the experimental observations!

4.3 Explanation by Quantum Theory

Building upon Einstein's theory that light can be quantized into photons, Compton proposed that the X-ray photons collide with the electrons in the graphite on a one-to-one basis.

The collision causes the X-ray photon to lose energy and momentum, and the electron to gain energy and momentum. The scattered X-ray photon has a longer wavelength than the incident X-ray photon, since the **energy of the photon is inversely proportional to its wavelength**.

$$\text{The energy loss of the photon} = \text{energy gained by the recoil electron}$$

4.4 Quantifying the Model

Using the **conservation of energy and momentum respectively**, we obtain a set of simultaneous equations, from which, we get

$$\lambda' - \lambda_0 = \Delta\lambda = \frac{h}{m_e c} (1 - \cos \theta) \quad (3)$$




Figure 3: The Compton Effect

In this case, the quantity $\Delta\lambda$ is known as the **Compton shift**. The scattered photon's wavelength is increased because some of its original energy is transferred to the electron, in which process it loses energy itself and hence increases in wavelength.

4.5 Explaining the Peaks

1. The larger peak comes from the X-ray photon losing energy and hence increasing in wavelength due to the collision with an electron. This positive change in wavelength is given by Equation (3)
2. The smaller peak is created in the occasions where the X-ray photon **collides with the whole of the carbon atom**, and not just the electron. In this case, the modified equation for $\Delta\lambda$ requires m_e to be changed to m_{carbon} , which is about 20000 times larger. This means that the change in wavelength is about 20000 times smaller, effectively 0. Thus, we get a smaller peak that roughly matches the original wavelength.

5 Alternative Photon-Interaction

The previously described effects of the photoelectric effect and the Compton effect are not the only ways photons can interact with matter. They only apply to photons in certain energy ranges.

- The photoelectric effect is most prominent among photons ranging from a few eV to a few keV.
- The Compton effect is most prominent among photons ranging from a few keV to 1 MeV.
- Between 1.022 and 2 MeV, the photons can interact with the nucleus of the atom to create an electron-positron pair. The energy of the incident photon must be at least sufficient to provide the rest mass energy of the particles in the electron-positron pair respectively.
- At even higher energies, **photodisintegration** can happen, where the photon is absorbed by the nucleus, entering into a high-energy state, and then decaying with the ejection of a neutron, proton, or alpha particle.

5.1 Matter Waves and the de Broglie Hypothesis

This hypothesis is the claim that electrons and other particles can exhibit wave-only properties such as diffraction and interference under the appropriate experimental condition.

Since photons move at relativistic speeds, one can use Einstein's relativistic equation linking momentum and energy and obtain Equation (2). This gives the de Broglie wavelength of a particle moving at a relativistic speed.

5.2 The Davisson-Germer Experiment

This experiment (1927) provided evidence for the wave nature of electrons, confirming Louis de Broglie's hypothesis that particles can exhibit wave-like properties.

- **Electron Beam:** Electrons were accelerated and directed at a nickel crystal target in a vacuum chamber.
- **Scattering and Detection:** The electrons scattered off the atoms in the crystal, and a detector was used to measure the number of electrons scattered at different angles.
- **Observation:** The experimenters observed that at certain angles, the intensity of the scattered electrons was much higher, indicating constructive interference — a hallmark of wave behavior.

Key result: The pattern of electron scattering closely matched the diffraction pattern expected if the electrons were behaving like waves rather than particles. The wavelength of the electrons, calculated from the diffraction pattern, matched the value predicted by de Broglie's equation $\lambda = \frac{h}{p}$. In this case, the momentum of the electrons was given by $p = \sqrt{2m_e eV}$, where m is the mass of the electron and v is its velocity. Invoking de Broglie's hypothesis, the wavelength of the electrons was given by the following, where

V is the p.d. through which the electron is accelerated

$$\lambda = \frac{h}{mv} = \frac{h}{\sqrt{2m_e eV}} \quad (4)$$

De Broglie's hypothesis is thus inconsistent with Bohr's model of the atom: De Broglie's states that electrons are represented by waves and thus cannot be localized/have a fixed position.

In the exam, when describing the De Broglie hypothesis, make sure to include both of the parts below:

- A moving particle has wave properties
- The De Broglie wavelength of a particle is given by $\lambda = \frac{h}{p}$

6 Exam Questions

6.1 Intensity and Frequency

A metallic surface is first irradiated with infrared radiation and photoelectrons are emitted from the surface. The infrared radiation is replaced by ultraviolet radiation of the same intensity.

What will be the change in the kinetic energy of the photoelectrons and the rate at which they are ejected?

1. Firstly, UV has a higher frequency and thus higher energy than IR. This means that the photoelectrons will have a higher kinetic energy when UV is used.
2. Since each UV photon possesses more energy, there must be fewer photons arriving per unit time at per unit area to maintain the same intensity; thus the rate of ejection decreases.

6.2 Evidence for Wave/Particle

Monochromatic light of very low intensity is incident on a metal surface. The light causes the emission of electrons almost instantaneously. Explain how this observation

(i) does not support the wave nature of light.

- Low intensity light would transfer energy to the electron at a low rate/slowly
- Time would be required for the electron to absorb the required energy to escape/be emitted

(ii) does support the photon nature of light.

- In the photon theory of light, the electron interacts with a single photon
- and absorbs all the energy and gets ejected immediately.

6.3 Photoelectric Experiment — IV Graphs

In a photoelectric effect experiment, a beam of light is incident on a metallic surface W in a vacuum.




Figure 4: Diagram

The graph shows how the current I varies with the potential difference V when three different beams X, Y, and Z are incident on W at different times.




Figure 5: Graph

- I. X and Y have the same frequency.
- II. Y and Z have different intensity.
- III. Y and Z have the same frequency.

Explanation:

- Recall that the stopping potential is a direct measure of the maximum kinetic energy of the emitted electrons and thus the frequency of the incident wave. The stopping potential is the x -intercept of the graph — it is when the electrons do not have enough energy to overcome the potential difference and thus the current is 0. Any pair that has the same root will have the same frequency.
- To compare the intensity, we can consider the relative positions of the graph. In this case, Y and Z, although they have the same root, line Y is clearly above line Z for all values of V , this indicates that the current is higher for Y than for Z, and this means that more electrons are ejected by Y than by Z.

6.4 Energy Transfer — Electron and Photon

An electron of non-relativistic speed v interacts with an atom. All the energy of the electron is transferred to an emitted photon of frequency f . An electron of speed $2v$ now interacts with the same atom and all its energy is transmitted to a second photon. What is the frequency of the second photon?

- For the electron, we should consider its kinetic energy. Suppose the first electron has $E = \frac{1}{2}m_e v^2$, then, the second electron has $E = \frac{1}{2}m_e (2v)^2 = 4\frac{1}{2}m_e v^2 = 4E$.
- Now, we have to invoke the $E = hf$ relationship. Suppose the frequency of the first photon is f . Since the energy of the second electron is 4 times that of the first, the frequency of the second photon is $4f$.

6.5 De Broglie Wavelength and Momentum

An atom of hydrogen ${}^1_1\text{H}$ and an atom of helium ${}^4_2\text{He}$ are both moving with the same kinetic energy.

What is $\frac{\lambda_{\text{H}}}{\lambda_{\text{He}}}$?

- If the kinetic energies equal, and $m_{\text{He}} = 4m_{\text{H}}$, then, the speed of the helium atom is half of the speed of the hydrogen atom (recall that the speed is squared and so a factor of 1/4 in v^2 to compensate the factor of 4 in m should be square rooted to find the factor for v).
- We know that the de Broglie wavelength is given by $\lambda = \frac{h}{p}$, where $p = mv$.
- The momentum of the helium atom is scaled by a factor of 4 from m and a factor of 1/2 from v , so the total scaling factor is 2.
- The wavelength is inversely proportional to the momentum, so the wavelength of the helium atom is half that of the hydrogen atom.
- Thus, $\frac{\lambda_{\text{H}}}{\lambda_{\text{He}}} = 2$.

6.6 Compton Scattering

In a Compton scattering experiment, an X-ray photon interacts with a free, stationary electron. The electron recoils with an energy of 500 eV. The wavelength of the scattered photon is 7.8×10^{-11} m.

- (a) Show that the energy of the scattered photon is about 16 keV.

$$E = \frac{hc}{\lambda} = \frac{1.24 \times 10^{-6} \text{ eVm}}{7.8 \times 10^{-11} \text{ m}} \\ = 15.9 \text{ keV}$$

- (b) Determine the wavelength of the incident photon.

- Because the energy from the incident photon is transferred to the electron and the scattered photon, adding up the energies gives the total energy of the incident photon.

$$E_{\text{incident}} = E_{\text{scattered}} + E_{\text{electron}} \\ = 15.9 \text{ keV} + 0.5 \text{ keV} \\ = 16.4 \text{ keV}$$

- Then, we use the formula

$$\lambda = \frac{hc}{E}$$

to find the wavelength of the incident photon.

$$\lambda = \frac{1.24 \times 10^{-6} \text{ eVm}}{16.4 \times 10^3 \text{ eV}} \\ = 7.56 \times 10^{-11} \text{ m}$$

- (c) Outline why the results of the experiment are inconsistent with the wave model of electromagnetic radiation.

- According to the wave model, the wavelength of the incident X-ray photon should not change when it scatters off the electron.
- But the wavelength does change because the photon loses energy and momentum to the electron.

(d) Calculate the scattering angle of the photon.

$$\begin{aligned}\Delta\lambda &= \frac{h}{m_e c}(1 - \cos\theta) \\ (7.8 - 7.56) \times 10^{-11} &= \frac{6.63 \times 10^{-34}}{9.11 \times 10^{-31} \times 3 \times 10^8}(1 - \cos\theta) \\ \theta &= 89^\circ\end{aligned}$$

6.7 Photoelectric Effect

A beam of electrons is incident on a thin crystalline sample of graphite. The electrons emerging from the sample form a pattern on a fluorescent screen. The pattern consists of a series of bright and dark rings concentric with the direction of the incident beam.

(a) Outline why the pattern observed on the screen is an evidence for matter waves.

(a) The pattern is formed when the electrons scattered from adjacent planes in the graphite crystal undergo interference / diffract

(b) and interference and diffraction are wave-only properties.

The beam is produced by accelerating electrons through an electric potential difference U .

(b) A typical interatomic distance in the graphite crystal is of the order of 2×10^{-10} m.

Estimate the minimum value of U for the pattern in (a) to be formed on the screen.

- This question links back to diffraction: To get the pattern, the wavelength of the electrons **must be comparable** to the interatomic distance. Since the command word is estimate, this is a hint that we should take the wavelength to be equal to the interatomic distance.

$$p = \frac{h}{\lambda} = \frac{h}{2 \times 10^{-10}}$$

$$E = \frac{p^2}{2m_e} = \frac{h^2}{2m_e(4 \times 10^{-20})}$$

- If this is the energy required to get the diffraction pattern, then, we can divide it by the charge of the electron to get the potential difference.

$$U = \frac{E}{e}$$

$$= \frac{(6.63 \times 10^{-34})^2}{2 \times 9.11 \times 10^{-31} \times 4 \times 10^{-20} \times 1.6 \times 10^{-19}}$$

$$= 37.69... \approx 40 \text{ V}$$

(c) Protons can also be accelerated by the same potential difference U . Compare, without calculation, the de Broglie wavelength of the protons to that of the electrons.

- Protons have the same charge but a much larger mass than electrons.
- Since $E_K = \frac{p^2}{2m}$ is the same for both (accelerated through the same p.d. and have the same charge magnitude), the proton, with a much greater mass, will have greater momentum as well.
- Given that $\lambda = \frac{h}{p}$, with smaller momentum, the wavelength of the proton will be shorter than that of the electron.

6.8 Millikan's Experiment

In an experiment to investigate the photoelectric effect, a beam of monochromatic light is incident on a metal surface. When the frequency of light is less than a certain threshold frequency f_0 , no electrons are emitted from the surface.

(a) Outline how this observation rejects the wave model of light.

- The wave model of light predicts that there is a continuous transfer of energy from the beam of light to the electrons in the metal.
- Thus, eventually, the electrons would gain enough energy to escape from the surface, even if the frequency of the light is less than f_0 .
- But this does not happen; no electrons are emitted until the frequency of the light is greater than f_0 .

The collecting plate of the photoelectric cell is at a variable potential V relative to the metal surface.




Figure 6: Diagram

The graph shows the variation of the current I in the ammeter with V , for light of wavelength 430 nm.




Figure 7: Graph

(b) Explain why I

(i) is zero for large negative V .

- The reversed potential creates an electric field that opposes the motion of the emitted electrons...
- so electrons are repelled from and do not have enough KE to reach the photocathode.
- The maximum KE is less than the work needed to transfer it to the collecting plate.

(ii) approaches a constant value for large positive V .

- The idea here is that with a fixed intensity, there is a maximum number of electrons that can be emitted per second.
- Suppose we raise the p.d. to a high enough value where all the emitted electrons can reach the collecting plate. Increasing the p.d. further will not increase the number of electrons emitted per second and will only affect how quickly they reach the collecting plate.

(c) Calculate, in eV

- (i) the energy of a photon of wavelength 430 nm.

$$E = \frac{hc}{\lambda} = \frac{1.24 \times 10^{-6}}{430 \times 10^{-9}} = 2.9 \text{ eV}$$

- (ii) the work function of the metal surface.

$$\begin{aligned}\phi &= hf - E_{\text{max}} \\ &= 2.9 - eV_{\text{stop}}\end{aligned}$$

The stopping voltage is the intercept on the V -axis at which the current is 0.

This is at -0.9 V . Hence, $V_{\text{stop}} = -0.9 \text{ V}$

$$\begin{aligned}\phi &= 2.9 - (0.9) \\ &= 2.0 \text{ eV}\end{aligned}$$