

Biological Conservation

Estimates and drivers of protected species bycatch in the California set gillnet fishery

--Manuscript Draft--

Manuscript Number:	BIOCON-D-24-01680
Article Type:	Full Length Article
Keywords:	gillnet; bycatch; marine mammals; seabirds; area-based management
Corresponding Author:	Yutian Fang University of California Santa Barbara Santa Barbara, California UNITED STATES
First Author:	Yutian Fang
Order of Authors:	Yutian Fang James V. Carretta Christopher M. Free
Abstract:	The identification of efficient management strategies that reduce protected species bycatch while also minimizing impacts on fishing livelihoods is a global conservation challenge. Identifying such strategies requires understanding levels of bycatch relative to management targets as well as the relationship between bycatch risk and potential management levers. In this study, we use ratio estimators to reconstruct bycatch of select marine mammal and seabird species in the California ≥3.5" set gillnet fishery from 1981-2022 and random forest models to identify drivers and hotspots of bycatch risk. We find that bycatch has dropped precipitously since the 1980s as a result of management, but at significant costs to fisheries participation and revenues. Recent marine mammal bycatch ranges from 0.1% to 4.0% of the potential biological removal and marine mammal populations are recovering. Spatial-temporal drivers of bycatch risk were more important than fishing-related drivers of risk, suggesting that spatial-temporal closures would be more effective than mesh size or soak time restrictions at further limiting bycatch. For each species, we identified 2-5 hotspots of elevated bycatch risk as candidates for temporary seasonal closures. Bycatch risk for harbor seal (<i>Phoca vitulina</i>) and California sea lion (<i>Zalophus californianus</i>), the species with the greatest bycatch risk, is especially high from April 1st to June 15th, suggesting that hotspot closures during this 2.5-month time period could be especially efficient. Our study also highlights the value of competing multiple modeling approaches to identify methods that best predict rare bycatch events.
Suggested Reviewers:	Margaret Siple NOAA Fisheries Alaska Fisheries Science Center margaret.siple@noaa.gov Expert on marine mammal bycatch assessment
	Maritza Sepúlveda Universidad de Valparaíso maritza.sepulveda@uv.cl Expert on marine mammal bycatch assessment
	Tessa Francis National Nature Assessment tfrancis@usgcrp.gov Expert on marine mammal bycatch assessment
	Arne Bjørge Institute of Marine Research arne.bjoerge@hi.no Expert on marine mammal bycatch assessment
	Elizabeth Babcock University of Miami ebabcock@miami.edu Expert on marine mammal bycatch assessment

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

BERKELEY • DAVIS • IRVINE • LOS ANGELES • MERCED • RIVERSIDE • SAN DIEGO • SAN FRANCISCO



SANTA BARBARA • SANTA CRUZ

Dear Editors,

On behalf of the author team, I'm pleased to submit our paper, "Estimates and drivers of protected species bycatch in the California set gillnet fishery", for consideration as a Research Article in *Biological Conservation*.

Bycatch, the incidental capture of non-target species in fisheries, is a significant global conservation and economic challenge. Ineffective bycatch management can reduce the profitability of fisheries and threaten endangered marine species. Area-based management, which closes large areas of ocean when bycatch levels are high, is a common bycatch reduction strategy. However, it can overly limit fishing grounds and unnecessarily reduce fishing effort and revenues. Alternatively, temporary hotspot closures or gear modifications can reduce bycatch of protected species while minimizing negative impacts to fisheries.

The California set gillnet fishery targets California halibut and white seabass along the California coast, and has had high bycatch of pinnipeds, small cetaceans, and seabirds. Depth and area closures since the 1980s have limited fishing grounds to southern California, reducing fishing revenues to 10% of historical levels. In this study, we use ratio estimation to reconstruct bycatch of six protected species in the set gillnet fishery and machine learning models to identify drivers and spatial-temporal hotspots of bycatch risk. We use these results to evaluate the effectiveness of previous management strategies and to provide recommendations on how to efficiently and effectively reduce bycatch with low impacts on the fishery.

We find that bycatch has dropped precipitously since the 1980s as a result of management, but at significant costs to fisheries participation and revenues. However, seasonally closing the predicted bycatch hotspots could provide equivalent bycatch reduction while releasing more fishing grounds to the fishery. These results are important for the future management of California set gillnet fishery, which is on the precipice of closure due to perceptions of high levels of protected species bycatch. Furthermore, by comparing two analytical tools in estimating spatial and temporal bycatch trends, we provide a transferable approach to the global fishing industry in assessing the effectiveness of area-based management.

The work in this study is all original research carried out by the authors, who all agree with the contents of the study and its submission to *Biological Conservation*. No part of the research has been published elsewhere, the manuscript is not being considered for publication elsewhere, and the manuscript has not previously been submitted to *Biological Conservation*. All funding sources are acknowledged. No ethics and other approvals were needed for this research.

Thank you for your consideration and we look forward to hearing from you.

Sincerely, on behalf of all authors,
Yutian Fang

Estimates and drivers of protected species bycatch in the California set gillnet fishery

Yutian Fang¹, James V. Carretta², Christopher M. Free^{3,1}

¹ Bren School of Environmental Science and Management, University of California, Santa Barbara, Santa Barbara, CA, 93106, USA

² Southwest Fisheries Science Center, National Marine Fisheries Service, La Jolla, CA, USA

³ Marine Science Institute, University of California, Santa Barbara, Santa Barbara, CA, 93106, USA

* **Corresponding author:** Bren School of Environmental Science and Management, University of California, Santa Barbara, 2400 Bren Hall, Santa Barbara, CA 93106-5131; yutianfang@umail.ucsb.edu

Acknowledgements

YF was funded by the Arnhold UC Santa Barbara-Conservation International Climate Solutions Collaborative and CMF was supported by The Nature Conservancy California. We are grateful to Paulo Serpa (CDFW), Miranda Haggerty (CDFW), and Kirsten Ramey (CDFW) for providing fishery observer and logbook data, and to Karin Forney (NOAA) for providing reports and observer data from the 1980s. We are also grateful to the hundreds of fishermen, observers, and agency scientists that have contributed to the collection of the analyzed data. Our visualizations of the data are compliant with the "rule-of-three" and all handling of the confidential data is compliant with our data sharing agreement. We thank Kinsey Matthews (CFGC), Kirsten Ramey (CDFW) and Susan Ashcraft (CFGC) for providing constructive feedback throughout this study. CDFW collects data from various sources for fisheries management and purposes, and data may be modified at any time to improve accuracy and as new data are required. CDFW may provide data upon request under a formal agreement. Data are provided as-is and in good faith, but CDFW does not endorse any particular analytical methods, interpretations or conclusions based upon the data it provides. Unless otherwise stated, use of CDFW's data does not constitute CDFW's professional advice or formal recommendation of any given analysis. CDFW recommends users consult with CDFW prior to data use regarding known limitations of certain data sets.

Declaration of Interest Statement

The authors declare no conflicts of interest related to this work.

1
2
3
4
56 1 **Estimates and drivers of protected species bycatch in the**
7 2 **California set gillnet fishery**
8
9
10
11
1213 3 **Abstract**
1415 4 The identification of efficient management strategies that reduce protected species bycatch while also
16 5 minimizing impacts on fishing livelihoods is a global conservation challenge. Identifying such strategies
17 6 requires understanding levels of bycatch relative to management targets as well as the relationship
18 7 between bycatch risk and potential management levers. In this study, we use ratio estimators to
19 8 reconstruct bycatch of select marine mammal and seabird species in the California $\geq 3.5''$ set gillnet
20 9 fishery from 1981-2022 and random forest models to identify drivers and hotspots of bycatch risk. We
21 10 find that bycatch has dropped precipitously since the 1980s as a result of management, but at significant
22 11 costs to fisheries participation and revenues. Recent marine mammal bycatch ranges from 0.1% to 4.0%
23 12 of the potential biological removal and marine mammal populations are recovering. Spatial-temporal
24 13 drivers of bycatch risk were more important than fishing-related drivers of risk, suggesting that spatial-
25 14 temporal closures would be more effective than mesh size or soak time restrictions at further limiting
26 15 bycatch. For each species, we identified 2-5 hotspots of elevated bycatch risk as candidates for temporary
27 16 seasonal closures. Bycatch risk for harbor seal (*Phoca vitulina*) and California sea lion (*Zalophus*
28 17 *californianus*), the species with the greatest bycatch risk, is especially high from April 1st to June 15th,
29 18 suggesting that hotspot closures during this 2.5-month time period could be especially efficient. Our study
30 19 also highlights the value of competing multiple modeling approaches to identify methods that best predict
31 20 rare bycatch events.
32
3334 21
35 22 **Keywords:** gillnet, bycatch, marine mammals, seabirds, area-based management
36
3738 23
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 24 **1. Introduction**

6 25 Bycatch, the accidental capture of non-target species in fisheries, presents a significant
7 26 conservation and economic challenge (Crowder and Murawski, 1998; Soykan et al., 2008). Bycatch of
8 27 large-bodied, slow-growing, low-productivity species such as marine mammals, sea turtles, and seabirds
9 28 (Crowder and Murawski, 1998; Read et al., 2006; Soykan et al., 2008) is of particularly high concern, as
10 29 the mortality of individuals in these vulnerable populations, often recovering from historical exploitation,
11 30 can threaten population collapse and even extinction (Geijer and Read, 2013; Read et al., 2006). As a
12 31 result, many countries have established strict mandates to limit bycatch of vulnerable species, which can
13 32 result in fisheries closures and other severe restrictions (Crowder and Murawski, 1998; Senko et al.,
14 33 2014). These management disruptions can have serious social, cultural, and economic impacts on fishing
15 34 communities (Smith et al., 2020). Due to the negative ecological, economic, and social impacts of fishery
16 35 bycatch, bycatch avoidance is an important objective for global fishery management. A sustainably
17 36 managed fishery with low bycatch can not only provide ecological benefits, but also social and economic
18 37 benefits by providing a sustainable source of income, food, and nutrition.

19 38
20 39 To guide effective bycatch reduction policies, it is important to understand the magnitude of
21 40 historical and recent bycatch as well as the drivers of bycatch in a fishery. Estimates of total bycatch are
22 41 needed to determine whether bycatch exceeds management targets or is on pace to exceed targets in the
23 42 near future (Bjørge et al., 2013; Geijer and Read, 2013; Read et al., 2006). Historical bycatch estimates
24 43 also offer insights into the effectiveness of past management interventions, which provide useful
25 44 benchmarks for adapting management in response to contemporary bycatch levels and trends.
26 45 Understanding the drivers of bycatch risk is critical to guiding effective and efficient management
27 46 adaptations. For example, determining whether bycatch is concentrated within specific areas or seasons
28 47 can support the design of time-area closures that prevent fishing when and where risk is high while
29 48 maintaining fishing opportunities elsewhere (Lewison et al., 2014; O’Keefe et al., 2023; Soykan et al.,
30 49 2008). Similarly, gear restrictions, soak time restrictions, or time of day restrictions can be used to curb
31 50 bycatch if there are strong relationships between bycatch risk and gear or other characteristics of fishing
32 51 (O’Keefe et al., 2023). Without this information, bycatch management must be precautionary to guarantee
33 52 compliance with protected species legislation, which could forego considerable fisheries yields.

34 53
35 54 Because observer programs, which place trained scientists on fishing vessels to collect bycatch
36 55 data, are costly and rarely cover all fishing trips, various analytical approaches have been developed to
37 56 estimate unobserved bycatch and to evaluate drivers of bycatch. Ratio estimation, which assumes that the
38 57 rate of bycatch in observed fishing trips is proportional to the rate for all fishing trips, is one of the most

common strategies (Cochran, 1977; Stock et al., 2019). The reliability of this simple approach increases if there are sufficient data to support estimates within meaningful spatial or temporal strata (e.g., regions, depth zones, seasons). However, ratio estimation can produce biased estimates if other factors (e.g., gear type, soak time, time of day) influence bycatch rates (ICES, 2007), if the observed trips are not representative of the unobserved trips, or if low sample sizes lead to spuriously low or high bycatch rates within a stratum (Martin et al., 2015; McCracken, 2004; Ortiz and Arocha, 2004; Rochet and Trenkel, 2005). Model-based approaches, which use statistical models to estimate bycatch, can overcome these limitations by incorporating a wider suite of covariates and are thought to produce better bycatch estimates (Stock et al., 2019). Additionally, these approaches can support management by identifying drivers of bycatch risk and by predicting detailed hotspots of risk (Long et al., 2024; Lopez et al., 2024; Stock et al., 2019).

The California set gillnet fishery would benefit from updated bycatch estimation due to growing concerns about the fishery's impact on protected marine mammals, which have led some conservation organizations to call for the fishery's closure (Birch et al., 2023; Birch and Shester, 2023). The fishery occurs in southern California and targets California halibut (*Paralichthys californicus*), white seabass (*Atractoscion nobilis*), and Pacific angel shark (*Squatina californica*), among other species. It is currently listed as a Category II fishery under the U.S. Marine Mammal Protection Act (MMPA), indicating that it presents a medium bycatch threat to protected marine species (NOAA, 2024). Bycatch of marine mammal and seabird species, including harbor porpoise (*Phocoena phocoena*), southern sea otter (*Enhydra lutris nereis*), and common murre (*Uria aalge*) was high during the 1980s and 1990s, prompting significant management interventions (Forney et al., 2001; Julian and Beeson, 1998). The fishery has also impacted pinniped species such as California sea lion (*Zalophus californianus*), harbor seal (*Phoca vitulina*), and northern elephant seal (*Mirounga angustirostris*). Total bycatch in the fishery has not been estimated since 2012 (Carretta et al., 2014) and there are concerns that bycatch may remain an issue (Birch et al., 2023; Birch and Shester, 2023).

A number of management actions have been taken to reduce bycatch in the California set gillnet fishery. During the 1980s, high bycatch of southern sea otters and common murres in central California (Barlow et al., 1994) led to a depth restriction that closed fishing inside of 40 fathoms (73 m) in 1987 (Forney et al., 2001). This restriction shut down the fishery in the San Francisco area, effectively pushing it south of Pigeon Point and into Monterey Bay and Morro Bay (**Fig. 1A-2**). In 1990, the state adopted Proposition 132 (CA Secretary of State, 1990), which went into effect in 1994 and banned the fishery in mainland state waters (0-3 nautical miles) and in waters within 1 nautical mile or 70 fathoms of depth,

1
2
3
4 92 whichever is less restrictive, around the Channel Islands to further reduce bycatch of protected marine
5 species (FGC §8610.1-8610.16). In 2002, the state expanded the existing depth restriction, closing fishing
6 inside of 60 fathoms (110 m) to avoid the harbor porpoise population in Central California (14 CCR
7 §104.1). This effectively closed the fishery in Monterey Bay and Morro Bay (**Fig. 1A-4**). Currently, the
8 fishery only operates in Southern California (south of Point Arguello) outside 3 nautical miles from the
9 mainland and outside 1 nautical mile or shallower than 70 fathoms (whichever is less) from the Channel
10 Islands.
11
12
13
14
15
16 99

17 100 Although these regulations have reduced bycatch in the set gillnet fishery, they have also greatly
18 reduced fishery participation. The implementation of the 40 fathom depth restriction in 1987 triggered a
19 precipitous decline in participation from ~400 vessels in 1987 to ~100 vessels in 1994. Since then,
20 participation has continued to decline, with only ~40 vessels active in 2022, and the vast majority (>90%)
21 of landings coming from just 13 vessels (CDFW, 2023) (**Fig. 1B**). Fishing effort has similarly decreased
22 from an estimated ~15,000 days of fishing in 1987 to 1,000 days in 2022 (**Fig. 1C**). This reduction in
23 effort has significantly reduced bycatch levels (Carretta et al., 2014) but at large costs to fishery revenues.
24 Fleetwide revenues decreased from US\$15 million in 1987 to only US\$1 million in 2022 (**Fig. 1C**; both
25 values in 2022 dollars). Despite declining fishing effort and bycatch, there are calls for additional
26 restrictions, including permanent closure, to further avoid bycatch (Birch et al., 2023; Birch and Shester,
27 2023). There is thus great need for scientific guidance on management regulations that are likely to
28 provide conservation benefits while also avoiding unnecessary burdens on the fishing industry.
29
30 112

31 113 In this study, we reconstruct the bycatch of select marine mammals and seabirds in the California
32 set gillnet fishery from 1981-2022 and identify drivers of bycatch that could be used to design
33 management measures that effectively and efficiently reduce bycatch. We focus on six protected species
34 encountered in the set gillnet fishery: California sea lion, harbor seal, northern elephant seal, harbor
35 porpoise, common murre, and Brandt's cormorant. Southern sea otter, despite being one of the original
36 species of management concern, is not included in this analysis due to data limitations (**Fig. 2A**). We use
37 ratio estimation methods to reconstruct historical bycatch levels. These methods, which have been used to
38 estimate bycatch in the fishery at various points in the past, provide a complete time series of bycatch
39 estimates using methods approved for stock assessment and management. We then use random forest
40 models to evaluate drivers of bycatch risk and to make predictions of spatial bycatch risk for four species
41 of concern; Brandt's cormorant and harbor porpoise were excluded due to poor model performance
42 (**Table 2**). Based on these results, we make recommendations for how management could effectively
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58 124
59
60
61
62
63
64
65

1
2
3
4 125 manage bycatch risk through measures such as seasonal or spatial closures, depth restrictions, or gear
5
6 126 restrictions.
7
8
9

10 127 **2. Methods**
11
12

13 128 **2.1 The fishery**
14

15 129 We defined the fishery using the definition in the MMPA List of Fisheries (NOAA, 2023): the
16 130 ≥ 3.5 inch mesh set gillnet fishery targeting California halibut, white seabass, Pacific angel shark, and
17 other species. Although this definition deviates from historical studies, which frequently focused on the
18 portion of the fishery using mesh sizes larger than 8.0 or 8.5 inches (**Table 1**), the MMPA definition
19 provides the legal basis for bycatch management and is more consistent with historical regulations.
20
21 132 Specifically, a minimum mesh size of 3.5 inches was set for white seabass in 1941, though it was
22 133 increased to 6.0 inches in 1988 (FGC §8623(d)). Since 1989, California halibut and Pacific angel shark
23 have been targeted using a minimum mesh size of 8.5 inches (FGC §8625(a)). The set gillnet fishery
24 principally excluded by this definition is that for Pacific herring (*Clupea pallasii*), which occurs in
25 California's four largest herring spawning areas — San Francisco Bay, Tomales Bay, Humboldt Bay, and
26 Crescent City Harbor — using mesh sizes of 2.0 to 2.5 inches (CDFW, 2019).
27
28
29
30
31
32
33
34

35 140 **2.2 Data**
36
37

38 141 **2.2.1 Observer data**
39

40 142 We received observer data from 1983 to 2017 from the California Department of Fish and
41 143 Wildlife (CDFW). There was observer coverage in the California set gillnet fishery from 1983-1995
42 (coastwide), 1999-2000 (Monterey Bay area only), 2010-2013 (south of Point Conception only), and in
43 144 2017 (south of Point Conception only) (**Fig. 2**). The observer program was run by CDFW from 1983-
44 145 1989 and by the National Oceanic and Atmospheric Administration (NOAA) from 1990 onwards.
45
46 146 Observers collected information on the amount and fate of catch (kept, discarded, or damaged), the length
47 147 composition of the catch, the location and time of the catch, and characteristics of the gear used to target
48 148 the catch (**Fig. S2**). In the state observer data, we defined individual gillnet sets based on the date of
49 149 fishing, the vessel, and the set number and built a unique identifier to link set-level information across
50 state datasets (i.e., YYYY-MM-DD-VesselID-Set#). In the federal observer data, we defined individual
51 150 gillnet sets based on the observer trip number and the set number and built a unique identifier to link set-
52 151 level information across federal datasets (i.e., TripID-Set#).
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 154
5
6 155 Although historical reports document low levels of observer coverage in Morro Bay, Monterey
7 Bay, and San Francisco in the 1980s (**Table 1**), the data that we received from CDFW excluded most of
8 these observations. We digitized a small portion of the missing raw data – observations from Monterey
9 Bay from 1987-1989 (**Fig. 2**) – from original CDFW data sheets that were given to a colleague at the
10 Southwest Fisheries Science Center during the late 1990s for a reanalysis of historic bycatch rates in that
11 region (Forney et al., 2001). We extracted summaries of bycatch rates from historical reports (**Table 1**)
12 for years and regions missing raw data to support the ratio estimation analysis. **Figure 2C** illustrates the
13 coverage of the available, recovered, and lost observer data.
14
15 161
16 162
17
18 163
19
20
21 164 We developed a series of simple assumptions to impute missing values for a few key variables
22 165 (GPS coordinates, fishing depth, soak hour, mesh size) reported in the observer data (**Fig. S3A**). We
23 166 assigned missing GPS coordinates using the median coordinates for observed trips within the statistical
24 167 block most frequently visited by the vessel – in order of preference – that week, month, or year based on
25 the logbook data (described below). We derived missing fishing depths by extracting depths from 25-
26 168 meter resolution bathymetry data (CDFW, 2002) (**Fig. S3B**). We reassigned missing soak hours the mode
27 169 value for a vessel and target species (**Fig S3C**). We reassigned missing mesh sizes the mode for – in order
28 170 of preference – the vessel and target species, the target species, or all vessels (**Fig. S3DE**). We assigned
29 171 each GPS coordinate to the nearest statistical reporting block (see **Fig. 1A**), which allows points
30 172 erroneously falling on land to be assigned a likely statistical block. We derived the distance from shore, a
31 173 covariate used to explain bycatch rates in the random forest model, as the distance of each set to the
32 174 nearest point on shore.
33
34 175
35
36 176
37
38 177 For each species of marine mammal, seabird, and sea turtle documented in the observer data, we
39 178 calculated the total number of captures observed in the California set gillnet fishery (**Fig. 2**). Throughout
40 179 this analysis, we focus on the six species with more than 50 observations: Common murre (2,381),
41 180 California sea lion (1,372), harbor seal (519), Brandt's cormorant (118), Northern elephant seal (78), and
42 181 harbor porpoise (97). Unfortunately, this excludes southern sea otter, which was a species of significant
43 182 conservation concern in the 1980s, but whose bycatch was only documented in the lost observer data.
44
45
46
47
48
49
50
51
52
53
54 183 2.2.2 Logbook data
55
56
57 184 We received logbook data from the commercial gillnet fishery from 1981 to 2022 from CDFW
58 185 (**Fig. 1**). These data describe vessel information (vessel name, unique identifier, permit number); when
59 186 (date), where (block id), and how long (hours) a vessel fished; what fish it targeted; what type of gear it
60
61
62
63
64
65

1
2
3
4 187 used (drift or set gillnet) and characteristics of this gear (length, mesh size, fishing depth; **Fig. S2**); what
5 species it caught; and the amount (number and/or weight) and fate of this catch (kept, released, or lost,
6 including the identity of predators preying on released fish). We attempted to identify individual fishing
7 sets as the unique combination of vessel administration information (vessel name, either vessel id or boat
8 number, permit number), where, when, and how long a vessel fished (block id, date, and fishing hours),
9 and characteristics of the gear (net length, mesh size, and fishing depth). This analysis revealed an
10 average of 1 set per trip, which is inconsistent with the 3 sets per trip documented in the more accurate
11 observer data. We term these unique identifiers “pseudo-sets” and view them as roughly equivalent to a
12 fishing trip (**Fig. S4**). We derived the distance from shore, a covariate used to explain bycatch rates in the
13 random forest model, as the median distance from shore of observed trips within the reported block given
14 that exact locations are not reported.
15
16 198
17
18 199 We developed a series of simple assumptions to impute missing or unrealistic values for a few
19 key variables (fishing depth, soak hour, mesh size) reported in the logbook data (**Fig. S5A**). We
20 reassigned both missing (including 0 values) and unrealistic fishing depths, which we defined as depths
21 exceeding the maximum depth in the reported fishing block, the median depth of the fishing block (**Fig.**
22 **S5B**). We computed the median and maximum depths of each fishing block using 25-meter resolution
23 bathymetry data (CDFW, 2002). We reassigned missing soak hours (including 0 values) the mode value
24 for a vessel. We capped rare and unlikely soak times exceeding 96 hours (4 days) at 96 hours; however,
25 such soak times could theoretically occur during rough weather when it is unsafe to haul gear (**Fig. S5C**).
26 We reassigned missing (including 0 values) and unrealistic mesh sizes, which we defined as mesh sizes
27 exceeding 20 inches, using a hierarchical procedure (**Fig. S5DE**). For logbooks with both vessel
28 identification and target species information, we assigned the mesh size most commonly used by the
29 vessel when targeting that target species. For logbooks with only target species information (no vessel
30 identification), we assigned the mesh size most commonly used when targeting that target species across
31 all vessels (**Figs. S5 & S6**).
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49 213 2.2.3 Sea surface temperature
50
51 214 Because sea surface temperature (SST) is a common driver of the distributions of both target and
52 bycatch species (Hazen et al., 2018), we used SST as an environmental covariate in the random forest
53 models described below. We derived the SST associated with each set documented in the observer and
54 logbook data using the NOAA 1/4° Daily Optimum Interpolation Sea Surface Temperature (OISST)
55 dataset, which interpolates observations from different monitoring platforms (e.g., satellites, ships, buoys,
56 and Argo floats) to provide a globally complete grid of SST from September 1, 1981 to present (Huang et
57
58 219
59
60
61
62
63
64
65

50
51 213 2.2.3 Sea surface temperature
52
53 214 Because sea surface temperature (SST) is a common driver of the distributions of both target and
54 bycatch species (Hazen et al., 2018), we used SST as an environmental covariate in the random forest
55 models described below. We derived the SST associated with each set documented in the observer and
56 logbook data using the NOAA 1/4° Daily Optimum Interpolation Sea Surface Temperature (OISST)
57 dataset, which interpolates observations from different monitoring platforms (e.g., satellites, ships, buoys,
58 and Argo floats) to provide a globally complete grid of SST from September 1, 1981 to present (Huang et
59
60
61
62
63
64
65

al., 2021). For sets reported in the observer data, we extracted the SST at the reported GPS location on the reported day of fishing. For sets reported in logbooks, we calculated the average SST in the reported block on the reported day of fishing.

2.3 Ratio estimation

We estimated annual bycatch for each study species using ratio estimators. Ratio estimators assume that the rate of bycatch in observed fishing trips is proportional to the rate of bycatch within all fishing trips within a given stratum (Cochran, 1977). This assumption requires that the characteristics of observed trips do not systematically differ from the characteristics of all trips, which was confirmed by a two-sided Kolmogorov-Smirnov test (**Fig. S7**). We used trips rather than sets as the sampling unit given the inability to identify unique sets in the logbooks data (**Fig. S4**). Under this approach, the bycatch rate for species s in stratum i ($r_{s,i}$) is thus calculated as:

$$r_{s,i} = \frac{k_{s,i}}{d_{s,i}}$$

Where $k_{s,i}$ is the total number of individuals of species s captured in observed trips occurring in strata i and $d_{s,i}$ is the total number of observed trips occurring in strata i . The total estimate of bycatch of species s in strata i ($m_{s,i}$) is then calculated by multiplying the bycatch rate ($r_{s,i}$) by the total number of trips to have occurred in the strata (D_i):

$$m_{s,i} = D_i * r_{s,i}$$

Where the total number of trips (D_i) is derived from the logbook data.

We calculated annual bycatch estimates using a seven-region stratification scheme (**Figs. 1A, S8**). This stratification scheme combines the scheme by Diamond and Hanan (1986) and Julian (1993) for areas north and south of Point Conception, respectively. Although early efforts to estimate bycatch in the California set gillnet fishery often stratified estimates by region and season (**Table 1**), later efforts found that observer coverage was often too limited to employ complex temporal stratification and that estimates between temporally stratified and unstratified approaches were generally similar (**Table 1**). Stratum-specific bycatch rates for years without observer coverage in the stratum are borrowed from the closest year (forwards or backwards) with observer coverage in the stratum (**Fig. 2C & S9**). We collated annual bycatch estimates from past studies (**Table 1**) for comparison with our updated estimates (**Figs. S10-S12**).

1
2
3
4 252 Although there are methods for estimating the uncertainty of bycatch estimates generated through
5 ratio estimation (Julian and Beeson, 1998), we were unable to implement these methods because they rely
6 on bootstrap procedures that sample from the bycatch rates of observed trips. Because these procedures
7 require raw observer data, we cannot use them for (1) species-strata-years where summary values from
8 historical reports are used because the raw data have been lost or (2) species-strata-years without observer
9 data. An exploration of the uncertainty estimates from historical reports (**Table 1**) suggest that the median
10 coefficient of estimate for annual bycatch estimates ranges from 0.14 for harbor seal to 0.47 for harbor
11 porpoise (**Figure S10B**).
12
13
14
15
16
17
18
19 260 2.4 Random forest modeling
20
21
22 261 2.4.1 Model training
23
24
25 262 We used random forest classification models trained on the observer data to identify drivers of
26 bycatch risk for each of the six study species. We considered nine attributes of fishing as potential drivers
27 of bycatch risk: haul depth (fathoms), mesh size (inches), soak time (hours), latitude ($^{\circ}$ N), longitude ($^{\circ}$ W),
28 distance from shore (km), Julian day, sea surface temperature ($^{\circ}$ C), and whether the fishing occurs near an
29 island (i.e., within 10 km of island coast). For each species, we classified an observed set as having (1) or
30 not having (0) bycatch, and trained a classification model assuming a Bernoulli distribution in the
31 response variable.
32
33
34
35
36 269
37
38 270 Because bycatch of protected species is rare, observed fishing sets show strong class imbalance
39 towards sets without bycatch compared to sets with bycatch. To illustrate, the percent of observed sets
40 with bycatch is as follows: California sea lion (1.02%), common murre (0.43%), harbor seal (0.43%),
41 Brandt's cormorant (0.09%), northern elephant seal (0.076%), and harbor porpoise (0.074%). Therefore,
42 without a proper sample balancing method, predictions are likely to be biased towards the majority class
43 (sets without bycatch), leading to an underestimation of bycatch risk. For this reason, we considered four
44 approaches for accounting for class imbalance resulting from bycatch rarity.
45
46 277
47
48 278 The first three approaches employ different sample balancing methods: (1) downsampling, (2)
49 upsampling, and (3) synthetic minority over-sampling (SMOTE), which uses a mixture of down and
50 upsampling (More and Rana, 2017). The downsampling approach randomly removes observations of the
51 majority group (sets without bycatch) to obtain equal representation of the majority and minority (sets
52 with bycatch) group. The upsampling approach randomly samples observations from the minority group
53 with replacement to obtain equal representation with the majority group. The synthetic minority over-
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 284 sampling (SMOTE) approach both up-samples the minority group and down-samples the majority group.
5
6 285 It up-samples the minority class by synthesizing new cases from its nearest five neighbors and down-
7 samples the majority class by randomly drawing samples from that group. We created each balanced
8 dataset using the *themis* package in R (Hvitfeldt, 2023) and fit random forest models to these datasets
9 using the *randomForests* package in R (Liaw and Wiener, 2002).
10
11 288
12 289
13

14 290 The fourth approach employs weighted random forests, which use observation weighting rather
15 than sample balancing to elevate the importance of the minority class. In this “cost-sensitive” learning
16 approach (More and Rana, 2017), higher weights are assigned to minority observations so that the model
17 receives a higher penalty for misclassifying these observations, helping to reduce bias towards the
18 majority class. We evaluated multiple weighting schemes to optimize the predictive skill of this approach.
19 Specifically, we assigned majority observations (sets without bycatch) a weight of 1 and assigned
20 minority observations (sets with bycatch) weights of 25 to 200 in increments of 25. Thus, a total of eight
21 candidate weighted random forest models were evaluated as described below. We fit the weighted
22 random forest model using the *ranger* package in R (Wright and Ziegler, 2017).
23
24 295
25
26 297
27
28 298
29 299
30

31 300 We trained each of the eleven candidate models (three balanced random forest models, eight
32 weighted random forest models) on 80% of the observer data, withholding the remaining 20% for model
33 testing. In training the models, we performed a grid search to identify the “mtry” hyperparameter – the
34 number of variables to randomly sample as candidates at each node split – that maximizes Cohen’s kappa
35 under 10-fold cross validation (**Fig. S13**). While accuracy measures the proportion of correctly classified
36 categorizations, Cohen’s kappa measures the proportion of correct classifications while accounting for the
37 probability of being correct by chance and is a better measure of predictive skill, especially for
38 imbalanced datasets (Cohen, 1968). Although there are no definitive rules for interpreting Cohen’s kappa,
39 general guidelines suggest that values above 0.7 are ‘excellent’, 0.4-0.7 are ‘good’, 0.2-0.4 are ‘fair’, and
40 below 0.2 are ‘poor’ (Fleiss et al., 2013; Landis and Koch, 1977). We identified the best fitting model as
41 the model generating the highest Cohen’s kappa on the training data. We applied this model to the test
42 dataset for an independent evaluation of its predictive power. We only evaluated four species (California
43 sea lion, common murre, harbor seal and northern elephant seal) whose best models exhibited “fair” or
44 better performance in their training dataset and close to “fair” performance on the test dataset for the rest
45 of the analysis (**Table 2**).
46
47
48
49
50
51
52
53
54
55

56
57
58
59
60
61
62
63
64
65

1
2
3
4 315 2.4.2 Model evaluation and prediction
5

6 316 We evaluated the drivers of bycatch risk for each species by inspecting the variable importance
7 and the marginal effects of each variable as estimated in the best fitting model. Variable importance was
8 evaluated as the total decrease in node impurities from splitting on the variable averaged over all trees.
9
10 318 The impurity measure is corrected when building the model to reduce its bias towards continuous
11 variables (Nembrini et al., 2018). Marginal effects measure the impact of the changes in one variable on
12 the response variable while all other variables are held constant. The marginal effects plots provide the
13 scientific basis for our discussions of management regulations that could effectively and efficiently
14
15 322 reduce bycatch risk.
16
17 323
18
19 324

21 325 We used the best fitting model to generate annual estimates of protected species bycatch from
22 1981 to 2022 by predicting whether “pseudo-sets” recorded in logbooks were likely to have captured each
23 study species. We predict to pseudo-sets rather than trips because the random forest model is trained on
24 set-level covariates in the observer data. We used the best fitting model for each species to estimate the
25 probability that a logged pseudo-set included bycatch of a species then categorized the pseudo-set as with
26 or without bycatch using a species-specific probability threshold. We derived the species-specific
27 probability thresholds as the threshold that maximizes Cohen’s kappa when applied to the training
28 datasets (**Fig. S14**). We selected the probability threshold based on Cohen’s kappa rather than the area
29 under the receiver operator curve (AUC) because (1) the models were tuned and selected based on
30 Cohen’s kappa and (2) simulation work shows that deriving thresholds based on AUC tends to
31 overestimates the prevalence of rare events while it underestimates the prevalence of common events
32 (Freeman and Moisen, 2008; Manel et al., 2001). We summed the number of pseudo-sets predicted to
33 have bycatch each year, converted this sum to “true sets” assuming three sets per pseudo-set (**Fig. S4AB**),
34 and multiplied this sum by the median number of captures when a capture occurs to generate estimates of
35 the total number of captured animals (**Fig. S4C**). We opted not to employ a more complex two-stage or
36 hurdle model approach, where a second model estimates the number of captured individuals when
37 bycatch occurs, given the rarity of bycatch events larger than one for all species but common murre (**Fig.**
38 **S4C**).
39
40 343
41
42 344 To generate maps of spatial bycatch risk, we used the best fitting model to predict risk to a 0.02°
43 grid spanning southern California ($32\text{-}35^{\circ}\text{N}$ and $117\text{-}121^{\circ}\text{W}$), the only area where the fishery can operate
44 under current regulations. Conceptually, our metric of spatial bycatch risk represents the probability of
45 bycatch at a given location under recent average conditions. We derived this metric by first predicting the
46 probability of bycatch on every calendar day (Julian day 1 to 365) for each grid cell. The eight input
47
48 348
49
50 342
51
52 343
53
54 344
55
56 345
57
58 346
59
60 347
61
62
63
64
65

variables for making these predictions were derived as follows: (1) *latitude*, (2) *longitude*, (3) *distance from shore*, and (4) *depth* were derived based on the centroid of the grid cell; (5) *soak time* was set to 24 hours, the logbook mode; (6) *mesh size* was set to 8.5 inches, the logbook mode; (7) *Julian day* was the input day; and (8) *sea surface temperature* (SST) was the average SST for that day and grid cell from 2010-2022. We then calculated the weighted average of the daily risk weighting by the amount of logbook entries (a metric of fishing effort) occurring on that Julian day from 2010-2022. We cropped the predictions to areas within 20 km of shore as this is the farthest offshore that the fishery has operated (**Fig. S7**).

3. Results

3.1 Ratio estimation

In general, estimated bycatch peaked in the mid-1980s, steadily declined following the 40-fathom depth restriction implemented in 1987, with a temporary increase in the late-1990s followed by continued decline (**Fig. 3**). This pattern reflects trends in fishing effort, which also declined after a peak in 1985, with a brief expansion in the late-1990s followed by continued decline (**Fig. 1C**). Estimated bycatch of Brandt's cormorant follows a similar pattern but lags behind the patterns for other species: estimated bycatch precipitously declines after a peak in 1989 with a temporary increase occurring in the early-2000s (**Fig. 3**). This pattern is driven by a steep increase in Brandt's cormorant bycatch rates in the Channel Islands region in 1990 that is assumed to have persisted to today (**Fig. S9**). None of the other species experienced such a pronounced and impactful change in bycatch rates (**Fig. S9**). Estimated bycatch of harbor porpoise and common murre declined especially sharply following the 2002 exclusion of fishing from waters shallower than 60 fathoms. This regulation effectively closed the fishery in Monterey Bay and Morro Bay and restricted it to only Southern California (**Fig. 1A**), where harbor porpoise do not occur and where common murre occur only in winter in low densities (**Fig. S15**).

The sustainability of estimated marine mammal bycatch can be weighed against their potential biological removal (PBR) for each stock, which is defined under the MMPA as the maximum number of animals, not including natural mortalities, that may be removed from a marine mammal stock while allowing that stock to reach or maintain its maximum sustainable population (**Fig. 7**). Based on this sustainability reference point, bycatch concerns, in order of decreasing threat, are as follows: harbor seal (65 individuals per year = 4.0% of a PBR of 1,641 individuals), harbor porpoise (1 individual per year = 1.5% of a PBR of 65 individuals in the Morro Bay stock, where the bycatch took place), California sea

lion (194 individuals per year = 1.4% of a PBR of 14,011 individuals), and northern elephant seal (7 individuals per year = 0.1% of a PBR of 5,122 individuals) (**Fig. 7**). The assessment that bycatch during the last 10 years poses the greatest risk to harbor seals is supported by the fact that the harbor seal stock size has been stable or declining in recent years while all of the other marine mammal populations have been undergoing sustained population growth (**Fig. 7**). The sustainability of estimated seabird bycatch is more difficult to evaluate given more limited population monitoring data (**Fig. 7**) and the lack of legally binding reference points for defining allowed incidental take. However, increasing Brandt's cormorant nests from 1980 to 2020 (**Fig. 7**) and steeply reduced bycatch of common murre (**Fig. 3**) suggests low risks posed to these species.

Our estimates of annual bycatch are generally aligned with estimates from historical studies (**Fig. S11**). Slight differences between our estimates and those from historical studies are driven by a mixture of differences in our methods and input data. While we apply a consistent approach for defining the fishery, stratifying the data, and estimating bycatch, historical studies have employed variable fishery definitions, stratification schemes, and estimation methodologies (**Table 1**). First, we consistently defined the fishery as using $\geq 3.5"$ set gillnets, while historical studies have considered set gillnets $\geq 8"$, $\geq 8.5"$, or of unspecified sizes (**Table 1**). Furthermore, it is unclear whether historical studies filtered out fishing sets based on the reported target species and whether this decision was consistent. We do not define the fishery based on reported target species given extreme heterogeneity in how this information is reported. Differences in fishery definitions, as well as differences in data cleaning methods, likely lead to the slight differences in effort and observed bycatch attributed to the fishery (**Fig. S12**), which inevitably causes differences in bycatch estimates. Second, we used a consistent six-region stratification scheme, while historical studies used a mixture of stratification schemes ranging from no stratification to spatial stratification to spatial-temporal stratification (**Table 1**). This impacts the ratio estimators and the magnitudes of bycatch. Finally, we used vessel days (trips) as the sample unit, whereas historical studies oscillated between trips and sets as the preferred sample unit (**Table 1**). This is problematic not only because of the lack of consistency but also because sets cannot be uniquely identified in the logbook data.

3.2 Random forest modeling

3.2.1 Model performance

The best fitting model performed well for four of the six study species: common murre, harbor seal, California sea lion, and northern elephant seal. All four species utilized weighted random forests but with different case weights (**Table 2; Fig. S13**). The models for two species, Brandt's cormorant and

1
2
3
4 412 harbor porpoise, were excluded from further consideration because of their poor performance (**Table 2**).
5
6 413 Weighted random forest models performed best for California sea lion, common murre, northern elephant
7 seal, and harbor seal with case weights of 25, 25, 25, and 75, respectively (**Table 2**). This highlights the
8 importance of evaluating multiple modeling approaches when predicting rare bycatch events. Model
9 performance was strongly correlated with the frequency of bycatch observations (**Table 2**). The best
10 fitting model for common murre, the most common bycatch species, exhibited a Cohen's kappa value
11 greater than 0.4 indicating "good" performance while the best models for harbor seal and California sea
12 lion, somewhat common bycatch species, exhibited values greater than 0.2 indicating "fair" performance.
13
14 418 The best fitting models for Brandt's cormorant and harbor porpoise, rare bycatch species, exhibited
15 Cohen's kappa values less than 0.2 on the test dataset, indicating "poor" performance, and were not
16 evaluated any further. Although northern elephant seal bycatch is also rare, its best fitting model exhibited
17 "fair" performance, and it was evaluated further. Cohen's kappa values were correlated with the area
18 under the receiver operator curve (AUC), indicating minimal tradeoffs in using this metric for model
19 selection (**Table 2**).
20
21
22
23
24
25
26
27
28
29 426 3.2.2 Drivers of bycatch risk
30
31 427 The importance of the evaluated explanatory variables in determining bycatch risk varied by
32 species but some general patterns emerged (**Fig. 4**). In general, spatial (latitude, longitude, depth, and
33 428 distance from shore) and temporal (Julian day) variables were more influential than variables associated
34 429 with the environment (sea surface temperature) or the fishing methodology (soak time, mesh size).
35
36 430 Whether fishing occurred close to an island (a spatial variable) was the exception, as it was consistently
37 431 the least important variable. Sea surface temperature, which is closely related to space and time, was
38 432 generally more important than soak time and always more important than mesh size. These results suggest
39 433 that spatial-temporal management may have better ability to manage bycatch risk than gear modifications
40 434 or soak time regulations, though shorter soak times could reduce discard mortality for non-target fish
41 435 species.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

4 438 The species exhibit a mixture of correlated and non-correlated responses to the explanatory
5 variables (**Fig. 5**). California sea lion and harbor seal exhibit similar responses in bycatch risk. Both
6 species have higher bycatch risk in shallower depths in nearshore areas with a spike in risk occurring
7 around 34°N latitude. They also exhibited a pronounced increase in risk during the spring, lasting
8 approximately from Apr 1 (90th day of the year) to June 15 (166th day of the year). They are infrequently
9 caught in nets with mesh sizes smaller than 8.5 inches, though the use of such nets is rare (**Fig. S5E**).
10 Variability in bycatch risk for common murre and northern elephant seal is most strongly determined by
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 445 latitude and longitude (**Fig. 5**), with the only area of elevated risk in contemporary fishing grounds
5 occurring just north of Point Conception (**Fig. 6**). For all four species, bycatch risk exhibits an asymptotic
6 relationship with soak time and a dome-shaped relationship with temperature; however, the shapes of
7 these relationships differ by species.
8
9 448
10
11
12 449 3.2.3 Maps of bycatch risk
13
14 450 California sea lion bycatch risk is predicted to be highest in four areas: (1) on the northern coasts
15 of the northern Channel Islands, especially on the northern coast of Santa Rosa; (2) a small nearshore area
16 west of Santa Barbara; (3) the eastern coast of Santa Cruz Island; and (4) the northwestern shores of Santa
17
18 452 Catalina and San Clemente Islands; (**Fig. 6**). The high risk around the northern Channel Islands is likely
19
20 453 related to the large haulouts of sea lions in that area (**Fig. S15**).
21
22
23 455
24
25 456 Harbor seal bycatch risk is predicted to be highest in four areas: (1) the sliver of nearshore area
26 stretching from Santa Barbara to Point Sal; (2) the eastern coasts of Santa Cruz Island; (3) a broad coastal
27
28 458 area near Point Mugu; and (4) the sliver of nearshore area stretching from Point Mugu to Point Vicente
29
30 459 (**Fig. 6**). Most of these areas correspond to the locations of large harbor seal haulouts on Santa Cruz
31
32 460 Island and near Point Mugu (**Fig. S15**).
33
34 461
35
36 462 Common murre bycatch risk is predicted to be negligible throughout most of southern California
37
38 463 (**Fig. 6**). It is only predicted to be high in a small patch near Point Sal and even there, the maximum risk
39
40 464 index is much lower than for the other evaluated species. This is consistent with the distribution of the
41
42 465 species, which has no breeding colonies or permanent foraging grounds in southern California (**Fig. S15**).
43
44 466 The absence of common murre in southern California largely explains the significant drop in common
45
46 467 murre bycatch (**Fig. 3**) since the fishery was pushed out of central California (**Fig. 1**).
47
48
49 469 Like common murre, northern elephant seal bycatch risk is also predicted to be negligible
50 throughout most of southern California except in the region near Point Sal (**Fig. 6**). Although northern
51
52 471 elephant seals breed on the Channel Islands and near San Simeon/Cambria (**Fig. S15**) from December to
53
54 473 March, they disperse to their distant foraging grounds (males to Alaska and females to oceanic waters far
55
56 West of California) before the fishing season peaks from April to June, significantly reducing their
57
58 vulnerability to the gillnet fishery.
59
60
61
62
63
64
65

1
2
3
4 475 3.2.4 Temporal trends in estimated bycatch
5

6 476 The random forest models estimate trends in bycatch that are similar to the estimates from the
7 ratio estimator from 2000-2022 (**Fig. 3**). The agreement between these two different approaches
8 underscores the result that bycatch has decreased to low levels as a result of management and reduced
9 fishing effort. However, the estimates produced by the two approaches diverge from 1981-2000 by
10 various extents. While they generally agree for California sea lion back to 2000, the random forest model
11 underpredicts bycatch relative to the ratio estimator in the late 1990s and overpredicts in the 1980s and
12 early 1990s (**Fig. 3**). While the approaches generally agree for harbor seal back to 1995, the random forest
13 model underpredicts bycatch relative to the ratio estimator before 1995, especially in the Channel Islands
14 and Ventura strata (**Fig. S16**). For common murre, the random forest model overpredicts bycatch relative
15 to the ratio estimator in the mid- to late-1990s and underpredicts relative to the ratio estimator in earlier
16 years, especially in Morro and Monterey Bays. These underpredictions likely occur because of the
17 unequal impacts of lost data from the northern strata in the 1980s (**Fig. 2**). While the ratio estimation
18 method leverages summarized observer data from old reports, meaning that it sees data from this time
19 period, the lack of the raw observer data from this time period means that the random forest model does
20 learn from this period. As a result, it is likely to underpredict risk in early years in northern strata because
21 it has largely learned from late years in southern California, where risk has been lower. For this reason,
22 we recommend the use of bycatch estimates from the ratio estimator over the random forest model until a
23 time when the 1980s observer data is rediscovered.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38 494 4. Discussion
39
40

41 495 Our study provides the first update to total estimates of protected species bycatch in the
42 California set gillnet fishery since 2012 (Carretta et al., 2014). We find that bycatch, once high and
43 unsustainable for some species (Forney et al., 2021, 2001), is now well below the “zero mortality rate
44 goal” (ZMRG) of 10% of the potential biological removal (50 C.F.R. § 229.2). Recent marine mammal
45 bycatch estimates range from 0.1-4.0% of their potential biological removals and common murre bycatch
46 has been effectively eliminated. All of the evaluated populations, including the once declining and heavily
47 depleted Morro Bay harbor porpoise population, are growing or stable. These advances, while directly
48 attributable to management interventions, are more due to reductions in fishing effort than to reductions in
49 bycatch rates (i.e., bycatch per unit effort). This highlights a steep tradeoff between conservation and
50 fisheries objectives under the current management regime: while populations of protected species have
51 undergone sustained growth, fishing opportunities have undergone sustained declines. Despite this, there
52 have been calls for more bycatch-motivated restrictions to the fishery (Birch et al., 2023; Birch and
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 507 Shester, 2023). Our results indicate that current fishing operations do not pose a threat to the evaluated
5 species, which suggests that current management is sufficient at limiting bycatch. However, targeting
6 management toward spatial-temporal bycatch hotspots could improve economic outcomes while keeping
7
8 509 bycatch low.
9
10

11 511
12 512 Our results suggest that spatial-temporal management could more efficiently and effectively
13 manage bycatch risk than gear modifications or soak time regulations. Specifically, bycatch rates for
14 California sea lions and harbor seals are greatest from April 1 to June 15, suggesting that a 2.5 month
15 seasonal closure of bycatch spots for these two species could reduce bycatch risk. Spatially, our findings
16 indicate that the existing state water closure (0-3 nautical miles offshore) in Southern California could be
17 adjusted to target specific bycatch hotspots around the Channel Islands, where both California sea lions
18 and harbor seals face higher bycatch risk compared to nearshore areas (except for the nearshore strip from
19 Santa Barbara to Point Sal). This adjustment could potentially open up more fishing grounds in the
20 nearshore region. However, we caution that such hotspot closures could exacerbate bycatch problems if
21 fishing effort is displaced and concentrated in areas of secondarily high risk (Free et al., 2023). Therefore,
22 monitoring of fishing effort and bycatch rates are important for verifying that seasonal closures achieve
23 their conservation and fisheries objectives. Additionally, any changes in current management strategies
24 must take into account spatial-temporal patterns of bycatch and relative sensitivity of each species.
25
26 519
27
28 520
29
30 521
31
32 522
33
34 523
35
36 524
37
38 525
39
40 526 The completion and safekeeping of accurate fisheries-dependent data is fundamental to producing
41 accurate bycatch assessments and effective management strategies. While summaries of historical data
42 facilitated reliable bycatch estimates through ratio estimation, the loss of raw observer data from the
43 1980s likely impeded our ability to accurately estimate bycatch in the northern strata using random
44 forests, an approach often thought to be more accurate than ratio estimation (Stock et al., 2019). Notably,
45 the lost data document a period when fishing was allowed in shallower, more inshore, and more northern
46 waters (**Fig. 1A**). Recovering this data would enhance our ability to assess the drivers of bycatch in the
47 northern region and could provide insights for re-evaluating previous management strategies.
48
49 533 Furthermore, missing meta-data on critical gear characteristics (e.g., mesh size, net length, net height, net
50 material; **Fig. S3**) in the available observer data also limited our ability to identify the potential for these
51 management levers to reduce bycatch risk. Ensuring the complete documentation of gear characteristics,
52 perhaps by prioritizing characteristics known to impact bycatch risk in other gillnet fisheries (Northridge
53 et al., 2017), is important to maximizing the utility of expensive, and sometimes controversial, observer
54 programs (Suuronen and Gilman, 2020). Finally, the ability to delineate individual sets in the logbooks
55 and improved documentation of the characteristics of logged sets would enhance future bycatch estimates
56
57 539
58
59 540
60
61
62
63
64
65

1
2
3
4 541 by allowing sets to be the sampling unit and by avoiding assumptions about missing data, respectively.
5
6 542 This could be achieved by redesigning logbooks, training fishers on completing logbooks, expanding
7
8 543 electronic monitoring, and/or demonstrating that better data can actually lead to fewer restrictions.

9 544
10
11 545 Our results highlight the importance of considering multiple modeling approaches when
12
13 546 estimating and evaluating rare bycatch events. Although model-based methods (e.g., random forests) for
14
15 547 estimating bycatch are often preferred to sample-based methods (e.g., ratio estimators) (Stock et al.,
16
17 548 2019), we find complementary value in using both approaches. While ratio estimation generated more
18
19 549 reliable bycatch estimates due to its ability to leverage both raw and summarized data, the random forest
20
21 550 model provided the empirical basis for assessing drivers of bycatch risk. Additionally, agreement between
22
23 551 the two approaches over the past two decades reinforces predictions that recent bycatch levels have been
24
25 552 low for the evaluated species. Furthermore, our results highlight the value of considering multiple sample
26
27 553 balancing approaches when evaluating bycatch using model-based methods, as the specification of the
28
29 554 best performing model varied by species. Recent efforts to estimate bycatch in West Coast fisheries using
30
31 555 random forests have used only a single sample balancing technique (e.g., Carretta, 2023); we encourage
32
33 556 future efforts to compete multiple approaches to optimize estimates of rare bycatch events (More and
34
35 557 Rana, 2017). Finally, we highlight our derivation of model-specific probability thresholds for classifying
36
37 558 logged fishing efforts as having or not having bycatch using Cohen's kappa as being aligned with
38
39 559 recommended best practices for classifying rare events (Freeman and Moisen, 2008; Manel et al., 2001).
40
41 560 Specifically, research shows that classification based on: (1) the default 0.5 probability threshold
42
43 561 underestimates rare events, (2) true skill statistics overestimates rare events; and (3) Cohen's kappa
44
45 562 provides the least biased predictions. This best practice should also be considered in future bycatch
46
47 563 estimation efforts.

48 564
49
50 565 The continued recovery of protected species will require management of stressors besides
51
52 566 fisheries bycatch, some of which may present even larger threats (Avila et al., 2018; Oldach et al., 2022).
53
54 567 For example, in Summer 2023, there were over 1,000 statewide strandings of California sea lions and
55
56 568 other pinnipeds attributed to domoic acid toxicosis resulting from an intense bloom of harmful diatoms in
57
58 569 the *Pseudo-nitzschia* genus (SCCOOS, 2023; Smith et al., 2023). Harmful algal blooms are increasing in
59
60 570 frequency, duration, and intensity on the West Coast (Hallegraeff et al., 2021) as a result of ocean
61
62 571 warming and eutrophication (McKibben et al., 2017) suggesting that, in the long-term, curbing climate
63
64 572 change and nutrient runoff may be the most important actions for stemming mortality for recovering
65
66 573 pinniped populations. Furthermore, harassment and shooting are some of the most commonly observed
67
68 574 sources of mortality and serious injury for California sea lions, harbor seals, and northern elephant seals

1
2
3
4 575 (Carretta, 2023) (**Fig. S17**), suggesting the need for greater outreach and enforcement to prevent these
5 gratuitous forms of mortality. Finally, it is important to understand the bycatch contributions of other
6 fisheries, many of which report higher levels of *observed* bycatch (**Fig. S17**) yet are not as heavily
7 prosecuted as the California set gillnet fishery. Modeling studies similar to this one are needed to
8 determine whether higher apparent bycatch in these fisheries is due to higher observer coverage or higher
9 bycatch rates. However, the sustained recovery of the evaluated populations suggests that total bycatch
10 across all fisheries is low.
11
12
13
14
15
16 582
17
18 583 There will always be tradeoffs between maximizing fishing opportunities and minimizing bycatch
19 of protected species (Samhouri et al., 2021). As a result, managers often seek to implement regulations
20 that maximize fishing outcomes while keeping bycatch below legally defined sustainable reference points
21 (Kirby and Ward, 2014). The identification of such strategies is seldom straightforward and depends on
22 substantial investments in data and scientific enterprises. Notably, they depend on monitoring populations
23 of protected species to support the assessment of their status and levels of allowable incidental take and
24 monitoring bycatch in key fisheries to support assessments of total bycatch, drivers of bycatch, and the
25 effectiveness of past management interventions (Kirby and Ward, 2014; Punt et al., 2021). In the absence
26 of such data, management must often be precautionary to ensure compliance with protected species
27 legislation (Punt et al., 2021). We illustrate the potential return on investment of supporting such
28 scientific enterprises as our results show that past management interventions have been successful at
29 reducing bycatch in the California set gillnet fishery well below target levels, opening the door for more
30 efficient restrictions and negating the need for unnecessary precaution. The continued demonstration that
31 monitoring programs can generate better outcomes for businesses could facilitate increased public support
32 and funding to identify win-win scenarios for fisheries and conservation in more fisheries.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 598 **References**
5
6

- 7 599 Avila, I.C., Kaschner, K., Dormann, C.F., 2018. Current global risks to marine mammals: Taking stock of
8 the threats. *Biol. Conserv.* 221, 44–58. <https://doi.org/10.1016/j.biocon.2018.02.021>
9 600
10 601 Barlow, J., Baird, R.W., Heyning, J.E., Wynne, K., Manville, A.M., Lowry, L.F., Hanan, D., Sease, J.,
11 Burkhanov, V., 1994. A Review of Cetacean and Pinniped Mortality in Coastal Fisheries Along
12 the West Coast of the USA and Canada and the East Coast of the Russian Federation. *Rep. Int.*
13 602
14 603
15 604
16 Whal. Comm.
- 17 605 Birch, C., Blacow-Draeger, A., Geoff Shester, PH.D., Webb, S., Purcell, E., 2023. The Net Consequence:
18 Impacts of Set Gillnets on California Ocean Biodiversity. Zenodo.
19 606
20 607 <https://doi.org/10.5281/ZENODO.7971874>
21
22 608 Birch, C., Shester, G., 2023. Underreporting of Marine Mammal Takes in the California Set Gillnet
23 Fishery Underscores the Need for Observers. *Oceana*.
24 609
25 610 Bjørge, A., Skern-Mauritzen, M., Rossman, M.C., 2013. Estimated bycatch of harbour porpoise
26 (*Phocoena phocoena*) in two coastal gillnet fisheries in Norway, 2006–2008. Mitigation and
27 implications for conservation. *Biol. Conserv.* 161, 164–173.
28 611
29 612
30 613 <https://doi.org/10.1016/j.biocon.2013.03.009>
31
32 614 CA Secretary of State, 1990. Proposition 132. Marine Resources. Initiative Constitutional Amendment.
33
34 615 Cameron, G.A., Forney, K.A., 2000. Preliminary Estimates of Cetacean Mortality in California/Oregon
35 Gillnet Fisheries for 1999 (International Whaling Commission Working Paper No. SC/52/O24).
36
37 616 Cameron, G.A., Forney, K.A., 1999. Preliminary Estimates of Cetacean Mortality in the California
38 Gillnet Fisheries for 1997 and 1998 (No. SC/51/04). Southwest Fisheries Science Center.
39 617
40 618 Capitolo, P.J., McChesney, G.J., Bechaver, C.A., Rhoades, S.J., Shore, A., Carter, H.R., Parker, M.W.,
41 Eigner, L.E., 2012. Breeding population trends of Brandt's and double-crested cormorants, Point
42 Sur to Point Mugu, California, 1979-2011.
43
44 621
45 622 Carretta, J.V., 2023. Sources of human-related injury and mortality for U.S. Pacific West Coast marine
46 mammal stock assessments, 2017-2021. <https://doi.org/10.25923/QWF2-9B97>
47 623
48
49 624 Carretta, J.V., 2002. Preliminary estimates of cetacean mortality in California gillnet fisheries for 2001
50 (No. SC/54/SM12). International Whaling Commission, Scientific Committee, Shimonoseki,
51 625
52 626 Japan.
53
54 627 Carretta, J.V., 2001. Preliminary estimates of cetacean mortality in California gillnet fisheries for 2000
55 (SC/53/SM9). International Whaling Commission, Scientific Committee, London, UK., London,
56 628
57 629 UK.
58
59 630 Carretta, J.V., Chivers, S.J., 2004. Preliminary estimates of marine mammal mortality and biological

1
2
3
4 631 sampling of cetaceans in California gillnet fisheries for 2003 (No. SC/56/SM1).
5

6 632 Carretta, J.V., Chivers, S.J., 2003. Preliminary estimates of marine mammal mortality and biological
7
8 633 sampling of cetaceans in California gillnet fisheries for 2002 (No. SC/55/SM3).

9 634 Carretta, J.V., Enriquez, L., 2012a. Marine mammal and seabird bycatch in California gillnet fisheries in
10
11 635 2010 (Administrative Report No. LJ-12-01). NOAA Southwest Fisheries Science Center, La
12
13 636 Jolla, CA.

14 637 Carretta, J.V., Enriquez, L., 2012b. Marine mammal and seabird bycatch in California gillnet fisheries in
15
16 638 2011 (NOAA Technical Memorandum NMFS No. NOAA-TM-NMFS-SWFSC-500). NOAA
17
18 639 Southwest Fisheries Science Center, La Jolla, CA.

19 640 Carretta, J.V., Enriquez, L., 2009. Marine mammal and seabird bycatch in observed California
20
21 641 commercial fisheries in 2007 (Administrative Report No. LJ-09-01). NOAA Southwest Fisheries
22
23 642 Science Center, La Jolla, CA.

24 643 Carretta, J.V., Enriquez, L., Villafana, C., 2014. Marine mammal, sea turtle, and seabird bycatch in
25
26 644 California gillnet fisheries in 2012 (NOAA Technical Memorandum NMFS No. NOAA-TM-
27
28 645 NMFS-SWFSC-526). NOAA Southwest Fisheries Science Center, La Jolla, CA.

29 646 Carretta, J.V., Oleson, E.M., Forney, K.A., Muto, M.M., Weller, D.W., Lang, A.R., Baker, J., Hanson, B.,
30
31 647 Orr, A.J., Barlow, J., Moore, J.E., Brownell Jr, R.L., 2022. U.S. Pacific marine mammal stock
32
33 648 assessments: 2021 (NOAA Technical Memorandum No. NOAA-TM-NMFS-SWFSC-663).
34
35 649 NOAA Southwest Fisheries Science Center, La Jolla, CA.

36 650 Carter, H.R., 2001. Population Trends of the Common Murre (*Uria aalge californica*), in: Biology and
37
38 651 Conservation of the Common Murre in California, Oregon, Washington, and British Columbia
39
40 652 Volume 1: Natural History and Population Trends.

41 653 CDFW, 2023. Evaluating Bycatch in the California Halibut Set Gill Net Fishery. California Department
42
43 654 of Fish and Wildlife, Sacramento, CA.

44 655 CDFW, 2021. California Wildlife Habitat Relationship System.

46 656 CDFW, 2019. California Pacific Herring Fishery Management Plan.

48 657 CDFW, 2014. Harbor Seals [ds106] GIS Dataset.

49 658 CDFW, 2010. Seabird Colonies: California, 2010.

51 659 CDFW, 2002. Bathymetry Project: 25m bathymetry dataset.

53 660 Cochran, W.G., 1977. Sampling Techniques. John Wiley and Sons.

54 661 Cohen, J., 1968. Weighted kappa: Nominal scale agreement provision for scaled disagreement or partial
55
56 662 credit. Psychol. Bull. 70, 213–220. <https://doi.org/10.1037/h0026256>

58 663 Condylios, S., 2023. priceR: Economics and Pricing Tools.

59 664 Crowder, L.B., Murawski, S.A., 1998. Fisheries Bycatch: Implications for Management. Fisheries 23, 8–

61
62
63
64
65

- 1
2
3
4 665 17. [https://doi.org/10.1577/1548-8446\(1998\)023<0008:FBIFM>2.0.CO;2](https://doi.org/10.1577/1548-8446(1998)023<0008:FBIFM>2.0.CO;2)
5
6 666 Diamond, S.L., Hanan, D.A., 1986. An Estimate of Harbor Porpoise Mortality in California Set Net
7 Fisheries, April 1, 1983 through March 31, 1984 (Administrative Report No. SWR-86-16).
8
9 668 NOAA Southwest Fisheries Science Center, La Jolla, CA.
10
11 669 Fleiss, J.L., Levin, B., Paik, M.C., 2013. Statistical Methods for Rates and Proportions. John Wiley &
12 Sons.
13
14 671 Forney, K.A., Benson, S.R., Cameron, G.A., 2001. Central California gillnet effort and bycatch of
15 sensitive species, 1990-1998, in: Melvin, E., Parrish, J.K. (Eds.), Seabird Bycatch: Trends,
16 Roadblocks, and Solutions. Alaska Sea Grant, University of Alaska Fairbanks, pp. 141–160.
17
18 673 <https://doi.org/10.4027/sbtrs.2001.08>
19
20 674 Forney, K.A., Carretta, J.V., Benson, S.R., 2014. Preliminary estimates of harbor porpoise abundance in
21 Pacific Coast waters of California, Oregon and Washington, 2007-2012.
22
23 675 Forney, K.A., Moore, J.E., Barlow, J., Carretta, J.V., Benson, S.R., 2021. A multidecadal Bayesian trend
24 analysis of harbor porpoise (*Phocoena phocoena*) populations off California relative to past
25 fishery bycatch. *Mar. Mammal Sci.* 37, 546–560. <https://doi.org/10.1111/mms.12764>
26
27 677 Free, C.M., Bellquist, L.F., Forney, K.A., Humberstone, J., Kauer, K., Lee, Q., Liu, O.R., Samhouri, J.F.,
28 Wilson, J.R., Bradley, D., 2023. Static management presents a simple solution to a dynamic
29 fishery and conservation challenge. *Biol. Conserv.* 285, 110249.
30
31 680 <https://doi.org/10.1016/j.biocon.2023.110249>
32
33 681 Freeman, E.A., Moisen, G.G., 2008. A comparison of the performance of threshold criteria for binary
34 classification in terms of predicted prevalence and kappa. *Ecol. Model.* 217, 48–58.
35
36 682 <https://doi.org/10.1016/j.ecolmodel.2008.05.015>
37
38 683 Geijer, C.K.A., Read, A.J., 2013. Mitigation of marine mammal bycatch in U.S. fisheries since 1994.
39
40 684 *Biol. Conserv.* 159, 54–60. <https://doi.org/10.1016/j.biocon.2012.11.009>
41
42 685 Hallegraeff, G.M., Anderson, D.M., Belin, C., Bottein, M.-Y.D., Bresnan, E., Chinain, M., Enevoldsen,
43 H., Iwataki, M., Karlson, B., McKenzie, C.H., Sunesen, I., Pitcher, G.C., Provoost, P.,
44 Richardson, A., Schweibold, L., Tester, P.A., Trainer, V.L., Yñiguez, A.T., Zingone, A., 2021.
45
46 686 Perceived global increase in algal blooms is attributable to intensified monitoring and emerging
47 bloom impacts. *Commun. Earth Environ.* 2, 117. <https://doi.org/10.1038/s43247-021-00178-8>
48
49 687 Hanan, D.A., Diamond, S.L., 1989. Estimates of Sea Lion, Harbor Seal, and Harbor Porpoise Mortalities
50 in California Set Net Fisheries for the 1986-87 Fishing Year.
51
52 688 Hanan, D.A., Diamond, S.L., Scholl, J.P., 1988. Estimates of Sea Lion and Harbor Seal Mortalities in
53 California Set Net Fisheries for 1983, 1984, and 1985.
54
55 689 Hanan, D.A., Diamond, S.L., Scholl, J.P., 1987. An Estimate of Harbor Porpoise Mortalities in California

1
2
3
4 699 Set Net Fisheries, April 1, 1985 through March 31, 1986 (Administrative Report No. SWR 87-5).
5
6 700 NOAA Southwest Fisheries Science Center, La Jolla, CA.

7 701 Hanan, D.A., Diamond, S.L., Scholl, J.P., 1986. An Estimate of Harbor Porpoise Mortality in California
8
9 702 Set Net Fisheries April 1, 1984 through March 31, 1985 (Administrative Report No. SWR 86-
10
11 703 16). NOAA Southwest Fisheries Science Center, La Jolla, CA.

12 704 Hazen, E.L., Scales, K.L., Maxwell, S.M., Briscoe, D.K., Welch, H., Bograd, S.J., Bailey, H., Benson,
13
14 705 S.R., Eguchi, T., Dewar, H., Kohin, S., Costa, D.P., Crowder, L.B., Lewison, R.L., 2018. A
15
16 706 dynamic ocean management tool to reduce bycatch and support sustainable fisheries. *Sci. Adv.* 4,
17
18 707 eaar3001. <https://doi.org/10.1126/sciadv.aar3001>

19 708 Huang, B., Liu, C., Banzon, V., Freeman, E., Graham, G., Hankins, B., Smith, T., Zhang, H.-M., 2021.
20
21 709 Improvements of the Daily Optimum Interpolation Sea Surface Temperature (DOISST) Version
22
23 710 2.1. *J. Clim.* 34, 2923–2939. <https://doi.org/10.1175/JCLI-D-20-0166.1>

24 711 Hvitfeldt, E., 2023. themis: Extra Recipes Steps for Dealing with Unbalanced Data. R package.
25
26 712 ICES, 2007. Report of the Workshop on Discard Raising Procedures (ICES CM No. 2007ACFM:06).
27
28 713 ICES, San Sebastian, Spain.

29 714 Julian, F., 1994. Pinniped and cetacean mortality in California gillnet fisheries: preliminary estimates for
30
31 715 1993 (International Whaling Commission No. SC/46/0). NOAA Southwest Fisheries Science
32
33 716 Center, La Jolla, CA.

34 717 Julian, F., 1993. Pinniped and Cetacean Mortality in California Gillnet Fisheries: Preliminary Estimates
35
36 718 for April 1 To June 30,1992 (Administrative Report No. LJ-93-19). NOAA Southwest Fisheries
37
38 719 Science Center, La Jolla, CA.

39 720 Julian, F., Beeson, M., 1998. Estimates of marine mammal, turtle, and seabird mortality for two
40
41 721 California gillnet fisheries: 1990-1995. *Fish. Bull.* 96, 271–284.

42
43 722 Kirby, D.S., Ward, P., 2014. Standards for the effective management of fisheries bycatch. *Mar. Policy* 44,
44
45 723 419–426. <https://doi.org/10.1016/j.marpol.2013.10.008>

46 724 Konno, E.S., 1990. Effort Estimates of Gill Net Fisheries in California that Incidentally Catch Marine
47
48 725 Mammals, for the 1987-88 Fishing Year. NOAA Southwest Fisheries Science Center, Terminal
49
50 726 Island, CA.

51 727 Laake, J.L., Lowry, M.S., DeLong, R.L., Melin, S.R., Carretta, J.V., 2018. Population growth and status
52
53 728 of California sea lions. *J. Wildl. Manag.* 82, 583–595. <https://doi.org/10.1002/jwmg.21405>

54 729 Landis, J.R., Koch, G.G., 1977. The Measurement of Observer Agreement for Categorical Data.
55
56 730 *Biometrics* 33, 159–174. <https://doi.org/10.2307/2529310>

57
58 731 Lennert, C., Kruse, S., Beeson, M., Barlow, J., 1994. Estimates of incidental marine mammal bycatch in
59
60 732 California gillnet fisheries for July through December, 1990 (Report of the International Whaling

- 1
2
3
4 733 Commission No. SC/43/O 3).
5
6 734 Lewison, R.L., Crowder, L.B., Wallace, B.P., Moore, J.E., Cox, T., Zydelis, R., McDonald, S., DiMatteo,
7 A., Dunn, D.C., Kot, C.Y., Bjorkland, R., Kelez, S., Soykan, C., Stewart, K.R., Sims, M.,
8 Boustany, A., Read, A.J., Halpin, P., Nichols, W.J., Safina, C., 2014. Global patterns of marine
9 mammal, seabird, and sea turtle bycatch reveal taxa-specific and cumulative megafauna hotspots.
10
11 737 Proc. Natl. Acad. Sci. 111, 5271–5276. <https://doi.org/10.1073/pnas.1318960111>
12
13 738 Liaw, A., Wiener, M., 2002. Classification and Regression by randomForest. R News 2, 18–22.
14
15 739 Long, C.A., Ahrens, R.N.M., Jones, T.T., Siders, Z.A., 2024. A machine learning approach for protected
16 species bycatch estimation. Front. Mar. Sci. 11. <https://doi.org/10.3389/fmars.2024.1331292>
17
18 740 Lopez, J., Griffiths, S., Wallace, B.P., Cáceres, V., Rodríguez, L.H., Abrego, M., Alfaro-Shigueto, J.,
19 Andraka, S., Brito, M.J., Bustos, L.C., Cari, I., Carvajal, J.M., Clavijo, L., Cocas, L., Paz, N. de,
20 Herrera, M., Mangel, J.C., Pérez-Huaripata, M., Piedra, R., Dávila, J.A.Q., Rendón, L., Rguez-
21 Baron, J.M., Santana, H., Suárez, J., Veelenturf, C., Vega, R., Zárate, P., 2024. Vulnerability of
22 the Critically Endangered leatherback turtle to fisheries bycatch in the eastern Pacific Ocean. I. A
23 machine-learning species distribution model. Endanger. Species Res. 53, 271–293.
24
25 741 <https://doi.org/10.3354/esr01288>
26
27 742 Lowry, M., Condit, R., Hatfield, B., Allen, S.G., Berger, R., Morris, P.A., Le Boeuf, B.J., Reiter, J., 2014.
28
29 743 Abundance, Distribution, and Population Growth of the Northern Elephant Seal (*Mirounga*
30 *angustirostris*) in the United States from 1991 to 2010. Aquat. Mamm. 40, 20–31.
31
32 744 <https://doi.org/10.1578/AM.40.1.2014.20>
33
34 745 Lowry, M.S., 2021. Abundance and distribution of pinnipeds at the Channel Islands in southern
35 California, central and northern California, and southern Oregon during summer 2016–2019
36 (NOAA Technical Memorandum NMFS No. NOAA-TM-NMFS-SWFSC-656). NOAA
37
38 746 Southwest Fisheries Science Center, La Jolla, CA.
39
40 747 Manel, S., Williams, H.C., Ormerod, S. j., 2001. Evaluating presence–absence models in ecology: the
41 need to account for prevalence. J. Appl. Ecol. 38, 921–931. <https://doi.org/10.1046/j.1365-2664.2001.00647.x>
42
43 748 Martin, S.L., Stohs, S.M., Moore, J.E., 2015. Bayesian inference and assessment for rare-event bycatch in
44 marine fisheries: a drift gillnet fishery case study. Ecol. Appl. 25, 416–429.
45
46 749 <https://doi.org/10.1890/14-0059.1>
47
48 750 McCracken, M.L., 2004. Modeling a Very Rare Event to Estimate Sea Turtle Bycatch: Lessons Learned
49 (NOAA Technical Memorandum No. NMFS-PIFSC-3). Pacific Islands Fisheries Science Center.
50
51 751 McKibben, S.M., Peterson, W., Wood, A.M., Trainer, V.L., Hunter, M., White, A.E., 2017. Climatic
52 regulation of the neurotoxin domoic acid. Proc. Natl. Acad. Sci. 114, 239–244.

- 1
2
3
4 767 https://doi.org/10.1073/pnas.1606798114
5
6 768 More, A.S., Rana, D.P., 2017. Review of random forest classification techniques to resolve data
7 imbalance, in: 2017 1st International Conference on Intelligent Systems and Information
8 Management (ICISIM). Presented at the 2017 1st International Conference on Intelligent Systems
9 and Information Management (ICISIM), IEEE, Aurangabad, pp. 72–78.
10
11 771 https://doi.org/10.1109/ICISIM.2017.8122151
12
13 772 Nembrini, S., König, I.R., Wright, M.N., 2018. The revival of the Gini importance? Bioinformatics 34,
14
15 3711–3718. https://doi.org/10.1093/bioinformatics/bty373
16
17 773 NOAA, 2024. List of Fisheries for 2024, Federal Register.
18
19 774 NOAA, 2023. List of Fisheries for 2024, Federal Register.
20
21 775 Northridge, S., Coram, A., Kingston, A., Crawford, R., 2017. Disentangling the causes of protected-
22 species bycatch in gillnet fisheries. Conserv. Biol. J. Soc. Conserv. Biol. 31, 686–695.
23
24 776 https://doi.org/10.1111/cobi.12741
25
26 777 O'Keefe, C.E., Cadrin, S.X., Glemarec, G., Rouxel, Y., 2023. Efficacy of Time-Area Fishing Restrictions
27 and Gear-Switching as Solutions for Reducing Seabird Bycatch in Gillnet Fisheries. Rev. Fish.
28
29 778 Sci. Aquac. 31, 29–46. https://doi.org/10.1080/23308249.2021.1988051
30
31 779 Oldach, E., Killeen, H., Shukla, P., Brauer, E., Carter, N., Fields, J., Thomsen, A., Cooper, C., Mellinger,
32 L., Wang, K., Hendrickson, C., Neumann, A., Bøving, P.S., Fangue, N., 2022. Managed and
33 unmanaged whale mortality in the California Current Ecosystem. Mar. Policy 140, 105039.
34
35 780 https://doi.org/10.1016/j.marpol.2022.105039
36
37 781 Ortiz, M., Arocha, F., 2004. Alternative error distribution models for standardization of catch rates of
38 non-target species from a pelagic longline fishery: billfish species in the Venezuelan tuna
39 longline fishery. Fish. Res., Models in Fisheries Research: GLMs, GAMS and GLMMs 70, 275–
40
41 782 297. https://doi.org/10.1016/j.fishres.2004.08.028
42
43 783 Perkins, P., Barlow, J., Beeson, M., 1994. Report on Pinniped and Cetacean Mortality in California
44 Gillnet Fisheries: 1988-1990 (Administrative Report No. LJ-94-11). NOAA Southwest Fisheries
45 Science Center, La Jolla, CA.
46
47 784 Perkins, P., Barlow, J., Beeson, M., 1992a. Report on Pinniped and Cetacean Mortality in California
48 Gillnet Fisheries: 1990-1991 (Administrative Report No. LJ-92-14). NOAA Southwest Fisheries
49 Science Center, La Jolla, CA.
50
51 785 Perkins, P., Barlow, J., Beeson, M., 1992b. Pinniped and cetacean mortality in California gillnet fisheries:
52 1991 (No. SC/44/SM14).
53
54 786 Punt, A.E., Siple, M.C., Francis, T.B., Hammond, P.S., Heinemann, D., Long, K.J., Moore, J., Sepúlveda,
55 M., Reeves, R.R., Sigurðsson, G.M., Víkingsson, G., Wade, P.R., Williams, R., Zerbini, A.N.,
56
57 787
58
59 788
60
61
62
63
64
65

- 1
2
3
4 801 2021. Can we manage marine mammal bycatch effectively in low-data environments? *J. Appl.*
5 *Ecol.* 58, 596–607. <https://doi.org/10.1111/1365-2664.13816>
6 802
7 803 Read, A.J., Drinker, P., Northridge, S., 2006. Bycatch of Marine Mammals in U.S. and Global Fisheries.
8 *Conserv. Biol.* 20, 163–169. <https://doi.org/10.1111/j.1523-1739.2006.00338.x>
9 804
10 805 Rochet, M.-J., Trenkel, V.M., 2005. Factors for the variability of discards: assumptions and field
11 evidence. *Can. J. Fish. Aquat. Sci.* 62, 224–235. <https://doi.org/10.1139/f04-185>
12 806
13 807 Samhouri, J.F., Feist, B.E., Fisher, M.C., Liu, O., Woodman, S.M., Abrahms, B., Forney, K.A., Hazen,
14 E.L., Lawson, D., Redfern, J., Saez, L.E., 2021. Marine heatwave challenges solutions to human–
15 wildlife conflict. *Proc. R. Soc. B Biol. Sci.* 288, 20211607.
16 808
17 809 <https://doi.org/10.1098/rspb.2021.1607>
18 810
19 811 SCCOOS, 2023. California HAB Bulletin: May-July 2023 [WWW Document]. URL
20 <https://sccoos.org/california-hab-bulletin/may-2023/> (accessed 5.31.24).
21 812
22 813 Senko, J., White, E.R., Heppell, S.S., Gerber, L.R., 2014. Comparing bycatch mitigation strategies for
23 vulnerable marine megafauna. *Anim. Conserv.* 17, 5–18. <https://doi.org/10.1111/acv.12051>
24 814
25 815 Smith, J., Cram, J.A., Berndt, M.P., Hoard, V., Shultz, D., Deming, A.C., 2023. Quantifying the linkages
26 between California sea lion (*Zalophus californianus*) strandings and particulate domoic acid
27 concentrations at piers across Southern California. *Front. Mar. Sci.* 10, 1278293.
28 816
29 817 <https://doi.org/10.3389/fmars.2023.1278293>
30 818
31 819 Smith, J.A., Tommasi, D., Sweeney, J., Brodie, S., Welch, H., Hazen, E.L., Muhling, B., Stohs, S.M.,
32 Jacox, M.G., 2020. Lost opportunity: Quantifying the dynamic economic impact of time-area
33 fishery closures. *J. Appl. Ecol.* 57, 502–513. <https://doi.org/10.1111/1365-2664.13565>
34 820
35 821 Soykan, C., Moore, J., Zydelis, R., Crowder, L., Safina, C., Lewison, R., 2008. Why study bycatch? An
36 introduction to the Theme Section on fisheries bycatch. *Endanger. Species Res.* 5, 91–102.
37 822
38 823 <https://doi.org/10.3354/esr00175>
39 824
40 825 Stock, B.C., Ward, E.J., Thorson, J.T., Jannot, J.E., Semmens, B.X., 2019. The utility of spatial model-
41 based estimators of unobserved bycatch. *ICES J. Mar. Sci.* 76, 255–267.
42 826
43 827 <https://doi.org/10.1093/icesjms/fsy153>
44 828
45 829 Suuronen, P., Gilman, E., 2020. Monitoring and managing fisheries discards: New technologies and
46 approaches. *Mar. Policy* 116, 103554. <https://doi.org/10.1016/j.marpol.2019.103554>
47 830
48 831 Wright, M.N., Ziegler, A., 2017. ranger: A Fast Implementation of Random Forests for High Dimensional
49 Data in C++ and R. *J. Stat. Softw.* 77, 1–17. <https://doi.org/10.18637/jss.v077.i01>
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6 832 Tables & Figures
7
8





Fig. 1. History of the California $\geq 3.5"$ set gillnet fishery. Panel A shows the spatial history of fishing effort during four regulatory periods. The horizontal lines delineate geographical strata used in the ratio estimation analysis; strata are labeled in the first plot. The thin coastal line marks state waters (less than 3 nautical miles from the coast). Blocks visited by fewer than three vessels during each regulatory period are hidden to maintain confidentiality and comply with the “rule-of-three.” The other panels show time series of fisheries (B) participation, (C) effort, and (D) revenues. Vertical lines mark years in which major regulations, labeled in Panel B, were implemented; these define the regulatory periods used in Panel A. These regulations became operative on April 15, 1987; January 1, 1994; and April 26, 2002. See Fig. S1 and the supplemental methods for details on estimating ex-vessel revenues from the fishery.



31 843

32 844 **Fig. 2.** History of observer coverage in the California $\geq 3.5"$ set gillnet fishery. Panel A shows the bycatch

33 845 of marine mammals, seabirds, and sea turtles recorded by observers from 1983-2017. We focus on species

34 846 with ≥ 50 observations, which are delineated by the horizontal dashed line. Note log-scale on x-axis. Panel

35 847 B shows the number of observed trips (vessel-days) over time. The dark labels show the estimated percent

36 848 of trips that were observed. Panel C shows the number of observed trips across the spatial (region) and

37 849 temporal (quarters) strata considered in the ratio estimation analysis. See Fig. S8 for a map of the spatial

38 850 strata. Quarters are defined as: 1 = JFM (winter), 2 = AMJ (spring), 3 = JAS (summer), and 4 = OND

39 851 (fall).

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60


61

62

63

64

65



852
853 **Fig. 3.** Estimated bycatch in the California $\geq 3.5''$ set gillnet fishery from 1981-2021 predicted by the ratio
854 estimation (bars) and random forest (line) modeling approaches. Average estimated annual bycatch rates
855 for the last 10 years (2012-2021) from the ratio estimator are marked in the top-right corner. Species are
856 listed in order of decreasing recent bycatch rates.





Fig. 4. Importance of the evaluated explanatory variables in the final random forest model for the study species with acceptable model performance. With the bias towards continuous variable corrected, variable importance is measured as the total decrease in node impurities from splitting on the variable averaged over all trees. Explanatory variables are colored based on the category of the variable.



862
863 **Fig. 5.** Marginal effect of the evaluated explanatory variables on bycatch risk as estimated by the best
864 fitting random forest model for the four study species with acceptable model performance. The marginal
865 effect of each variable represents the effect of the variable when the other variables are held at their mean
866 values. The dashed lines indicate 40 and 60 in Depth (fathoms), 8.5 in mesh size (cm), and 3 in shore
867 distance (km).




Fig. 6. Average spatial bycatch risk as estimated by the best fitting random forest model for the four study species with acceptable model performance. The spatial bycatch risk represents the probability of bycatch at a given location under recent (2010-2021) average conditions. Key landmarks for delineating bycatch hotspots are labeled in each panel.




Fig. 7. Estimated abundance of populations of the six study species. The potential biological removal (PBR) indicates the maximum number of animals, not including natural mortalities, that may be removed from a marine mammal stock while allowing the stock to reach or maintain its optimum sustainable population. PBRs are only calculated for marine mammals. Error bars indicate 95% confidence or credible intervals. Years without reported values were filled using linear interpolation. Vertical lines mark years in which major bycatch regulations were implemented. Population estimates are from the following sources: California sea lion (Laake et al., 2018), northern elephant seal (Carretta et al., 2022), harbor seal (Carretta et al., 2022), harbor porpoise (Forney et al., 2021), Brandt's cormorant (Capitolo et al., 2012), and common murre (Carter, 2001). Data from Carretta et al. (2022) were graphically digitized.

14
15
16
17
18
19
20 883
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Historical bycatch estimation studies and their characteristics (the period column indicates month and year as MM/YY).

Study	Seasons	Period	Species	Fishery definition	Stratification scheme	Sample unit
(Barlow et al., 1994)	1983-87	(see papers)	Pinnipeds/cetaceans	(see included papers)	(see included papers)	(see papers)
(Hanan et al., 1988)	1983-85	Apr 1-Mar 31	Sea lion, harbor seals	Set nets for halibut/flounder/sharks	5 regions	Sets
(Diamond and Hanan, 1986)	1983	4/83 - 3/84	Harbor porpoise	$\geq 8.0"$ set nets for halibut/flounder	3 regions	Sets
(Hanan et al., 1986)	1984	4/84 - 3/85	Harbor porpoise	Set nets for halibut/flounder/sharks	3 regions	Sets
(Hanan et al., 1987)	1985	4/85 - 3/86	Harbor porpoise	Set nets (but not for croaker)	3 regions	Sets
(Hanan and Diamond, 1989)	1986	4/86 - 3/87	Sea lion, harbor seal, harbor porpoise	Set nets for halibut/flounder/sharks	5 regions, seasons	Sets
(Konno, 1990)	1987	4/87 - 3/88	Sea lion, harbor seal, harbor porpoise	Set nets for halibut/flounder/sharks	5 regions, seasons	Sets
(Perkins et al., 1994)	1988-90	4/88 - 3/89 4/89 - 12/89 1/90 - 12/90	Pinnipeds/cetaceans	Set nets for halibut/angel shark	3 regions	Vessel-day
(Lennert et al., 1994)	1990	7/90 - 12/90	Marine mammals	Set nets for halibut/angel shark	3 regions	Vessel-day
(Perkins et al., 1992a)	1990	7/90 - 6/91	Pinnipeds/cetaceans	$\geq 8.0"$ nets for halibut/angel shark	3 regions	Sets
(Perkins et al., 1992b)	1991	1/91 - 12/91	Pinnipeds/cetaceans	Set nets for halibut/angel shark	SCal/Ventura: Quarters Other strata: Annual	Vessel-day
(Julian, 1993)	1992	1/92 - 12/92	Pinnipeds/cetaceans	Set nets for halibut/angel shark	4 regions: quarterly	Vessel-day
(Julian, 1994)	1993	1/93 - 12/93	Pinnipeds/cetaceans	Set nets for halibut	4 regions	Vessel-day
(Julian and Beeson, 1998)	1990-95	7/90-12/90 1-12, 91-95	Mammals/seabirds/turtles	$\geq 8.5"$ nets for halibut/angel shark	SCal/Ventura: Quarters Other strata: Annual	Vessel-day
(Cameron and Forney, 1999)	1997-98	1/97 - 12/97 1/98 - 12/98	Cetaceans	$\geq 8.5"$ nets for halibut/angel shark	1997: Geographical+seasonal 1998: Geographical only	Vessel-day
(Cameron and Forney, 2000)	1999	1/99 - 12/99	Cetaceans	$\geq 8.5"$ nets for halibut/angel shark	Geographical+seasonal	Vessel-day
(Carretta, 2001)	2000	1/00 - 12/00	Cetaceans	$\geq 8.5"$ nets for halibut/angel shark	SCal/Ventura: Quarters Other strata: Annual	Vessel-day
(Carretta, 2002)	2001	1/01 - 12/01	Cetaceans	Set nets for halibut/angel shark	SCal/Ventura: Quarters Other strata: Annual	Vessel-day
(Carretta and Chivers, 2003)	2002	1/02 - 12/02	Cetaceans	Set nets for halibut/angel shark	SCal/Ventura: Quarters Other regions: Annual	Vessel-day
(Carretta and Chivers, 2004)	2003	1/03 - 12/03	Marine mammals	$\geq 8.5"$ nets for halibut/angel shark	SCal/Ventura: Quarters Other regions: Annual	Vessel-day
(Carretta and Enriquez, 2009)	2007	1/07 - 12/07	Mammals/seabirds	Set nets for halibut/white seabass	No stratification	Sets
(Carretta and Enriquez, 2012a)	2010	1/10 - 12/10	Mammals/seabirds	Set nets for halibut/white seabass	No stratification	Vessel-day
(Carretta and Enriquez, 2012b)	2011	1/11 - 12/11	Mammals/seabirds	Set nets for halibut/white seabass	No stratification	Sets
(Carretta et al., 2014)	2012	1/12 - 12/12	Mammals/seabirds/turtles	Set nets for halibut/white seabass	No stratification	Vessel-day

14
15
16
17
18
1920 884 **Table 2.** Performance of the best fitting random forest model by species.
21
22
23

Species	# of bycatch observed	Best model	Mtry	Training data		Test data	
				Kappa	AUC	Kappa	AUC
California sea lion	1372	Weighted-25	3	0.24	0.78	0.23	0.78
Harbor seal	519	Weighted-75	2	0.25	0.83	0.15	0.81
Harbor porpoise	97	Weighted-50	2	0.34	0.98	-0.005	0.98
Common murre	2381	Weighted-25	6	0.71	0.99	0.61	0.97
Brandt's cormorant	118	Weighted-25	8	0.06	0.68	0.07	0.63
Northern elephant seal	78	Weighted-25	1	0.23	0.87	0.21	0.86

38 885
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5

6 886 Supplemental Methods
7
8
9


10 887 Identifying set gillnet landing receipts and revenues
11

12 888 We used landing receipts (a.k.a., fish tickets) to estimate ex-vessel revenues generated by the
13 California ≥ 3.5 " set gillnet fishery from 1981-2022. Among other information, landing receipts report the
14 date, value, species, and gear of commercial landings. We identified landing receipts associated with the
15 ≥ 3.5 " set gillnet fishery through a multi-step filtering process. First, we filtered the landing receipts to the
16 five gear types that could include ≥ 3.5 " mesh set gillnets: trammel nets, set gillnets, small-mesh set
17 gillnets, large-mesh set gillnets, or entangling nets (**Fig. S1A**). Entangling nets, which encompass both set
18 and drift gillnets, were a widely used gear type from 1984-1993. As a result, this filter retained many
19 swordfish landings and other landings associated with drift gillnets. It also retained many herring landings
20 and other landings associated with set gillnets with mesh sizes smaller than 3.5 inches. To remove landing
21 receipts associated with drift gillnets and set gillnets with mesh sizes smaller than 3.5 inches, we used
22 gillnet logbooks to identify landing receipts associated with known set gillnet vessels and logged set
23 gillnet trips. We began by further filtering to only include vessels documented as using ≥ 3.5 " set gillnets
24 in the gillnet logbooks (**Fig. S1B**). After this filter, a large amount of swordfish and herring landings
25 remained, indicating that many ≥ 3.5 " set gillnet vessels use other gears. Thus, to further tie landing
26 receipts with known ≥ 3.5 " set gillnet trips, we explored four related approaches for linking landing
27 receipts to logged ≥ 3.5 " set gillnet trips. The first approach was the most strict and only considered
28 landing receipts reported on the exact day of logged ≥ 3.5 " set gillnet trips (**Fig. S1C**). This filter
29 eliminated swordfish and herring landings but is likely to be overly restrictive. The date of landing may
30 differ from the date of fishing because of misreporting, multi-day trips, or delayed sales. Thus, we
31 explored three progressively less restrictive rules, which attributed landing receipts recorded within one
32 (**Fig. S1D**), two (**Fig. S1E**), or three (**Fig. S1F**) days of logged ≥ 3.5 " set gillnet trips to the fishery. We
33 selected the landing receipts associated with the 3-day buffer as the final set of landing receipts associated
34 with the fishery because it effectively eliminated landings of species not associated with the ≥ 3.5 " set
35 gillnet fishery (i.e., swordfish and herring) while being inclusive-within-reason of potential ≥ 3.5 " set
36 gillnet fishery landings. Finally, we adjusted daily ex-vessel landings values for inflation by converting
37 all values to January 1, 2022 US dollars using the *priceR* package in R (Condylios, 2023).
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 914 Mapping species ranges
5
6

7 915 We mapped the range of the study species using range maps from the California Wildlife Habitat
8 916 Relationships (CWHR) System (CDFW, 2021). The CWHR ranges were developed by species-specific
9 experts. Range maps were not developed for harbor porpoise as part of the CWHR effort. We developed a
10 917 range map for harbor porpoise assuming that harbor porpoise occur primarily in waters shallower than 50
11 918 fathoms (92 meters) north of Point Conception (Forney et al., 2014). Harbor seal haulouts were mapped
12 919 using the CDFW Harbor Seal Haulout GIS dataset (CDFW, 2014). CDFW conducted aerial surveys of all
13 920 known haulout sites in 2001, 2002, and 2003 and counted the number of harbor seals observed in aerial
14 921 photographs of each site. We mapped northern elephant seal rookery size in 2010 using a database of
15 922 counts developed by (Lowry et al., 2014). Counts were generated through a review of ground and aerial
16 923 photographic surveys. We mapped California sea lion haulouts using data from (Lowry, 2021). Haulouts
17 924 were mapped in the Channels Islands between 2016-2019 using aerial photographic surveys. Sea lion
18 925 haulouts occur along the California coast but were not mapped to single sites in this study and therefore
19 926 not plotted in our range maps. We mapped seabird colonies using the 2010 CDFW Seabird Colonies
20 927 Database (CDFW, 2010). These data were collected as part of the Marine Life Protection Act (MLPA)
21 928 planning process and report the maximum number of seabirds of 26 species at all known colonies.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 930 Supplemental Tables & Figures
5
6



47
48 932 **Fig. S1.** Estimated ex-vessel revenues generated by the California $\geq 3.5"$ set gillnet fishery as estimated
49 through six different filtration procedures. The filtration procedures examine the sum annual ex-vessel
50 revenues reported on landing receipts from (A) vessels using various reported entangling net gears; (B)
51 vessels using various entangling net gears that are known to use set gillnets based on logbooks; and (C-F)
52 vessels known to use set gillnets based on logbooks that are dated within various buffers of a logged set
53 gillnet trip. We adopted the final filter, which sums landing receipts date within 3 days of logged set
54 gillnet trip, as the best estimate of ex-vessel revenues for the fishery. Revenues have not been adjusted for
55 inflation (see Fig. 1D for the inflation adjusted ex-vessel revenues).
56
57
58
59
60
61
62
63
64
65






Fig. S2. Diagram of a typical California set gillnet illustrating the measurements reported in the logbook and observer data. Grey text indicates parts of the gillnet and black text indicates the reported measurements. The material, strength, and color of the net is also reported. Finally, the hanging ratio, a percentage that is calculated as the length of the mesh web divided by the length of the cork line, is reported.



32 946
33 947 **Fig. S3.** Traits of the observed set gillnet metadata. Panel A shows the level of completeness of gillnet
34 metadata. Panel B shows the distribution of reported depths. Panel C shows the distribution of reported
35 soak times; the maximum reported soak time is 240 hours (10 days). Panel D shows the distribution of
36 reported mesh sizes; the maximum reported mesh size is 26 inches. Panel E shows the distribution of
37 reported mesh sizes by reported target species. In the boxplots, the solid line indicates the median, the box
38 indicates the interquartile range (IQR; 25th to 75th percentiles), the whiskers indicate 1.5 times the IQR,
39 and points indicate outliers.
40
41
42
43
44



32 954
33 955 **Fig. S4.** The (A) number of sets per observed fishing trip by year and overall; (B) number of pseudo-sets
34 956 per logged fishing trip by year and overall; and (C) number of bycatch in sets with bycatch of each of the
35 957 species of interest. In the boxplots, the solid line indicates the median, the box indicates the interquartile
36 958 range (IQR; 25th to 75th percentiles), the whiskers indicate 1.5 times the IQR, and points indicate
37 959 outliers. In (A), the fill color indicates the number of observed fishing trips contributing to the annual
41 960 distribution.

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64




Fig. S5. Traits of the set gillnet sets documented in the logbook data. Panel **A** shows the level of completeness of gillnet logbook metadata. Panel **B** shows the distribution of reported depths. Panel **C** shows the distribution of reported soak times. The maximum reported soak time in the observer data is 10 days; rare values larger than this value were assumed to be unrealistic and were capped at the maximum. Panel **D** shows the distribution of reported mesh sizes by reported target species. In the boxplots, the solid line indicates the median, the box indicates the interquartile range (IQR; 25th to 75th percentiles), the whiskers indicate 1.5 times the IQR, and points indicate outliers. Panel **E** shows the distribution of reported mesh sizes. Panel **F** shows the distribution of reported net lengths.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26 970
27 971
28
29 972
30
31 973
32 974
33 975
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65






Fig. S6. Mesh size (inches) by target species in the logbook (red) and observer (blue) data. Since 1989, California halibut and Pacific angel shark can only be targeted using mesh sizes larger than 8.5 inches. White seabass are typically targeted using a minimum mesh size of 6.0 inches; however, a small amount of incidental take (<20% of catch and ≤ 10 individuals) in mesh sizes between 3.5 to 6.0 inches is allowed from June 16 to March 14 (14, § 155.10).



32 976
33 977 **Fig. S7.** Comparison of key set traits in the logbook (red) and observer (blue) data. We used a two-sided
34 Kolmogorov-Smirnov test to confirm that the traits of the logbook and observer data could have come
35 from the same probability distribution (all p-values < 0.001) could have come from the same probability
36 distribution (all p-values < 0.001).
37
38
39



981
982 **Fig. S8.** The regional stratification scheme used throughout the analysis. The stratification scheme north
983 of Point Conception was originally proposed by Diamond and Hanan (1986). The stratification scheme
984 south of Point Conception was originally proposed by Julian (1993). The dark black lines around the
985 Channel Islands show the 10 km buffer used to identify island-associated fishing trips.
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000






Fig. S9. Annual bycatch rates by species and regional strata as assumed in the ratio estimation analysis. Solid circles indicate years with observer data and open circles indicate years whose bycatch rates are borrowed from the closest year with observer data. See **Figs. 1A and S8** for maps of the regional strata.



990
991 **Fig. S10.** Estimates of (A) annual bycatch from historical studies and (B) the uncertainty of these
992 estimates expressed as the coefficient of variation (CV). In (A), error bars indicate 95% confidence
993 intervals. See **Table 1** for the sources of these estimates. In the boxplots, the solid line indicates the
994 median, the box indicates the interquartile range (IQR; 25th to 75th percentiles), the whiskers indicate 1.5
995 times the IQR, and points indicate outliers.



996
997 **Fig. S11.** A comparison of estimates of annual bycatch from our study (bars) and historical studies (points
998 and lines). Potential reasons for these differences are explored in **Fig. S12**.





Fig. S12. A comparison of historical (A) fishing effort (number of vessel days); (B) observer coverage (number of observed vessel days); and (C) observer records (number of observed bycatch) derived in our analysis (bars) and reported in historical studies (points and lines). These time series represent the key inputs into the ratio estimation analysis and help to explain the differences between the estimates of annual bycatch derived in our study as compared to historical studies (see Fig. S11). See Table 1 for additional details on historical studies.



1007
1008 **Fig. S13.** A comparison of the model performance as measured through cross-validation on the training
44 dataset between (A) balanced random forest modeling approaches and (B) weighted random forest
45 modeling approaches for all candidate bycatch species (i.e., observer records > 50). We evaluated the
46 model performance using Cohen's kappa and area under the receiver operator curve (ROC) and selected
47 the model with the highest Cohen's kappa as the best performing model (labeled in the plot). We
48 excluded Brandt's cormorant in the model prediction as their Cohen's kappa (<0.1) is too low to produce
49 reliable predictions.
50
51
52
53
54




Fig. S14. Illustration of the methods used to select a probability threshold for classifying a logged set as having or not having bycatch. We selected the probability threshold that maximizes Cohen's kappa when applied to the training data as the optimal threshold (labeled in plot). We highlight the performance of this threshold when used on the independent test data to illustrate performance on out-of-sample data.




Fig. S15. Ranges of the six study species and information on haulout and colony size for selected species.

Blue colors indicate year-round ranges and yellow colors (common murre only) indicate winter ranges.

The range maps for all species except harbor porpoise are from the California Wildlife Habitat Relationship System (CDFW, 2021). Harbor porpoises occur in waters less than 50 fathoms (92 meters) deep north of Point Conception (Forney et al., 2014). SFRR indicates the San Francisco-Russian River harbor porpoise stock. Common murre and Brandt's cormorant colony counts are from the California Seabird Colony Database (CDFW, 2010). There are no historical breeding records for common murre south of Point Sur except for at the Prince Island colony in Cuyler Harbor of San Miguel Island. Harbor seal haulout counts are from the CDFW Harbor Seal GIS dataset (CDFW, 2014).




Fig. S16. A comparison of estimated bycatch numbers between ratio estimation and random forest stratified by regions (Fig. S8). The major differences between years are highlighted in gray.




Fig. S17. Average annual observed mortality and serious injury incidents by source on the entire U.S.

West Coast between 2007-2022 (Carretta, 2023).

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Yutian Fang reports financial support was provided by Arnhold UC Santa Barbara-Conservation International Climate Solutions. Christopher M. Free reports financial support was provided by The Nature Conservancy California. If there are other authors, they declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.