

		% of amino acid surface buried in the interface						Note
res	AA	Symmetric dimer		Asymmetric dimer; ADP-bound		Asymmetric dimer; AMPPNP bound		
		4MNE_C	4MNE_B	9RTR_C Activator	9RTR_B Receiver	9RTS_C Activator	9RTS_B Receiver	
446	SER	0	0	0	0	20	0	NtA
447	SER	0	0	30	0	20	0	
448	ASP	0	0	70	30	70	40	
449	ASP	0	0	80	90	80	90	
450	TRP	80	80	100	100	90	110	conserved N lobe residues making intramolecular contacts
451	GLU	0	0	20	30	20	30	
452	ILE	0	0	90	80	110	90	
453	PRO	0	0	60	60	50	50	
456	GLN	0	0	70	90	80	90	
475	LYS	0	20	20	0	0	0	
476	TRP	60	70	100	100	100	100	
477	HIS	80	80	100	90	100	90	
478	GLY	60	80	0	20	0	0	
506	ARG	20	20	40	60	20	60	Asymmetric dimer-specific salt bridge with the Asp448 of the NtA - tested R506A and R506E
507	LYS	20	0	50	80	70	80	
508	THR	70	70	0	0	0	20	
509	ARG	100	100	100	100	100	100	
510	HIS	90	90	90	80	90	80	
511	VAL	60	60	70	80	80	70	
513	ILE	0	0	90	70	60	60	Main chain atoms: interface contacts. Side chain atoms: intramolecular contacts.
515	LEU	80	80	100	100	100	100	
516	PHE	100	100	0	80	50	50	
517	MET	100	90	100	100	100	100	
518	GLY	0	0	0	0	0	110	
530	GLN	80	80	50	70	70	60	
532	CYS	0	0	0	20	0	0	Mutated residue. Also mutated in >20 symmetric structures
562	GLN	90	60	/	/	/	/	
562	ARG	/	/	30	0	20	20	
565	ASP	70	60	0	0	0	0	Symmetric dimer specific - tested D565R
566	TYR	100	100	50	40	50	50	
569	ALA	30	30	0	0	0	50	
570	LYS	50	40	50	30	60	90	Asymmetric specific salt bridge between NtA (Asp448) of Activator and the Lys570 of the Receiver - tested K570A, K570R, K570E
571	SER	0	0	0	0	0	40	
585	HIS	30	40	0	0	0	0	
586	GLU	40	40	0	0	0	0	
587	ASP	0	0	0	20	0	0	
588	LEU	50	40	0	0	0	0	
589	THR	30	30	0	0	0	0	
591	LYS	0	0	80	90	80	70	Side chain makes an intramolecular H-bond with Val511 of the αC-β4 regulatory loop
600	GLU	0	0	0	0	0	20	
715	GLU	30	0	0	0	0	0	