



Oxford Cambridge and RSA

A Level Mathematics A

H240/01 Pure Mathematics

Practice Paper – Set 1

Time allowed: 2 hours

You must have:

- Printed Answer Booklet

You may use:

- a scientific or graphical calculator

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the barcodes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION

- The total mark for this paper is **100**.
- The marks for each question are shown in brackets [].
- **You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **16** pages. The Question Paper consists of **8** pages.

Formulae

A Level Mathematics A (H240)

Arithmetic series

$$S_n = \frac{1}{2}n(a + l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$S_\infty = \frac{a}{1-r} \quad \text{for } |r| < 1$$

Binomial series

$$(a+b)^n = a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_r a^{n-r}b^r + \dots + b^n \quad (n \in \mathbb{N})$$

$$\text{where } {}^nC_r = {}_nC_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots \quad (|x| < 1, n \in \mathbb{R})$$

Differentiation

$f(x)$	$f'(x)$
$\tan kx$	$k \sec^2 kx$
$\sec x$	$\sec x \tan x$
$\cot x$	$-\operatorname{cosec}^2 x$
$\operatorname{cosec} x$	$-\operatorname{cosec} x \cot x$

$$\text{Quotient rule } y = \frac{u}{v}, \quad \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

Differentiation from first principles

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Integration

$$\int \frac{f'(x)}{f(x)} dx = \ln|f(x)| + c$$

$$\int f'(x)(f(x))^n dx = \frac{1}{n+1}(f(x))^{n+1} + c$$

$$\text{Integration by parts } \int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

Small angle approximations

$$\sin \theta \approx \theta, \quad \cos \theta \approx 1 - \frac{1}{2}\theta^2, \quad \tan \theta \approx \theta \quad \text{where } \theta \text{ is measured in radians}$$

Trigonometric identities

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \quad \left(A \pm B \neq \left(k + \frac{1}{2}\right)\pi\right)$$

Numerical methods

Trapezium rule: $\int_a^b y \, dx \approx \frac{1}{2}h\{(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1})\}$, where $h = \frac{b-a}{n}$

The Newton-Raphson iteration for solving $f(x) = 0$: $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$

Probability

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cap B) = P(A)P(B|A) = P(B)P(A|B) \quad \text{or} \quad P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Standard deviation

$$\sqrt{\frac{\sum(x-\bar{x})^2}{n}} = \sqrt{\frac{\sum x^2}{n} - \bar{x}^2} \quad \text{or} \quad \sqrt{\frac{\sum f(x-\bar{x})^2}{\sum f}} = \sqrt{\frac{\sum fx^2}{\sum f} - \bar{x}^2}$$

The binomial distribution

If $X \sim B(n, p)$ then $P(X = x) = \binom{n}{x} p^x (1-p)^{n-x}$, Mean of X is np , Variance of X is $np(1-p)$

Hypothesis test for the mean of a normal distribution

If $X \sim N(\mu, \sigma^2)$ then $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$ and $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$

Percentage points of the normal distribution

If Z has a normal distribution with mean 0 and variance 1 then, for each value of p , the table gives the value of z such that $P(Z \leq z) = p$.

p	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.999	0.9995
z	0.674	1.282	1.645	1.960	2.326	2.576	2.807	3.090	3.291

Kinematics

Motion in a straight line

$$v = u + at$$

$$s = ut + \frac{1}{2}at^2$$

$$s = \frac{1}{2}(u + v)t$$

$$v^2 = u^2 + 2as$$

$$s = vt - \frac{1}{2}at^2$$

Motion in two dimensions

$$\mathbf{v} = \mathbf{u} + \mathbf{a}t$$

$$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$$

$$\mathbf{s} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$$

$$\mathbf{s} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2$$

Answer **all** the questions.

1 In this question you must show detailed reasoning.

Express each of the following in the form $a + b\sqrt{2}$, where a and b are integers.

(i) $\sqrt{3}(\sqrt{12} + \sqrt{54})$ [3]

(ii) $\frac{6}{2 + \sqrt{2}}$ [3]

2 In this question you must show detailed reasoning.

Solve the equation $3 \sin^2 \theta - 2 \cos \theta - 2 = 0$ for $0^\circ \leq \theta \leq 360^\circ$. [5]

3 (i) Express $2x^2 + 4x + 5$ in the form $p(x + q)^2 + r$, where p , q and r are integers. [4]

(ii) State the coordinates of the turning point on the curve $y = 2x^2 + 4x + 5$. [2]

(iii) Given that the equation $2x^2 + 4x + 5 = k$ has no real roots, state the set of possible values of the constant k . [1]

4 A sheet of metal is a square of side 21 cm. Equal squares of side x cm are cut from each corner, and the sheet is then folded to make an open box with vertical sides.

(i) Use calculus to find the value of x that maximises the volume of the box. Justify that the volume is a maximum. [6]

(ii) State an assumption that is needed when answering part (i). [1]

5 The cubic polynomial $f(x)$ is defined by $f(x) = x^3 + 4x^2 - 7x - 10$.

(i) Given that $f(-1) = 0$, express $f(x)$ in a fully factorised form. [3]

(ii) Show the equation $e^{3y} + 4e^{2y} - 7e^y - 10 = 0$ has exactly one real root. State the exact value of this root. [3]

- 6 The population of fish, P , in a lake is recorded at 10 day intervals. The table below shows the data collected, where t is the number of days since the population was first recorded.

t	0	10	20	30	40	50
P	20	24	29	34	42	50

It is proposed the population can be modelled by the equation $P = ab^t$, where a and b are constants.

- (i) Complete the table of values in the Printed Answer Booklet. Plot the final three values of $\log_{10} P$ against t on the axes provided. [1]
 - (ii) By drawing an appropriate straight line on your graph, find the values of a and b . [3]
 - (iii) Use the model to predict the population of fish when $t = 200$. [1]
 - (iv) Explain why this prediction may not be reliable. [1]
- 7 Find $\int (2x+1)\ln x \, dx$. [5]
- 8 The function f is defined as $f(x) = \frac{8}{x+2}$ for $x \geq 0$.
- (i) State the range of f . [1]
 - (ii) Find an expression for $f^{-1}(x)$. [2]
 - (iii) Solve the equation $f(x) = f^{-1}(x)$. [2]

- 9 Fig. 1 shows a garden that is to be designed to include a lawn and a flowerbed.




Fig. 1

The lawn can be modelled using four trapezia, as shown in Fig. 2. Each trapezium has a width of 1.5 m, and the lengths of the parallel sides are 8.0 m, 8.5 m, 8.2 m, 8.4 m and 8.6 m respectively.




Fig. 2

- (i) (a) Use the trapezium rule with 4 strips to estimate the area of the lawn. [2]
- (b) Given that lawn seed costs £0.49 per square metre, estimate the total cost of the lawn seed required. [1]
- (ii) Suggest two limitations of this model. [2]
- (iii) Suggest one possible refinement of this model. [1]

The flowerbed can be modelled as the segment of a circle with radius 3.2 m. Fertiliser costs £0.17 per square metre.

- (iv) Estimate the total cost of fertiliser required to cover the entire area of the flowerbed. [5]

- 10 The first term in an arithmetic series is $(5t + 3)$, where t is a positive integer. The last term is $(17t + 11)$ and the common difference is 4. Show that the sum of the series is divisible by 12 when, and only when, t is odd. [7]

- 11 (i) Find the first three terms in the expansion of $(1 + 2x)^{\frac{1}{2}}$ in ascending powers of x . [3]
 (ii) Obtain an estimate of $\sqrt{3}$ by substituting $x = 0.04$ into your answer to part (i). [3]
 (iii) Explain why using $x = 1$ in the expansion would not give a valid estimate of $\sqrt{3}$. [1]

12 In this question you must show detailed reasoning.

A curve has equation

$$x \sin y + \cos 2y = \frac{5}{2}$$

for $x \geq 0$ and $0 \leq y < 2\pi$.

Determine the exact coordinates of each point on the curve at which the tangent to the curve is parallel to the y -axis. [9]

- 13 The equation $x^3 - x - 2 = 0$ has exactly one real root α .

(i) Use the iterative formula $x_{n+1} = \sqrt[3]{x_n + 2}$ with $x_1 = 1$ to find α correct to 4 significant figures, showing the result of each iteration. [3]

(ii) An alternative iterative formula is $x_{n+1} = F(x_n)$, where $F(x_n) = \frac{x_n + 2}{x_n^2}$. By considering $F'(\alpha)$, explain why this iterative process will not converge to α . [3]

14 A curve is defined by the parametric equations $x = \frac{2t}{1+t}$ and $y = \frac{t^2}{1+t}$, $t \neq -1$.

(i) (a) Show that the curve passes through the origin. [1]

(b) Find the y -coordinate when $x = 1$. [1]

(ii) Show that the area enclosed by the curve, the x -axis and the line $x = 1$ is given by

$$\int_0^1 \frac{2t^2}{(1+t)^3} dt.$$
 [5]

(iii) In this question you must show detailed reasoning.

Hence use an appropriate substitution to find the exact area enclosed by the curve, the x -axis and the line $x = 1$. [6]

END OF QUESTION PAPER

OCR

Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.