



...day June 20XX – Morning/Afternoon

AS Level Further Mathematics A

Y531 Pure Core

SAMPLE MARK SCHEME

Duration: 1 hour 15 minutes

MAXIMUM MARK 60



This document consists of 16 pages

Text Instructions

1. Annotations and abbreviations

Annotation in scoris	Meaning
✓ and *	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining a result or establishing a given result
dep*	Mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
AG	Answer given
awrt	Anything which rounds to
BC	By Calculator
DR	This question included the instruction: In this question you must show detailed reasoning.

2. Subject-specific Marking Instructions for AS Level Further Mathematics A

- a Annotations should be used whenever appropriate during your marking. The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded. For subsequent marking you must make it clear how you have arrived at the mark you have awarded.
- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner. If you are in any doubt whatsoever you should contact your Team Leader.
- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E


Mark for explaining a result or establishing a given result. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep*' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only – differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is not the case please, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner. Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.
- f Unless units are specifically requested, there is no penalty for wrong or missing units as long as the answer is numerically correct and expressed either in SI or in the units of the question. (e.g. lengths will be assumed to be in metres unless in a particular question all the lengths are in km, when this would be assumed to be the unspecified unit.) We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so. When a value is given in the paper only accept an answer correct to at least as many significant figures as the given value. This rule should be applied to each case. When a value is not given in the paper accept any answer that agrees with the correct value to 2 s.f. Follow through should be used so that only one mark is lost for each distinct accuracy error, except for errors due to premature approximation which should be penalised only once in the examination. There is no penalty for using a wrong value for g. E marks will be lost except when results agree to the accuracy required in the question.
- g Rules for replaced work: if a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests; if there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others. NB Follow these maths-specific instructions rather than those in the assessor handbook.
- h For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some papers. This is achieved by withholding one A mark in the question. Marks designated as cao may be awarded as long as there are no other errors. E marks are lost unless, by chance, the given results are established by equivalent working. 'Fresh starts' will not affect an earlier decision about a misread. Note that a miscopy of the candidate's own working is not a misread but an accuracy error.
- i If a calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers (provided, of course, that there is nothing in the wording of the question specifying that analytical methods are required). Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.
- j If in any case the scheme operates with considerable unfairness consult your Team Leader.

Question			Answer	Marks	AO	Guidance	
1			DR $\alpha + \beta = -2, \alpha\beta = 5$ $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta, \alpha^2\beta^2 = (\alpha\beta)^2$ $\Rightarrow p = 6, q = 25$	B1 M1 A1 [3]	1.2 1.1a 1.1	Both used, must be seen	Allow alternative methods e.g. using complex roots or substituting $x = \sqrt{y}$
2	(i)		DR $z_1^* z_2 = (3 - 2i)(-1 - i) = -3 + 2i - 3i + 2i^2$ $= -5 - i$	M1 A1 [2]	1.1 1.1	Find conjugate, then multiply out brackets	Working must be seen
2	(ii)		DR $\frac{z_1 + 2z_2}{z_2} = \frac{3 + 2i - 2 - 2i}{-1 - i} = \frac{1}{-1 - i} \cdot \frac{-1 + i}{-1 + i}$ $= \frac{-1 + i}{2} = -\frac{1}{2} + \frac{1}{2}i$	M1 A1 [2]	1.1 1.1	Multiply by $\frac{-1 + i}{-1 + i}$	Must be seen

Question		Answer	Marks	AO	Guidance	
3	(i)	$\mathbf{AB} = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} -1 & 2 \\ 2 & -1 \end{pmatrix} = \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}$ <p>So $\mathbf{AB} = 3\mathbf{I}$ and $m = 3$</p>	M1 E1 [2]	1.1 1.1	Or BC	OR M1 $\mathbf{A}^{-1} = \frac{1}{1-4} \begin{pmatrix} 1 & -2 \\ -2 & 1 \end{pmatrix}$ $= \frac{1}{-3} \begin{pmatrix} 1 & -2 \\ -2 & 1 \end{pmatrix} = \frac{1}{3} \mathbf{B}$ E1 So $\mathbf{AB} = 3\mathbf{AA}^{-1} = 3\mathbf{I}$ and $m = 3$
3	(ii)	$\mathbf{CD} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}$ <p>so $\mathbf{C}^{-1} = \frac{1}{2}\mathbf{D}$ and $k = \frac{1}{2}$</p>	B1 E1 [2]	1.1 2.2a	BC	OR B1 $\mathbf{C}^{-1} = \begin{pmatrix} -2 & 4 & -1 \\ -2.5 & 4.5 & -0.5 \\ 1.5 & -2.5 & 0.5 \end{pmatrix}$ BC E1 so $\mathbf{C}^{-1} = \frac{1}{2}\mathbf{D}$ and $k = \frac{1}{2}$
3	(iii)	$\mathbf{EF} = \begin{pmatrix} k & k^2 \\ 3 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ k \end{pmatrix} = \begin{pmatrix} 2k + k^3 \\ 6 \end{pmatrix}$ $\Rightarrow \begin{pmatrix} 2k + k^3 \\ 6 \end{pmatrix} = \begin{pmatrix} -2k \\ 6 \end{pmatrix}$ $\Rightarrow 2k + k^3 = -2k$ $k = 0, 2i, -2i$ $\mathbf{F} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}, \begin{pmatrix} 2 \\ 2i \end{pmatrix} \text{ or } \begin{pmatrix} 2 \\ -2i \end{pmatrix}$	M1 M1 A1 A1 A1 [5]	1.1a 2.1 2.2a 1.1 1.1	Both multiplications attempted BC www If A0 , allow SC1 for one matrix only	

Question	Answer	Marks	AO	Guidance	
4	 <p>The shaded region is the required locus</p>	<p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>[4]</p>	<p>1.1</p> <p>1.1</p> <p>1.1</p> <p>1.1</p>	<p>A circle with centre $3 + 4i$, passing through origin.</p> <p>A line parallel to y-axis through 1</p> <p>Correct shading and continuous lines</p>	<p>Accept the correct minor arc alone</p> <p>Accept the correct line segment alone</p> <p>Must state which region is the required locus.</p>

Question		Answer	Marks	AO	Guidance	
5	(i)	A' is $\left(-\frac{3}{5}, \frac{4}{5}\right)$, B' is $\left(\frac{1}{5}, \frac{7}{5}\right)$ and C' is $\left(\frac{4}{5}, \frac{3}{5}\right)$	B1 B1 B1 [3]	2.2a 1.1 1.1	Coordinates of any two images seen Coordinates of third image seen; completely correct labelled diagram, must include indication of coordinates	
5	(ii)	$M \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}$ So $-\frac{3}{5}x + \frac{4}{5}y = x$ And $\frac{4}{5}x + \frac{3}{5}y = y$ Both of which lead to $y = 2x$	M1 M1 E1 [3]	1.1 1.1 2.2a	Seen or implied At least one seen Must conclude from both equations	OR M1 $A' \left(\frac{-3}{5}, \frac{4}{5}\right) A (1, 0)$ M1 \Rightarrow midpoint is $\left(\frac{1}{5}, \frac{2}{5}\right)$ E1 $\Rightarrow y = 2x$
5	(iii) (a)	Det $M = -1$	B1 [1]	1.1		
5	(iii) (b)	It means that area remains the same ... but that the orientation of the image has changed oe	B1 B1 [2]	2.2a 2.4	Accept 'sense', 'order of labelling' oe	

Question		Answer	Marks	AO	Guidance	
6	(i)	Let x, y and z be the amount invested in each account $\begin{pmatrix} 1 & 1 & 1 \\ 0.025 & 0.037 & 0.049 \\ 1 & -2 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2000 \\ 92 \\ 0 \end{pmatrix}$	B2 [2]	1.1 3.1b	B1 for writing down matrix alone	OR B2 Let s be the amount invested in the savings account $(0.025 \ 0.037 \ 0.049) \begin{pmatrix} 2s \\ s \\ 2000-3s \end{pmatrix} = 92$
6	(ii)	$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 0.025 & 0.037 & 0.049 \\ 1 & -2 & 0 \end{pmatrix}^{-1} \begin{pmatrix} 2000 \\ 92 \\ 0 \end{pmatrix}$ $x = 200, y = 100, z = 1700$ so he invests £200 in the current account £100 in the savings account £1700 in the supersaver account	M1 A1 [2]	3.1a 1.1	Attempt to find x, y, z BC Interpret in context	M1 by multiplying out and solving for s . A1 $s = 100$ so he invests £200 in the current account £100 in the savings account £1700 in the supersaver account
6	(iii)	The 92 in part (ii) should be 92 ± 0.5 , therefore giving a range of answers for each account	E1 E1 [2]	3.1b 3.2b		

Y531

Mark Scheme

June 20XX

Question			Answer	Marks	AO	Guidance	
7			$f(5) = 0$	M1	3.1a	DR	
			$\Rightarrow f(x) = (x-5)(x^2 + bx + c)$	A1	2.4		
			$\Rightarrow f(x) = (x-5)(x^2 - 8x + 25)$	B1	1.1		
			Attempt to solve their quadratic	M1	2.1		
			$x = 4 \pm 3i$	A1	1.1		
			The roots are 5, $4+3i$ and $4-3i$				
			Attempt to find moduli	M1	3.1a		
			$ 5 = 5$				
			$ 4 + 3i = \sqrt{4^2 + 3^2} = 5$	A1	1.1		
			$ 4 - 3i = \sqrt{4^2 + 3^2} = 5$	A1	1.1		
			Distance from origin for all roots is 5 units, so all roots lie on $ z = k$, where $k=5$	E1	2.4	FT	
				[9]			

Question			Answer	Marks	AO	Guidance	
8			Let $n = 4$, then	B1	2.1	Basis case for proof by induction	Must state that $r + 1 > 2$ oe A <i>formal</i> proof is required for full marks Accept other <i>complete</i> methods
			$4! = 24$ and $2^4 = 16$ so $4! > 2^4$	M1	2.1	Assumption	
			Assume true for $n = r$				
			$r! > 2^r$ for $r \geq 4$	M1	1.1	Add next statement	
			Then for $n = r + 1$				
			$(r + 1)! = (r + 1) \times r! > (r + 1) \times 2^r$ by assumption				
			Since $r + 1 > 2$, $(r + 1) \times 2^r > 2 \times 2^r = 2^{r+1}$	E1	2.2a	Sufficient working to establish true for $r + 1$	
			so $(r + 1)! > 2^{r+1}$	E1	2.4	Clear conclusion for induction process	
			If true for r then true for $r + 1$.				
			Hence, given basis case, the statement is true for all positive integers.				
				[5]			
9	(i)		$-1 \times -2 + 2 \times 3 + 1 \times k = 0$	M1	1.1	Attempt the scalar product and set equal to zero so	Allow use of i, j, k notation
			$\Rightarrow k = -4$	A1	1.1		
				[2]			
9	(ii)		Equate x and y coordinates:	M1	3.1a	Use coordinates to find μ and λ .	
			$3 + \lambda = 6 + 2\mu \Rightarrow \lambda - 2\mu = 3$				
			$2 - \lambda = 5 + \mu \Rightarrow \lambda + \mu = -3$				
			$\Rightarrow \mu = -2, \lambda = -1$	A1	1.1		
			Consistent with z coordinates since	E1	3.1a	Check consistency with third coordinate	
			$7 + 3 \times (-1) = 4$ and $2 - (-2) = 4$				
			So the point of intersection is $(2, 3, 4)$	A1	1.1		
				[4]			

Y531

Mark Scheme

June 20XX

Question		Answer	Marks	AO	Guidance	
9	(iii)	<p>The vector product find a mutual perpendicular</p> $\begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix} \times \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} = \begin{pmatrix} -2 \\ 7 \\ 3 \end{pmatrix}$ $\begin{pmatrix} 1 \\ a \\ b \end{pmatrix} = \lambda \begin{pmatrix} -2 \\ 7 \\ 3 \end{pmatrix}$ $\lambda = -\frac{1}{2}$ $a = -3.5, b = -1.5$	<p>M1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[5]</p>	<p>3.1a</p> <p>1.1</p> <p>3.1a</p> <p>1.1</p> <p>1.1</p>	<p>Attempt the vector product, by any valid method</p> <p>BC</p> 	<p>OR M1A1</p> $\begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ a \\ b \end{pmatrix} = 0 \text{ and } \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ a \\ b \end{pmatrix} = 0$ <p>A1 $1 - a + 3b = 0$ and $2 + a - b = 0$</p> <p>M1 Solve simultaneous equations</p> <p>A1 $a = -3.5, b = -1.5$</p>

Assessment Objectives (AO) Grid

Question	AO1	AO2	AO3(PS)	AO3(M)	Total
1	3				3
2(i)	2				2
2(ii)	2				2
3(i)	2				2
3(ii)	1	1			2
3(iii)	3	2			5
4	4				4
5(i)	2	1			3
5(ii)	2	1			3
5(iii)(a)	1				1
5(iii)(b)		2			2
6(i)	1		1		2
6(ii)	1		1		2
6(iii)			2		2
7	4	3	2		9
8	1	4			5
9(i)	2				2
9(ii)	2		2		4
9(iii)	4	1			5
Totals	37	15	8	0	60

PS = Problem Solving

M = Modelling

Summary of Updates

Date	Version	Change
October 2019	2	Amendments to the front cover rubric instructions to candidates

BLANK PAGE

BLANK PAGE

BLANK PAGE

AS Level Further Mathematics A

Y531 Pure Core

Printed Answer Booklet

Version 2

Date – Morning/Afternoon

Time allowed: 1 hour 15 minutes

You must have:

- Question Paper Y531 (inserted)
- Formulae AS Level Further Mathematics A

You may have:

- a scientific or graphical calculator



First name

Last name

Centre
numberCandidate
number**INSTRUCTIONS**

- The Question Paper will be found inside the Printed Answer Booklet.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION


- **You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **12** pages. The Question Paper consists of **4** pages.

1	
2(i)	
2(ii)	

3(i)	
3(ii)	
3(iii)	
(answer space continued on next page)	

3(iii) (continued)	
4	

5(i)



5(ii)

5(iii)(a)	
5(iii)(b)	

PLEASE DO NOT WRITE IN THIS SPACE

6(i)	
6(ii)	
6(iii)	

7	

8

9(i)	
9(ii)	

9(iii)

PLEASE DO NOT WRITE ON THIS PAGE

Copyright Information:

OCR is committed to seeking permission to reproduce all third-party content that it uses in the assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.