



Oxford Cambridge and RSA

AS Level Further Mathematics A

Y533/01 Mechanics

Practice Paper – Set 1

Time allowed: 1 hour 15 minutes

You must have:

- Printed Answer Booklet
- Formulae AS Level Further Mathematics A

You may use:

- a scientific or graphical calculator

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Booklet. The question number(s) must be clearly shown.
- Do **not** write in the barcodes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION

- The total mark for this paper is **60**.
- The marks for each question are shown in brackets [].
- **You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **12** pages. The Question Paper consists of **4** pages.

Answer **all** the questions.

- 1 A climber of mass 65 kg climbs from the bottom to the top of a vertical cliff which is 78 m in height. The climb takes 90 minutes so the velocity of the climber can be neglected.


(i) Calculate the work done by the climber in climbing the cliff. [2]

(ii) Calculate the average power generated by the climber in climbing the cliff. [2]

- 2 The universal law of gravitation states that $F = \frac{Gm_1m_2}{r^2}$ where F is the magnitude of the force between two objects of masses m_1 and m_2 which are a distance r apart and G is a constant.

Find the dimensions of G . [4]

3




A particle P of mass 1.5 kg is attached to one end of a light inextensible string of length 2.4 m. The other end of the string is attached to a fixed point O . The particle is initially at rest directly below O . A horizontal impulse of magnitude 9.3 N s is applied to P . In the subsequent motion the string remains taut and makes an angle of θ radians with the downwards vertical at O , as shown in the diagram.

(i) Find the speed of P when $\theta = \frac{1}{6}\pi$. [6]

(ii) Determine whether P will reach the same horizontal level as O . [2]

4



A and *B* are two long straight parallel horizontal sections of railway track. An engine on track *A* is attached to a carriage of mass 6000 kg on track *B* by a light inextensible chain which remains horizontal and taut in the ensuing motion. The chain is 13 m in length and the points of attachment on the engine and carriage are a perpendicular distance of 5 m apart. The engine and carriage start at rest and then the engine accelerates uniformly to a speed of 5.6 m s^{-1} while travelling 250 m. It is assumed that any resistance to motion can be ignored.

(i) Find the work done on the carriage by the tension in the chain. [3]

(ii) Find the magnitude of the tension in the chain. [3]

The mass of the engine is 10 000 kg.

(iii) At a point further along the track the engine and the carriage are moving at a speed of 8.4 m s^{-1} and the power of the engine is 68 kW. Find the acceleration of the engine at this instant. [3]

5 Two discs, *A* and *B*, have masses 1.4 kg and 2.1 kg respectively. They are sliding towards each other in the same straight line across a large sheet of horizontal ice. Immediately before the collision *A* has speed 2 m s^{-1} and *B* has speed 3 m s^{-1} . Immediately after the collision *A*'s speed is 4 m s^{-1} .

(i) Explain why it is impossible for *A* to be travelling in the same direction after the collision as it was before the collision. [2]

(ii) Find the velocity of *B* immediately after the collision. [4]


(iii) Calculate the coefficient of restitution between *A* and *B*. [2]

(iv) State what your answer to part (iii) means about the kinetic energy of the system. [1]

The discs are made from the same material. The discs will be damaged if subjected to an impulse of magnitude greater than 6.5 N s.

(v) Determine whether *B* will be damaged as a result of the collision. [3]

(vi) Explain why *A* will be damaged if, and only if, *B* is damaged. [1]



A particle of mass 0.2 kg is attached to one end of a light inextensible string. The other end of the string is attached to a fixed point O which is 1.8 m above a smooth horizontal table. The particle moves on the table in a circular path at constant speed with the string taut (see diagram).

The particle has a speed of 0.5 m s^{-1} and its angular velocity is 0.625 rad s^{-1} .

- (i) Show that the radius of the circular path is 0.8 m . [1]
- (ii) Find the magnitude of the normal contact force between the particle and the table. [6]

The speed is changed to $v \text{ m s}^{-1}$. At this speed the particle is just about to lose contact with the table.

- (iii) Find the value of v . [3]

- 7 The masses of two particles A and B are m and $2m$ respectively. They are moving towards each other on a smooth horizontal table. Just before they collide their speeds are u and $2u$ respectively. After the collision the kinetic energy of A is 8 times the kinetic energy of B . Find the coefficient of restitution between A and B . [12]

END OF QUESTION PAPER

OCR

Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.