



Oxford Cambridge and RSA

AS Level Further Mathematics A

Y531/01 Pure Core

Practice Paper – Set 1

Time allowed: 1 hour 15 minutes

You must have:

- Printed Answer Booklet
- Formulae Further Mathematics A

You may use:

- a scientific or graphical calculator

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** If additional space is required, you should use the lined page(s) at the end of this booklet. The question number(s) must be clearly shown.
- Do **not** write in the barcodes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

INFORMATION

- The total mark for this paper is **60**.
- The marks for each question are shown in brackets [].
- **You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **12** pages. The Question Paper consists of **4** pages.

Answer **all** the questions.

1 The matrix **A** is given by $\mathbf{A} = \begin{pmatrix} -3 & 3 & 2 \\ 5 & -4 & -3 \\ -1 & 1 & 1 \end{pmatrix}$.

(i) Find \mathbf{A}^{-1} . [1]

(ii) Solve the simultaneous equations

$$-3x + 3y + 2z = 12a$$

$$5x - 4y - 3z = -6$$

$$-x + y + z = 7$$

giving your solution in terms of a . [3]

2 The loci C_1 and C_2 are given by $|z - (3 + 2i)| = 2$ and $\arg(z - (3 + 2i)) = \frac{5\pi}{6}$ respectively.

(i) Sketch C_1 and C_2 on a single Argand diagram. [4]

(ii) Find, in surd form, the number represented by the point of intersection of C_1 and C_2 . [3]

(iii) Indicate, by shading, the region of the Argand diagram for which

$$|z - (3 + 2i)| \leq 2 \text{ and } \frac{5\pi}{6} \leq \arg(z - (3 + 2i)) \leq \pi. \quad [2]$$

3 Two lines, l_1 and l_2 , have the following equations.

$$l_1: \mathbf{r} = \begin{pmatrix} -11 \\ 10 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$$

$$l_2: \mathbf{r} = \begin{pmatrix} 5 \\ 2 \\ 4 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ 1 \\ -2 \end{pmatrix}$$

P is the point of intersection of l_1 and l_2 .

(i) Find the position vector of P . [3]

(ii) Find, correct to 1 decimal place, the acute angle between l_1 and l_2 . [3]

Q is a point on l_1 which is 12 metres away from P . R is the point on l_2 such that QR is perpendicular to l_1 .

(iii) Determine the length QR . [2]

4 In this question you must show detailed reasoning.

The distinct numbers ω_1 and ω_2 both satisfy the quadratic equation $4x^2 + 4x + 17 = 0$.

(i) Write down the value of $\omega_1\omega_2$. [1]

(ii) A , B and C are the points on an Argand diagram which represent ω_1 , ω_2 and $\omega_1\omega_2$. Find the area of triangle ABC . [6]

5 In this question you must show detailed reasoning.

The equation $x^3 + 3x^2 - 2x + 4 = 0$ has roots α , β and γ .

(i) Using the identity $\alpha^3 + \beta^3 + \gamma^3 \equiv (\alpha + \beta + \gamma)^3 - 3(\alpha\beta + \beta\gamma + \gamma\alpha)(\alpha + \beta + \gamma) + 3\alpha\beta\gamma$ find the value of $\alpha^3 + \beta^3 + \gamma^3$. [3]

(ii) Given that $\alpha^3\beta^3 + \beta^3\gamma^3 + \gamma^3\alpha^3 = 112$ find a cubic equation whose roots are α^3 , β^3 and γ^3 . [4]

6 Prove by induction that $n! \geq 6n$ for $n \geq 4$. [5]

7 A transformation is equivalent to a shear parallel to the x -axis followed by a shear parallel to the y -axis and is represented by the matrix $\begin{pmatrix} 1 & s \\ t & 0 \end{pmatrix}$.

Find in terms of s the matrices which represent each of the shears. [7]

8 (i) (a) Find, in terms of x , a vector which is perpendicular to the vectors $\begin{pmatrix} x-2 \\ 5 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} x \\ 6 \\ 2 \end{pmatrix}$. [2]

(b) Find the shortest possible vector of the form $\begin{pmatrix} 1 \\ a \\ b \end{pmatrix}$ which is perpendicular to the vectors

$\begin{pmatrix} x-2 \\ 5 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} x \\ 6 \\ 2 \end{pmatrix}$. [5]

(ii) Vector \mathbf{v} is perpendicular to both $\begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ p \\ p^2 \end{pmatrix}$ where p is a real number. Show that it is impossible for \mathbf{v} to be perpendicular to the vector $\begin{pmatrix} 1 \\ 1 \\ p-1 \end{pmatrix}$. [6]

END OF QUESTION PAPER

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.