



Oxford Cambridge and RSA

AS Level Further Mathematics A

Y533/01 Mechanics

Practice Paper – Set 2

Time allowed: 1 hour 15 minutes

You must have:

- Printed Answer Booklet
- Formulae AS Level Further Mathematics A

You may use:

- a scientific or graphical calculator

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Booklet. The question number(s) must be clearly shown.
- Do **not** write in the barcodes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION

- The total mark for this paper is **60**.
- The marks for each question are shown in brackets [].
- **You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **12** pages. The Question Paper consists of **4** pages.

Answer **all** the questions.

- 1 A particle P of mass 2.4 kg is attached to one end of a light inextensible string of length 1.4 m . The other end of the string is attached to a fixed point O on a smooth horizontal table. P moves on the table at constant speed along a circular path with O at its centre. The magnitude of the tension in the string is 21 N .

(i) (a) Find the magnitude of the acceleration of P . [2]

(b) State the direction of the acceleration of P . [1]

(ii) Find the speed of P . [2]

(iii) Find the time taken for P to complete a single revolution. [2]

- 2 A pump is pumping still water from the base of a well at a constant rate of 300 kg per minute. The well is 4.5 m deep and water is released from the pump at ground level in a horizontal jet with a speed of 6.2 m s^{-1} .

Ignoring any energy losses due to resistance, calculate the power generated by the pump. [5]

- 3 A student is investigating fluid flowing through a pipe.

In her first model she assumes a relationship of the form $P = S\rho^\alpha g^\beta h^\gamma$ where ρ is the density of the fluid, h is the length of the pipe, P is the pressure difference between the ends of the pipe, g is the acceleration due to gravity and S is a dimensionless constant. You are given that ρ is measured in kg m^{-3} .

(i) Use the fact that pressure is force per unit area to show that $[P] = \text{ML}^{-1}\text{T}^{-2}$. [2]

(ii) Find the values of α , β and γ . [6]

The density of the fluid the student is using is 540 kg m^{-3} . In her experiment she finds that when the length of the pipe is 1.40 m the pressure difference between the ends of the pipe is 3.25 N m^{-2} .

(iii) Find the length of the pipe for which her first model would predict a pressure difference between the ends of the pipe of 4.65 N m^{-2} . [2]

In an alternative model the student suggests a modified relationship of the form $P = S\rho^\alpha g^\beta h^\gamma + \frac{1}{2}h\nu^2$, where ν is the average velocity of the fluid in the pipe.

(iv) Use dimensional analysis to assess the validity of her alternative model. [2]

- 4 A car has a mass of 850 kg and its engine can generate a maximum power of 35 kW. The total resistance to motion of the car is modelled as kv N where $v \text{ m s}^{-1}$ is the speed of the car and k is a constant.

When the car is moving in a straight line on a straight horizontal road, the greatest constant speed that it can attain is 25 m s^{-1} .

(i) Show that $k = 56$. [2]

(ii) Find the greatest possible acceleration of the car on the road at an instant when it is moving with a speed of 15 m s^{-1} . [3]

A trailer of mass 240 kg is attached to the car by means of a light inextensible tow bar which is parallel to the surface of the road. The resistance to motion of the trailer is modelled as a constant force of magnitude 350 N.

The car and trailer move on the horizontal road. At a certain instant the car's engine is working at a rate of 30 kW and the acceleration of the car is 0.2 m s^{-2} .


(iii) (a) Find the speed of the car at this instant. [4]

(b) Find the magnitude of the tension in the tow bar at this instant. [2]

The car and trailer now move in a straight line on a straight road inclined at 8° to the horizontal.

(iv) Find the difference between their greatest possible constant speed travelling up the slope and their greatest possible constant speed travelling down the slope. [5]

- 5 Two particles A and B are on a smooth horizontal floor with B between A and a vertical wall. The masses of A and B are 4 kg and 11 kg respectively. Initially, B is at rest and A is moving towards B with a speed of $u \text{ m s}^{-1}$ (see diagram). A collides directly with B . The coefficient of restitution between A and B is e .




(i) Show that immediately after the collision the speed of B is $\frac{4}{15}u(1+e)$. [4]

After the collision between A and B the direction of motion of A is reversed. B subsequently collides directly with the vertical wall. The coefficient of restitution between B and the wall is $\frac{1}{2}e$.

(ii) Given that there is a second collision between A and B , find the range of possible values of e . [7]

- 6 A fairground game involves a player kicking a ball, B , from rest so as to project it with a horizontal velocity of magnitude $u \text{ m s}^{-1}$. The ball is attached to one end of a light rod of length $l \text{ m}$. The other end of the rod is smoothly hinged at a fixed point O so that B can only move in the vertical plane which contains O , a fixed barrier and a bell which is fixed $l \text{ m}$ vertically above O .

Initially B is vertically below O . The barrier is positioned so that when B collides directly with the barrier, OB makes an angle θ with the downwards vertical through O (see diagram).



The coefficient of restitution between B and the barrier is e . B rebounds from the barrier, passes through its original position and continues on a circular path towards the bell. The bell will only ring if the ball strikes it with a speed of at least $V \text{ m s}^{-1}$. The player wins the game if the player causes the bell to ring having kicked B so that it first collides with the barrier. You may assume that B and the bell are small and that the barrier has negligible thickness.

Show that, whatever the position of the barrier, the player cannot win the game if $u^2 < 4gl + \frac{V^2}{e^2}$. [9]

END OF QUESTION PAPER

OCR

Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.