

A Level Further Mathematics A**Y540 Pure Core 1****Sample Question Paper**

Version 2

Date – Morning/Afternoon

Time allowed: 1 hour 30 minutes

**You must have:**

- Printed Answer Booklet
- Formulae A Level Further Mathematics A

You may use:

- a scientific or graphical calculator

**INSTRUCTIONS**

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION

- The total number of marks for this paper is **75**.
- The marks for each question are shown in brackets [].
- **You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **12** pages. The Question Paper consists of **8** pages.

Answer **all** the questions.

1 Show that $\frac{5}{2-4i} = \frac{1}{2} + i$. [2]

2 In this question you must show detailed reasoning.


The equation $f(x) = 0$, where $f(x) = x^4 + 2x^3 + 2x^2 + 26x + 169$, has a root $x = 2 + 3i$.

(i) Express $f(x)$ as a product of two quadratic factors. [4]

(ii) Hence write down all the roots of the equation $f(x) = 0$. [1]

3 In this question you must show detailed reasoning.

The diagram below shows the curve $r = 2\cos 4\theta$ for $-k\pi \leq \theta \leq k\pi$ where k is a constant to be determined.



Calculate the exact area enclosed by the curve. [6]

4 Draw the region in an Argand diagram for which $|z| \leq 2$ and $|z| > |z - 3i|$. [3]

5 (i) Show that $\frac{d}{dx}(\sinh^{-1}(2x)) = \frac{2}{\sqrt{4x^2 + 1}}$. [2]

(ii) Find $\int \frac{1}{\sqrt{2-2x+x^2}} dx$. [3]

6 The equation $x^3 + 2x^2 + x + 3 = 0$ has roots α , β and γ .

The equation $x^3 + px^2 + qx + r = 0$ has roots $\alpha\beta$, $\beta\gamma$ and $\gamma\alpha$.

Find the values of p , q and r . [5]

7 The lines l_1 and l_2 have equations $\frac{x-3}{1} = \frac{y-5}{2} = \frac{z+2}{-3}$ and $\frac{x-4}{2} = \frac{y+2}{-1} = \frac{z-7}{4}$.

(i) Find the shortest distance between l_1 and l_2 . [5]

(ii) Find a cartesian equation of the plane which contains l_1 and is parallel to l_2 . [2]

8 (i) Find the solution to the following simultaneous equations.

$$\begin{array}{rrcr} x + & y + & z = & 3 \\ 2x + & 4y + & 5z = & 9 \\ 7x + & 11y + & 12z = & 20 \end{array}$$

[2]

(ii) Determine the values of p and k for which there are an infinity of solutions to the following simultaneous equations.

$$\begin{array}{rrcr} x + & y + & z = & 3 \\ 2x + & 4y + & 5z = & 9 \\ 7x + & 11y + & pz = & k \end{array}$$

[6]


9 Prove by induction that, for all positive integers n ,

$$\sum_{r=1}^n \frac{5-4r}{5^r} = \frac{n}{5^n}.$$

[5]

- 10** The Argand diagram below shows the origin O and pentagon $ABCDE$, where A, B, C, D and E are the points that represent the complex numbers a, b, c, d and e , and where a is a positive real number.

You are given that these five complex numbers are the roots of the equation $z^5 - a^5 = 0$.



- (i) Justify each of the following statements.

- (a) A, B, C, D and E lie on a circle with centre O . [1]
- (b) $ABCDE$ is a regular pentagon. [2]
- (c) $b \times e^{\frac{2i\pi}{5}} = c$ [1]
- (d) $b^* = e$ [1]
- (e) $a + b + c + d + e = 0$ [2]

- (ii) The midpoints of sides AB, BC, CD, DE and EA represent the complex numbers p, q, r, s and t . Determine a polynomial equation, with real coefficients, that has roots p, q, r, s and t . [3]

- 11** A company is required to weigh any goods before exporting them overseas. When a crate is placed on a set of weighing scales, the mass displayed takes time to settle down to its final value.

The company wishes to model the mass, m kg, which is displayed t seconds after a crate X is placed on the scales.

For the displayed mass it is assumed that the rate of change of the quantity $\left(0.5\frac{dm}{dt} + m\right)$ with respect to time is proportional to $(80 - m)$.

- (i) Show that $\frac{d^2m}{dt^2} + 2\frac{dm}{dt} + 2km = 160k$, where k is a real constant. [2]

It is given that the complementary function for the differential equation in part (i) is $e^{\lambda t}(A\cos 2t + B\sin 2t)$, where A and B are arbitrary constants.

- (ii) Show that $k = \frac{5}{2}$ and state the value of the constant λ . [4]

When X is initially placed on the scales the displayed mass is zero and the rate of increase of the displayed mass is 160 kg s^{-1} .

- (iii) Find m in terms of t . [7]

- (iv) Describe the long term behaviour of m . [1]

- (v) With reference to your answer to part (iv), comment on a limitation of the model. [1]

- (vi) (a) Find the value of m that corresponds to the stationary point on the curve $m = f(t)$ with the smallest positive value of t . [2]

- (b) Interpret this value of m in the context of the model. [1]

- (vii) Adapt the differential equation $\frac{d^2m}{dt^2} + 2\frac{dm}{dt} + 5m = 400$ to model the mass displayed t seconds after a crate Y, of mass 100 kg, is placed on the scales. [1]

END OF QUESTION PAPER

BLANK PAGE

BLANK PAGE

Copyright Information:

OCR is committed to seeking permission to reproduce all third-party content that it uses in the assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact OCR, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.