

This document consists of 12 pages

Text Instructions

1. Annotations and abbreviations

Annotation in scoris	Meaning
✓and ×	-
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in	Meaning
mark scheme	
E1	Mark for explaining a regult or establishing a given regult
	Mark for explaining a result or establishing a given result
dep*	Mark dependent on a previous mark, indicated by *
dep*	Mark dependent on a previous mark, indicated by *
dep* cao	Mark dependent on a previous mark, indicated by * Correct answer only
dep* cao oe	Mark dependent on a previous mark, indicated by * Correct answer only Or equivalent Rounded or truncated Seen or implied
dep* cao oe rot	Mark dependent on a previous mark, indicated by * Correct answer only Or equivalent Rounded or truncated
dep* cao oe rot soi	Mark dependent on a previous mark, indicated by * Correct answer only Or equivalent Rounded or truncated Seen or implied Without wrong working Answer given
dep* cao oe rot soi www	Mark dependent on a previous mark, indicated by * Correct answer only Or equivalent Rounded or truncated Seen or implied Without wrong working
dep* cao oe rot soi www AG	Mark dependent on a previous mark, indicated by * Correct answer only Or equivalent Rounded or truncated Seen or implied Without wrong working Answer given

2. Subject-specific Marking Instructions for AS Level Mathematics A

- a Annotations should be used whenever appropriate during your marking. The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded. For subsequent marking you must make it clear how you have arrived at the mark you have awarded.
- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.

 If you are in any doubt whatsoever you should contact your Team Leader.
- c The following types of marks are available.

М

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

Α

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

В

Mark for a correct result or statement independent of Method marks.

Ε

Mark for explaining a result or establishing a given result. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep*' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is **not** the case please, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.

 Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is **not** shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.
- In Unless units are specifically requested, there is no penalty for wrong or missing units as long as the answer is numerically correct and expressed either in SI or in the units of the question. (e.g. lengths will be assumed to be in metres unless in a particular question all the lengths are in km, when this would be assumed to be the unspecified unit.) We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so. When a value is given in the paper only accept an answer correct to at least as many significant figures as the given value. This rule should be applied to each case. When a value is not given in the paper accept any answer that agrees with the correct value to 2 s.f. Follow through should be used so that only one mark is lost for each distinct accuracy error, except for errors due to premature approximation which should be penalised only once in the examination. There is no penalty for using a wrong value for g. E marks will be lost except when results agree to the accuracy required in the question.
- Rules for replaced work: if a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests; if there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others. NB Follow these maths-specific instructions rather than those in the assessor handbook.
- h For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A mark in the question. Marks designated as cao may be awarded as long as there are no other errors. E marks are lost unless, by chance, the given results are established by equivalent working. 'Fresh starts' will not affect an earlier decision about a misread. Note that a miscopy of the candidate's own working is not a misread but an accuracy error.
- If a calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers (provided, of course, that there is nothing in the wording of the question specifying that analytical methods are required). Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.
- j If in any case the scheme operates with considerable unfairness consult your Team Leader.

	Questic	on	Answer	Marks	AOs	Guida	nce
1	(i)	(a)		M1	1.1	A horizontal translation only is seen	
			Coordinates $(-3, 0)$, $(-1, 0)$ and $(-2, 2)$	A1	1.1		
			seen				
				[2]			
1	(i)	(b)	Reflection in the <i>x</i> -axis	B1	1.2	Must be a complete statement	
				[1]			
1	(ii)	(a)	(2, 12)	B1	1.1		
_	(**)	4.	(.)	[1]			
1	(ii)	(b)	$\left(\frac{1}{2},3\right)$	B1	1.1		
				[1]			
2			DR				
			$2(1-\sin^2 x) = 2-\sin x$	M1	3.1a	Use $\cos^2 x = 1 - \sin^2 x$ and simplify	One step of simplification must be seen
			$2\sin^2 x - \sin x = 0$	A1	1.1	Obtain $2\sin^2 x - 1\sin x = 0$	
			$\sin x(2\sin x - 1) = 0$	M1	1.1a	Attempt to solve a 2 term quadratic in	Use any valid method
						sin x and use correct order of	Must be seen
						operations to obtain x	
			$\sin x = \frac{1}{2}$ so $x = 30$ or $x = 150$	A1	1.1	Both values are required	
1			$\sin x = 0 \text{ so } x = 0 \text{ or } x = 180$	A1	1.1	Both values are required	
				[5]			

	Duestic	'n	Answer	Marks	AOs	Guidance	\neg
	_)II 	*****				-
3	(i)		The model is exponential so the rate of	M1	1.1	Gradient of $e^{kx} = ke^{kx}$	
			change of <i>m</i> is proportional to <i>m</i>	17:1	22	T	
			In this case, the rate of change of m is $2m$	E1	2.2a	In context	
	(8.8)			[2]			
3	(ii)		The initial membership	B1	1.1		
				[1]			
3	(iii)		$60000 = 150e^{2t}$	M1	3.4	Correct equation and use correct order	
						of operations	
			ln 400 = 2t	A1	1.1	Obtain correct intermediate step	
						Or $\ln 60000 = \ln 150 + 2t$	
			2.995 = t and hence 3	A1	1.1	Obtain correct answer	
				[3]			
3	(iv)		E.g. When the graph reaches 60 000 the	B1	3.5c	Correct suggestion	
	\ <i>'</i>		graph becomes constant.				
				[1]			
4	(i)		(4)	B1	1.1		
-	(-)		$\overrightarrow{BC} = \begin{pmatrix} 4 \\ -2 \end{pmatrix}$				
			(-2)				
			$\begin{pmatrix} 4 \\ -2 \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} -2 \\ 1 \end{pmatrix} = \mathbf{d} - \mathbf{a} = \overline{AD}$	M1	3.1a	soi	
			$\begin{vmatrix} $				
			$\overrightarrow{OD} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$	A1	1.1		
			$ OD = _{-1}$				
				ray	ļ		
<u> </u>	(**)			[3]			_
4	(ii)		$ \overrightarrow{OM}_{-}(4) $	B1	1.1		
			$OM = \begin{pmatrix} 4 \end{pmatrix}$				
				M1	1.1	soi	
			$ \overrightarrow{AM} = \overrightarrow{OM} - \overrightarrow{OA} = $	IVII	1.1	SOI	
			$\overrightarrow{OM} = \begin{pmatrix} 4\\4 \end{pmatrix}$ $\overrightarrow{AM} = \overrightarrow{OM} - \overrightarrow{OA} = \begin{pmatrix} 6\\3 \end{pmatrix}$ $ \overrightarrow{AM} = \sqrt{6^2 + 3^2} = 3\sqrt{5}$				
				A1	2.2a	Accept 6.71	
			$ AM = \sqrt{6^2 + 3^2} = 3\sqrt{5}$			1100pt 01/1	
1				[3]			

	Question	Answer	Marks	AOs	Guid	ance
5	(i)	$\log_{10} y = \log_{10} p + x \log_{10} q$	B1	2.1		
		$m = \log_{10} q$, $c = \log_{10} p$	B1	2.4		
			[2]			
5	(ii)	E.g. $\log_{10} q = \frac{2.4 - 1.6}{1 - 5} = -0.2$	M1	3.3	Measure gradient from graph and identify it as $\log q$	
		$q = 10^{-0.2} = 0.63$	A1	1.1		Accept q in $[0.6, 0.7]$
		$\log_{10} p = 2.5 \text{ so } p = 380$	B1	1.1		Accept pin [320, 400]
			[3]		4	
5	(iii)	$\log_{10} 20 = 1.3$ so week 7	B1	3.4		
		E.g. Extrapolation is unjustified because it assumes that the assumptions made in the model will hold true in the long term	E1	3.5b	One valid explanation	
6	(i)	31 gives $3^2 + 1^2 = 10$	M1	2.1		OR
		10 is even and hence the suggestion is false	E1	2.1		M1 37 gives $3^2 + 7^2 = 58$ E1 58 is even and hence the suggestion is false
6	(ii)	$n^2 + (n+1)^2 + (n+2)^2$	Mī	2.1	Any valid expressions for three consecutive integers	
		$3n^2+6n+5$	A1FT	1.1	FT their expressions	
		$3(n^2+2n+1)+2$ which always leaves a	E1	2.1	Correct conclusion.	
		remainder of 2 and so cannot be divided by 3	[3]			

	Questio	n	Answer	Marks	AOs	Guidance		
7			$f(x+h) = x^4 + 4x^3h + 6x^2h^2 + 4xh^3 + h^4$	M1	1.1	Attempt at expansion with product of powers of <i>x</i> and <i>h</i> summing to 4 and some attempt at coefficients, not		
			2 2 2 4	3.64		necessarily correct		
			$\frac{f(x+h)-f(x)}{h} = \frac{4x^3h + 6x^2h^2 + 4xh^3 + h^4}{h}$	M1	1.1	Attempt $\frac{f(x+h)-f(x)}{h}$ Allow at most two errors		
			$=4x^3+6x^2h+4xh^2+h^3$	A1	1.1	All terms correct		
			As $h \to 0$ all the terms in h tend to zero. Therefore $f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h} = 4x^3$	A1	2.4	Accept some indication that as h tends to 0, the terms involving h vanish and leave $4x^3$		
			n→0 N	E1	2.1	Award for good use of language, and of limit and function notation	Only requires the two M1 marks to be awarded.	
				[5]				

C	Question	Answer	Marks	AOs	Guida	nce
8		2x + 3y = 0	241	2.1	Identify gradient of line $\left(=-\frac{2}{3}\right)$	Allow sign slip
		$\Rightarrow y = -\frac{2}{3}x$ and gradient $-\frac{2}{3}$	M1	3.1a	anywhere	
		Hence, gradient of the tangent is $\frac{3}{2}$	A1FT	1.1	Use $m_1 m_2 = -1$ anywhere $\left(=\frac{3}{2}\right)$ FT	
			M1	1.1a	their gradient Attempt differentiation	The power must be seen to decrease
		$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{3}{2}kx^{\frac{1}{2}}$	A1	1.1	Obtain $\frac{3}{2}kx^{\frac{1}{2}}$	
		At $x = 4$, $\frac{3}{2}k(4)^{\frac{1}{2}} = 3k$	M1	1.1	Substitute $x = 4$ and equate to the normal gradient	Tangent gradients may also be used i.e. $-\frac{1}{3k} = -\frac{2}{3}$
		Hence $3k = \frac{3}{2}$, so $k = \frac{1}{2}$	E1	1.1	AG	
		At P , $y = \frac{1}{2}(4)^{\frac{3}{2}} = 4$ so $P = (4, 4)$ so	M1	3.1a	Identify coordinates, gradient of	Accept $y = 4$
		equation of normal through P is $(y-4) = -\frac{2}{3}(x-4)$			normal and form equation with their coordinates	
		When $y = 0$, $x = 10$ so $Q = (10, 0)$	A1	1.1	Substitute $y = 0$ and obtain $x = 10$	
		Using $P(4, 4)$ and $Q(10, 0)$ $PQ^2 = (10-4)^2 + (0-4)^2$	M1	1.1	Use Pythagoras to obtain length PQ^2	
		Circle equation is $(x-4)^2 + (y-4)^2 = 52$	A1FT [10]	1.1	Accept equivalent forms FT their coordinates for P and Q	

	Questic	on	Answer	Marks	AOs	Guida	nce
9	(i)		(4-0)	M1	1.1	Attempt at acceleration calculation	Or use of $v = u + at$ with
			$-\left(\frac{4-0}{8-5}\right)$			with at most one error	v = 0, u = 4 and $t = 3$ with at least
							2 values correct
			Acceleration = $-\frac{4}{3}$	A1	1.1	Or equivalent	
				[2]			
9	(ii)		Distance travelled		1.1	Attempt at both areas; the trapezium	
			$=\frac{1}{2}(5+8)(4)+\frac{1}{2}(12)(3)=44$	M1		and triangle	
			<u>2</u> 2				
			Displacement	4.1		Either distance travelled or	
			$=\frac{1}{2}(5+8)(4)-\frac{1}{2}(12)(3)=8$	A1	1.1	displacement correct	
			2 $44 m > 40 m$ so distance travelled is more	E1	2.2a	Must see relevant comparison	
			than five times the displacement	EI	2.2a	Must see relevant comparison	
			than five times the displacement				
				[3]	*		
10	(i)		$s = 9(6)^2 - 1.5(6)^3 = 0$	M1	1.1		
			E.g. The boomerang is at O	E1	3.4		
			E.g. She catches the boomerang			/	
				[2]			
10	(ii)		$v = 18t - \frac{9}{2}t^2$	M1	1.1		
			When $v = 0$,	M1	1.1	Imply deduction that greatest distance	
			$\begin{array}{l} \text{when } v = 0, \\ t = 0 \text{ or } t = 4 \end{array}$	A1	1.1	is when velocity $= 0$ and solve	
			At $t = 0$, $s = 0$, so maximum displacement	E1	3.4	is when versely o und serve	
			must be when $t = 4$ giving $s = 48$ m				
				[4]			
10	(iii)		t = 5	B1	3.1b		
			v = -22.5	B1	3.4		
10	(:-)		10. 04	[2]	11		-
10	(iv)		a = 18 - 9t Acceleration = -27 (m s ⁻²)	M1 A1	1.1 1.1		
				[2]	1.1		

	Questio	n	Answer	Marks	AOs	Guida	nce
11	(i)		Resultant force from the tug boats is positive so it is moving east	E1	2.2a	(600i)	
			There is zero resultant force in the j direction, so it is not moving north or south	E1	2.2a		
				[2]			
11	(ii)		350 + 250 - 200 = 100000a	M1	3.3	Use $F = ma$. Allow sign errors and	
			2			one missing force	
			Obtain 0.004 m s ⁻²	A1	1.1		
11	(***)		,	[2]	2.11		
11	(iii)		$400 = 1.5t + \frac{1}{2}(0.004)t^2$	M1	3.1b	Use $s = ut + \frac{1}{2}at^2$	
			$0.002t^2 + 1.5t - 400 = 0$	A1	1.1	Obtain correct quadratic. Any equivalent form	
				M1	3.4	Use any method to solve their quadratic	Including BC
			Obtain 209 (seconds)	A1	1.1	If negative root given (-958.63088) this must be clearly discarded	Accept better (208.630877) but not 208
			$v^2 = 1.5^2 + 2(0.004)(400)$	M1	3.4	Use $v^2 = u^2 + 2as$ with their a	
						or $v = u + at$ with their a and t	
			Obtain 2.33 (m s ⁻¹)	A1	1.1	Accept better (2.3345235)	
				[6]			

Assessment Objectives (AO) Grid

Question	AO1	AO2	AO3(PS)	AO3(M)	Total
Pure					
1(i)(a)	2				2
1(i)(b)	1				1
1(ii)(a)	1				1
1(ii)(b)	1				1
2	4		1		5
3(i)	1	1			2
3(ii)	1				1
3(iii)	2			1	3
3(iv)				1	1
4(i)	2		1		3
4(ii)	2	1			3
5(i)		2			2
5(ii)	2			1	3 2
5(iii)				2	
6(i)		2			2
6(ii)	1	2			3 5
7	3	2			
8	8		2		10
Mechanics					
9(i)	2				2
9(ii)	2	1			3
10(i)	1			1	2
10(ii)	3		AX	1	4
10(iii)			1	1	2
11(i)		2			2
11(ii)	1			1	2
11(iii)	3		1	2	6
Totals	45	13	6	11	75

PS = Problem Solving M = Modelling





AS Level Mathematics AH230/02 Pure Mathematics and Mechanics

Printed Answer Booklet

Date - Morning/Afternoon

Time allowed: 1 hour 30 minutes

OCR supplied materials:

· Printed Answer Booklet

You must have:

- · Printed Answer Booklet
- · Scientific or graphical calculator



First name										
Last name										
Centre number	Candidate number									

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer all the questions.
- Write your answer to each question in the space provided in the Printed Answer Booklet
- Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by g m s-2. Unless otherwise instructed, when a numerical value is needed, use g = 9.8.

INFORMATION

- You are reminded of the need for clear presentation in your answers.
- The Printed Answer Booklet consists of 12 pages. The Question Paper consists of 8 pages.



Section A: Pure Mathematics

1(i)(a)							y					
						5	<u> </u>					:
			<u>.</u>	<u>:</u> :	<u>.</u>	4					<u></u>	<u> </u>
2											<u></u>	
	1 1 >>,										<u></u>	
		. 5	4	3	2	1 <i>O</i>		1	2	3	4	5
				<u>.</u>	<u>.</u>	-2						
				: :	<u></u>	-3						
	000000			<u>.</u> : : :	 	- 4	• • • • • • • • • • • • • • • • • • • •		<u> </u>	: :		
		:	:	:	:	- 5						
						_	7					
					-6							
					X							
1(i)(b)												
1(ii)(a)												
1(ii)(b)												

2	

3(i)	
3(ii)	
3(iii)	
·	
·	
3(iv)	

4(i)	
4(ii)	
5(i)	

5(ii)	
	<u>^</u>
	3
	2
	$\log_{10} y$
	1
	0
	0 1 2 3 4 5 6 7 8
5(iii)	
_	

6(i)	
6(ii)	
7	

8	

Section B: Mechanics

9(i)	
9(ii)	
10(i)	

10(ii)	
10(iii)	
10(iv)	
11(i)	

11(ii)	
11(11)	
11(iii)	

PLEASE DO NOT WRITE ON THIS PAGE

Copyright Information:

OCR is committed to seeking permission to reproduce all third-party content that it uses in the assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations