



Circum-Arctic Map of Permafrost and Ground Ice Conditions

Summary

The Circum-Arctic permafrost and ground ice map is available via ftp in ESRI Shapefile format and Equal-Area Scalable Earth Grid (EASE-Grid) format. See the [Format](#) section for an explanation of the files provided via ftp.

The circumpolar permafrost and ground ice data contribute to a unified international data set that depicts the distribution and properties of permafrost and ground ice in the Northern Hemisphere (20°N to 90°N). The re-gridded data set shows discontinuous, sporadic, or isolated permafrost boundaries. Permafrost extent is estimated in percent area (90-100%, 50-90%, 10-50%, <10%, and no permafrost). Relative abundance of ground ice in the upper 20 m is estimated in percent volume (>20%, 10-20%, <10%, and 0%). The data set also contains the location of subsea and relict permafrost. The gridded data are gridded at 12.5 km, 25 km, and 0.5 degree resolution. The shapefiles were derived from the original 1:10,000,000 paper map ([Brown et al. 1997](#))

Permafrost, or permanently frozen ground, is ground (soil, sediment, or rock) that remains at or below 0°C for at least two years (Permafrost Subcommittee, 1988). It occurs both on land and beneath offshore arctic continental shelves, and underlies about 22% of the Earth's land surface.

For more information on the creation of the original map, see [Heginbottom et al. \(1993\)](#). The original paper map also includes information on the relative abundance of ice wedges, massive ice bodies and Pingos, ranges of permafrost temperature and thickness ([Brown et al. 1997](#)).

Citing These Data:

Brown, J., O.J. Ferrians, Jr., J.A. Heginbottom, and E.S. Melnikov. 1998, revised February 2001.
Circum-arctic map of permafrost and ground ice conditions. Boulder, CO: National Snow and Ice Data Center/World Data Center for Glaciology. Digital media.

Overview Table

Category	Description
Data format	ESRI Shapefiles and EASE-Grid
Spatial coverage	25N - 90N, 180W-180E
Grid type and size	Lambert Azimuthal projection, EASE-Grid
File naming convention	*.avl = legend *.dbf = attribute data *.prj = projection info *.shp = feature geometry *.shx = feature geometry index *.byte = binary raster files *.hdr = header files used to bring the binary raster files into a GIS program
File size	94 bytes - 20.2 MB
Parameter(s)	discontinuous, sporadic, or isolated permafrost boundaries permafrost extent relative abundance of ground ice in the upper 20 m location of subsea and relict permafrost
Procedures for obtaining data	Data are available via FTP

Table of Contents

- [1. Contacts](#)
- [2. Detailed Data Description](#)
- [3. Data Access and Tools](#)
- [4. Data Acquisition and Processing](#)
- [5. References and Related Publications](#)
- [6. Document Information](#)

1. Contacts:

Investigator(s) Name and Title:

Tingjun Zhang
National Snow and Ice Data Center
CIRES, 449 UCB
University of Colorado
Boulder, CO, USA

Technical Contact:

NSIDC User Services
National Snow and Ice Data Center
CIRES, 449 UCB
University of Colorado
Boulder, CO 80309-0449 USA
phone: +1 303.492.6199
fax: +1 303.492.2468
form: [Contact NSIDC User Services](#)
e-mail: nsidc@nsidc.org

2. Detailed Data Description:

Format:

The Circum-Arctic permafrost and ground ice map data set contains the following files:

permaice.avl = ESRI Shapefile legend file for permaice coverage based on NUM_CODE field in the polygon attribute table
permaice.dbf = attribute data
permaice.prj = projection info
permaice.shp = feature geometry
permaice.shx = feature geometry index
subsea.avl: ESRI Shapefile legend file for subsea coverage
subsea.dbf = attribute data
subsea.prj = projection info
subsea.shp = feature geometry
subsea.shx = feature geometry index
treeline.avl: ESRI Shapefile legend for treeline map
treeline.dbf = attribute data
treeline.prj = projection info
treeline.shp = feature geometry
treeline.shx = feature geometry index
l1ipa.byte = binary raster file, 0.5 degree by 0.5 degree version
l1ipa.hdr = header file used to bring the binary raster files into a GIS program
nh1pa.byte = binary raster file, 2.5 km by 12.5 km EASE-Grid version
nh1pa.hdr = header file
nl1pa.byte = binary raster file, 25 km by 25 km EASE-Grid version
nl1pa.hdr = header file
readpf.f = fortran program

Each datum within a file is a single byte. Numerical data values are listed below.

Value	Definition
0	No information
1 - chf	Continuous permafrost extent with high ground ice content and thick overburden
2 - dhf	Discontinuous permafrost extent with high ground ice content and thick overburden
3 - shf	Sporadic permafrost extent with high ground ice content and thick overburden
4 - ihf	Isolated patches of permafrost extent with high ground ice content and thick overburden
5 - cmf	Continuous permafrost extent with medium ground ice content and thick overburden
6 -	Discontinuous permafrost extent with medium ground ice content and thick overburden

dmf	
7 - smf	Sporadic permafrost extent with medium ground ice content and thick overburden
8 - imf	Isolated patches of permafrost extent with medium ground ice content and thick overburden
9 - clf	Continuous permafrost extent with low ground ice content and thick overburden
10 - dlf	Discontinuous permafrost extent with low ground ice content and thick overburden
11 - slf	Sporadic permafrost extent with low ground ice content and thick overburden
12 - ilf	Isolated patches of permafrost extent with low ground ice content and thick overburden
13 - chr	Continuous permafrost extent with high ground ice content and thin overburden and exposed bedrock
14 - dhr	Discontinuous permafrost extent with high ground ice content and thin overburden and exposed bedrock
15 - shr	Sporadic permafrost extent with high ground ice content and thin overburden and exposed bedrock
16 - ihr	Isolated patches of permafrost extent with high ground ice content and thin overburden and exposed bedrock
17 - clr	Continuous permafrost extent with low ground ice content and thin overburden and exposed bedrock
18 - dlr	Discontinuous permafrost extent with low ground ice content and thin overburden and exposed bedrock
19 - slr	Sporadic permafrost extent with low ground ice content and thin overburden and exposed bedrock
20 - ilr	Isolated patches of permafrost extent with low ground ice content and thin overburden and exposed bedrock
21 - g	Glaciers
22 - r	Relict permafrost
23 - l	Inland lakes
24 - o	Ocean/inland seas
25 - ld	Land

Permafrost extent codes

c = continuous	(90-100%)
d = discontinuous	(50- 90%)
s = sporadic	(10- 50%)
i = isolated patches	(0 - 10%)

Ground ice content codes

h = high	(>20% for "f" landform codes) (>10% for "r" landform codes)
m = medium	(10-20%)
l = low	(0-10%)

Landform (terrain and overburden) codes

f	lowlands, highlands, and intra- and intermontane depressions characterized by thick overburden cover (>5-10m)
r	mountains, highlands ridges, and plateaus characterized by thin overburden cover (>5-10m) and exposed bedrock

Projection:

Projection for the raster (*.byte) files is:


Projection: Lambert Azimuthal
Units: meters
Spheroid: defined
Major Axis: 6371228.00000
Minor Axis: 6371228.000
Parameters:
radius of the sphere of reference: 6371228.00000
longitude of center of projection: 0
latitude of center of projection: 90
false easting (meters): 0.00000
false northing (meters): 0.00000

Projection of the shapefiles is:

Projection: Lambert Azimuthal
Datum: none
Units: meters
Spheroid: defined
Major Axis: 6370997.00000
Minor Axis: 0.00000
Parameters:
radius of the sphere of reference: 6370997.00000
longitude of center of projection: 180
latitude of center of projection: 90
false easting (meters): 0.00000
false northing (meters): 0.00000

Spatial Coverage:**Spatial Coverage Map:**

spatial coverage map: 25°N - 90°N, 180°W-180°E



Legend for EASE-Grid Permafrost and Ground Ice Map

Permafrost Extent (percent of area)	Ground Ice Content (visible ice in the upper 10-20 m of the ground; percent by volume)				
	Lowlands, highlands, and intra-and intermontane depressions characterized by thick overburden cover (>5-10m)			Mountains, highlands, ridges, and plateaus characterized by thin overburden cover (<5-10 m) and exposed bedrock)	
	High (> 20%)	Medium (10-20%)	Low (0-10%)	High to medium (>10%)	Low (0-10%)
Continuous (90-100%)	Ch	Cm	Cl	Ch	Cl
Discontinuous (50-90%)	Dh	Dm	Dl	Dh	Dl
Sporadic (10-50%)	Sh	Sm	Sl	Sh	Sl
Isolated Patches (0-10%)	Ih	Im	Il	Ih	Il

Variations in the extent of permafrost are shown by the different colors; variations in the amount of ground ice are shown by the different intensities of color. Letter codes assist in determining to which basic permafrost and ground ice class any particular unit belongs. Letter codes are defined in the documentation that accompanies the data files.



Ice caps and glaciers

3. Data Access and Tools:

Data Access:

Data are available via [ftp](#).

Related Data Collections:

- [All About EASE-Grid](#)
- [NSIDC'S EASE-Grid Geolocation Tools Description](#)

4. Data Acquisition and Processing:

Processing Steps:

The investigators initialized the Eurasian section of the map at the United Nations Environmental Programme/Global Resource Information Data Base (UNEP/GRID)-Arendal in Norway. The Norwegian Mapping Authority scanned and vectorized the polygon arcs, and georeferenced, corrected, and attributed the vectorized map at UNEP/GRID-Arendal.

The North America and Greenland sections were also initialized at UNEP/GRID-Arendal. The investigators manually digitized permafrost boundaries with an Root Mean Square (RMS) error of approximately 1 mm. Any coastline not coinciding with a permafrost unit was digitized as a generalized line. They digitized glacial boundaries as generalized lines and attributed closed polygons. They updated and corrected North America and Greenland at the Cold Regions Research and Engineering Laboratory (CRREL) in Hanover, New Hampshire, and the United States Geological Survey (USGS) in Woods Hole, Massachusetts. They superimposed digital coastline and glacial boundaries using Arc/Info's Identity command, and corrected and updated attributes. Finally, they joined the data set of North America and Greenland with the compiled Eurasian area.

The investigators refined the full map at CRREL and USGS by appending digital coastlines and country lines to the map region south of 50°N. These data only extend down to 25°N, the limit of Northern Hemisphere permafrost. Finally, they superimposed digital data sets of select deep, large lakes with no underlying permafrost.

Data Source:

Following are input data used in processing the digital map:

- The original 1:10,000,000 paper map "Circum-Arctic map of permafrost and ground-ice conditions" ([Brown et al. 1997](#)).
- **[Environmental Systems Research Institute \(ESRI\)](#):**
 - Coastline arc data north and south of 50°N ("cntry95.shp")
 - Country arc data south of 50°N ("cntry95.shp")
 - Digital Chart of the World data set generalized by UNEP/Grid-Arendal
 - Eurasian lakes polygon data set from the Digital Chart of the World
- **[USGS](#)**: Alaskan deep lakes polygon data set
- **[Canadian Soil Information System](#)**: Canadian deep lakes polygon data set ("Hydro" coverage)
- **[Geological Survey of Canada](#)**: Glacier arc data set

5. References and Related Publications:

Baranov, I. Ya. 1982. "Geocryological map of the USSR" (In Russian). Moscow: Main Department for Geodesy and Cartography (GUGK), scale 1:7,500,000.

Baranov, I. Ya., ed. 1965. "Principles of geocryological zonation of the permafrost region" (In Russian). Moscow: Nauka.

Baranov, I. Ya. 1959. *Geographical distribution of seasonally frozen ground and permafrost*. Ottawa: National Research Council of Canada, Technical Translation No. 1121.

Baranov, I. Ya. 1959. *Geographical distribution of seasonally frozen ground and permafrost*. Ottawa: National Research Council of Canada, Technical Translation.

Baranov, I. Ya., ed. 1956. "Geocryological map of the USSR" (In Russian). Moscow: Main Department for Geodesy and Cartography (GUGK), scale 1:10,000,000.

Baulin., V.V., ed. 1982. "Map of geocryological regions of the West Siberian Plain" (In Russian), USSR Ministry of Geology, VSEGINGEO, scale 1:1,500,000, 4 sheets.

Black, R.F. 1954. Permafrost -- a review. *Bulletin of the Geological Society of America* 65: 839-856.

Bliss, L.C. 1979. Vegetation and revegetation within permafrost terrain in *Proceedings of the Third International Conference on Permafrost*. Ottawa: National Research Council of Canada, Vol. 2, pp. 31-50.

Bostock, H.S. 1970. *Physiographic regions of Canada*. Geological Survey of Canada, Map 1254A, scale 1:5,000,000.

Brodzik, M.J. 1998. The [EASE-Grid, A Versatile Set of Equal-Area Projections and Grids](#). Unpublished report to the National Snow and Ice Data Center. Boulder, CO.

Brown, J., O.J. Ferrians, Jr., J.A. Heginbottom, and E.S. Melnikov, eds. 1997. Circum-Arctic map of permafrost and ground-ice conditions. Washington, DC: U.S. Geological Survey in Cooperation with the Circum-Pacific Council for Energy and Mineral Resources. Circum-Pacific Map Series CP-45, scale 1:10,000,000, 1 sheet.

Brown, R.J.E. 1978. Permafrost in *Hydrological Atlas of Canada*. Ottawa: Department of Fisheries and the Environment, Plate 32, scale 1:10,000,000.

Brown, R.J.E. 1973. Permafrost in *National Atlas of Canada*. Ottawa: Department of Energy, Mines and Resources, Plate 11-12, scale 1:15,000,000.

Brown, R.J.E. 1967. *Permafrost in Canada*. Ottawa: National Research Council of Canada, Publication 9769, and Ottawa: Geological Survey of Canada, Map 1246A, scale 1:7,603,200.

Crawford, C.B., and G.H. Johnston. 1971. Construction on permafrost. *Canadian Geotechnical Journal* 8: 236-251.

Ershov, E.D., ed. 1988, 1989. "Geocryology of the USSR" (In Russian). 5 volumes. Moscow: Nauka.

Ferrians, O.J., Jr. 1965. Permafrost map of Alaska in *Miscellaneous Geologic Investigations*. U.S. Geological Survey, Map I-445, scale 1:2,500,000.

Fotiev, S.M., N.S. Danilova, and N.S. Shevleva. 1978. Zonal and regional characteristics of permafrost in central Siberia in *Permafrost -- The USSR Contribution to the Second International Conference*, National

Academy of Sciences, Washington, D.C., pp. 104-110.

Fotiev, S.M. 1978. Effect of long-term cryometamorphism of earth materials on the formation of ground water in *Proceedings of the Third International Conference on Permafrost*. Ottawa: National Research Council of Canada, Vol. 1, pp. 181-187.

Gorbunov, A.P. 1978. Permafrost investigations in high-mountain regions. *Arctic and Alpine Research* 10: 283-294.

Heginbottom, J.A., and M.A. Dubreuil. 1995. Canada -- Permafrost in *National Atlas of Canada*, 5th edition, Plate 2.1 (MCR 4177), scale 1:7,500,000.

Heginbottom, J.A., and M.A. Dubreuil. 1993. A new permafrost and ground ice map for the National Atlas of Canada in *Proceedings of the Sixth International Conference on Permafrost*, Beijing, Vol. 1, pp. 255-260.

Heginbottom, J.A., J. Brown, E.S. Melnikov, and O.J. Ferrians, Jr. 1993. [Circum-arctic map of permafrost and ground ice conditions](#) in *Proceedings of the Sixth International Conference on Permafrost*, Wushan, Guangzhou, China: South China University Press, Vol. 2: 1132-1136. Revised December 1997. Boulder, CO: National Snow and Ice Data Center/World Data Center for Glaciology.

Heginbottom, J.A., and L.K. Radburn. 1992. *Permafrost and ground ice conditions of Northwestern Canada*. Geological Survey of Canada, Map 1691A, scale 1:1,000,000.

Heginbottom, J.A. 1984. The mapping of permafrost. *Canadian Geographer* XXVIII: 78-83.

Hughes, T. 1973. Glacial permafrost and Pleistocene ice ages in *Permafrost -- the North American Contribution to the Second International Conference*, National Academy of Sciences, Washington, D.C., pp. 213-223.

Hughes, O.L. 1969. *Distribution of open system pingos in central Yukon Territory with respect to glacial limits*. Paper 69-34. Ottawa: Geological Survey of Canada.

International Permafrost Association, Data and Information Working Group. 1998. comp. Circumpolar Active-Layer Permafrost System (CAPS), version 1.0. Boulder, CO: National Snow and Ice Data Center. CD-ROM.

Judge, A.S. 1973. Deep temperature observations in the Canadian north in *Permafrost -- the North American Contribution to the Second International Conference*, National Academy of Sciences, Washington, D.C., pp. 35-40.

Knowles, K. 1992. Points, Pixels, Grids, and Cells: A Mapping and Gridding Primer. Unpublished report to the National Snow and Ice Data Center, Boulder, CO.

Kudryavtsev, V.A., K.A. Kondrat'eva, and A.G. Gavrilov. 1978. "Geocryological map of the USSR" (In Russian). General Permafrost Studies; Materials for the Third International Conference on Permafrost. Novosibirsk: Nauka, scale 1:2,500,000.

Mackay, J.R. and S.R. Dallimore. 1992. Massive ice of the Tuktoyaktuk area, western Arctic coast, Canada. *Canadian Journal of Earth Sciences* 29: 1235-1249.

Mackay, J.R. 1972. The world of underground ice. *Annals of the Association of American Geographers* 62: 1-22.

Melnikov, E.S. 1988. Natural geosystems of the plain cryolithozones in *Proceedings of the Fifth International Conference on Permafrost*. Trondheim, Norway: Tapir Publishers, Vol. 1, pp. 208-212.

Melnikov, P.I. 1966. "Geocryological map, Yakustkoi, A.S.S.R." (In Russian). Moscow: Akademia Nauk SSSR, scale 1:5,000,000.

Nikiforoff, C. 1928. The perpetually frozen subsoil of Siberia. *Soil Science* 26: 61-81.

Permafrost Subcommittee. 1988. Glossary of Permafrost and Related ground ice Terms. National Research Council of Canada. Technical Memorandum. No. 142. p. 156.

Pewe, T.L. 1983. Alpine permafrost in the contiguous United States: a review. *Arctic and Alpine Research* 15: 145-156.

Pewe, T.L. 1982. *Geologic hazards of the Fairbanks area*. Special Report 15. Fairbanks: Alaska Division of Geological and Geophysical Surveys.

Popov, A.I., et al. 1990. "Cryolithological map of North America" (In Russian). Moscow: Faculty of Geography, M.V. Lomonosov University, scale 1:6,000,000, 4 sheets.

Popov, A.I., et al. 1985. "Map of cryolithology of the USSR" (In Russian). Moscow: Faculty of Geography, M.V. Lomonosov University, scale 1:4,000,000, 4 sheets.

Popov, A.I., S.P. Kachurin, and N.A. Grave. 1966. Features of the development of frozen geomorphology in northern Eurasia. in *Proceedings of the Permafrost International Conference*, National Academy of Sciences, Washington, D.C., NRC Publication 1287, pp. 481-487.

Rapp, A., and L. Annersten. 1969. Permafrost and tundra polygons in northern Sweden in *The Periglacial Environment*. Edited by T.L. Pewe. Montreal: McGill-Queen's University Press, pp. 65-91.

Shumskiy, P.A, and B.I. Vtyurin. 1966. Underground ice in *Proceedings of the Permafrost International Conference*, National Academy of Sciences, Washington, D.C. NRC Publication 1287, pp. 108-113.

Shi Yafeng, and Mi Disheng, eds. 1988. "Map of snow, ice and frozen ground in China" (In Chinese). Lanzhou Institute of Glaciology and Geocryology, Academia Sinica, scale 1:4,000,000, 2 sheets.

Sodnom, N., and A.L. Yanshin, eds. 1990. Geocryology and Geocryological zonation in *National Atlas of Mongolia*. Moscow: GUGK, Ulan Bator, Plates 40 and 41, scales 1:4,500,000 and 1:12,000,000.

Tong Boliang, et al. 1982. "Map of permafrost along the Qinghai-Xizang Highway" (In Chinese). Lanzhou Institute of Glaciology and Geocryology, Academia Sinica, scale 1:600,000.

Vtyurin, B.I. 1975. "Ground ice of the USSR" (In Russian). Moscow: Nauka.

Wahrhaftig, C. 1965. *Physiographic divisions of Alaska*. U.S. Geological Survey, Professional Paper 482, 52 pp.

Washburn, A.L. 1980. *Geocryology*. New York: John Wiley and Sons, Halstead Press.

Weidick, A. 1968. Observations on some Holocene glacier fluctuations in West Greenland. *Meddelelser om Grønland* 165(6), 202 pp.

Williams, R.B.G. 1969. Permafrost and temperature conditions in England during the last glacial period in *The Periglacial Environment*. Edited by T.L. Pewe. Montreal: McGill-Queen's University Press.

Zhang, T., R.G. Barry, K. Knowles, J.A. Heginbottom, J. Brown. 1999. Statistics and characteristics of Permafrost and ground ice Distribution in the Northern Hemisphere. *Polar Geography* 23(2):132-154.

Zhang, T., J.A. Heginbottom, R.G. Barry, J. Brown. 2000. Further Statistics on the Distribution of Permafrost and Ground Ice in the Northern Hemisphere. *Polar Geography* 24(2):126-131.

6. Document Information:

Document Creation Date:

February 2001

Document Revision Date:

October 2002

14 December 2005

Document URL:

<http://nsidc.org/data/ggd318.html>