

Note For Noether's Theorem

Ting-Kai Hsu

December 28, 2023

Contents

1	Noether Theorem in Classical Mechanics	1
1.1	Lagrangian Formalism	1

1 Noether Theorem in Classical Mechanics

1.1 Lagrangian Formalism

In Lagrangian formalism, we've defined the action by lagrangian,

$$S = \int_{t_1}^{t_2} L[x(t), \dot{x}(t)] dt \quad (1.1)$$

If action follows the *least action principle*, we would have the lagrangian be associated with the correct *equation of motion* of the system given by eq(1.1).

$$\frac{\partial L}{\partial x_i} - \frac{d}{dt} \frac{\partial L}{\partial \dot{x}_i} = 0 \quad (1.2)$$

Noether's theorem states that if given system has continuous symmetry, then there would be a conserved quantity associated with this symmetry.

What is continuous symmetry? One that leaves the action invariant even when the dynamical equations (equations of motion) are *not* satisfied, we call it **infinitesimal symmetry transformation**. We denote the infinitesimal symmetry transformation as δ_S to separate it from the variation δ .

We're interested in the case when the action doesn't change under a symmetry, and this implies there is special property behind the dynamics of the system.

If we would like to discuss the transformation on time, we should use another parameter,

$$\begin{aligned} t &= t(\tau) \\ q(t(\tau)) &= \mathcal{Q}(\tau) \end{aligned} \quad (1.3)$$

Generally, the transformation would be

$$\begin{aligned} t &\rightarrow t' = \tau'(t') \\ q &\rightarrow q'(t') = \mathcal{Q}(\tau') \end{aligned} \quad (1.4)$$

and the variation of position and time¹,

$$\begin{aligned} \delta_S q &= q'(t) - q(t) = \mathcal{Q}'(\tau) - \mathcal{Q}(\tau) = \delta_S \mathcal{Q} \\ \delta_S t &= t' - t = \tau'(t) - \tau(t) = \delta_S \tau \end{aligned} \quad (1.5)$$

Rewrite the original action,

$$S = \int_{t_1}^{t_2} dt L[q(t), \dot{q}(t)] = \int_{\tau_1}^{\tau_2} d\tau \left(\frac{dt}{d\tau} \right) L[\mathcal{Q}, \frac{d}{d\tau} \mathcal{Q}, t] \quad (1.6)$$

Redefine the new lagrangian,

$$\mathbb{L} \left[\mathcal{Q}, \frac{d}{d\tau} \mathcal{Q}, t, \frac{dt}{d\tau} \right] = \frac{dt}{d\tau} L \quad (1.7)$$

Now consider the infinitesimal symmetry transformation, and the new action would become,

$$S' = \int d\tau \mathbb{L} \left[\mathcal{Q}', \frac{d\mathcal{Q}'}{d\tau}, t', \frac{dt'}{d\tau} \right] + \int d\tau \frac{dK}{d\tau} \quad (1.8)$$

Adding a term of total derivative for general consideration, and we know it wouldn't affect the equation of motion. Therefore we can do the variation of action and lagrangian,

$$\delta S = S' - S = 0 \quad (1.9)$$

The variation of action should vanish because we assume it is symmetry infinitesimal transformation.

¹The parameter should be the same.